

Active ingredient: Chlorpyrifos (0,0-dimethyl 0-(3,5-bis-trichloro-2-pyridyl) phosphorothioate) 0.25%
 Inert ingredients 99.75%
 100.00%

HEAD to TAIL[®]

FLEA AND TICK CONTROL FOR DOGS

KILLS FLEAS FOR 60 DAYS. KILLS TICKS FOR 16 DAYS. LEAVES COAT WITH NATURAL LOOK AND FEEL.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Spray in well ventilated area. **FOR MAXIMUM EFFECTIVENESS IT IS IMPORTANT THAT THE DOG'S ENTIRE COAT BE THOROUGHLY WETTED RIGHT DOWN TO THE SKIN. APPLY AS FOLLOWS:**

1. Apply to clean, dry coat. Shampooing and drying prior to application are suggested. Do not repeat application for at least 30 days.
2. The coat should be thoroughly wetted down to the skin with HEAD TO TAIL. Spray against the lay of the hair, rolling the coat on long-haired dogs. Be sure to wet neck, chest, abdomen and legs thoroughly. Avoid spraying in eyes and genitalia. Do not rinse dog.
3. Allow dog to air dry or blow dry. Do not towel dry as this will tend to remove the product. The coat will air dry quickly, usually in 10 to 15 minutes. When dry any tangles or mats will disappear.
4. When dry, comb or brush dog as usual. The coat will feel natural to the touch with no stiffness or sticky residue.
5. If dog is then only wetted with water (as in swimming), product is still effective for 28 days against fleas. Wetting may prevent satisfactory control of ticks; however, the number of ticks will be substantially reduced during the period. Frequent bathing, if necessary, according to directions.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

HAZARDS TO HUMANS: Spray in a well ventilated area. Harmful if swallowed or inhaled. May cause eye irritation. Avoid contact with eyes. Wash thoroughly. Avoid or minimize contact with food. Wash hands and skin exposed to the product with soap and water after use.

HAZARDS TO ANIMALS: Do not use on cats. Do not use on birds, dogs, or small animals under 6 weeks of age. Avoid spraying in eyes. Do not use on sick or injured animals. Do not use on dogs or cats that are pregnant or nursing. Do not use on dogs or cats that are being treated with other flea and tick products. Do not use on dogs or cats that are being treated with an anthelmintic. Do not use on dogs or cats that are being treated with a flea and tick collar. Do not use on dogs or cats that are being treated with a flea and tick shampoo.

Keep this and all other pesticides out of the hands of children. Use only as directed on the label. See back panel for additional information.

PHYSICAL OR CHEMICAL HAZARDS

Flammable liquid and vapor (n-pentane)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. Do not give anything by mouth to someone who has swallowed this product. If you are having difficulty breathing, call a physician.

IF IN EYES: Flush eyes with water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

STORAGE AND DISPOSAL


STORAGE: Store in original container.

DISPOSAL: Do not reuse or refill this container. Do not pour into drains or on the ground. Do not burn.

Distributed by: Bayer Animal Health, Inc.
 Grand Island, NY 11732-0001

800-843-8888
 609-261-2100
 U.S. Patent No. 4,478,615

Kills fleas for 60 days



Coats hairs with an invisible shield that protects the entire dog


HEAD to TAIL[®]

FLEA & TICK CONTROL FOR DOGS

Coats hairs with an invisible terpolymer that protects the entire dog

DO NOT USE ON CATS
KEEP OUT OF REACH OF CHILDREN
WARNING: See back panel for additional precautionary statements

8 FL. OZ. (237 mL)



Flea and Tick Control for Dogs KILLS FLEAS FOR 60 DAYS. KILLS TICKS FOR 16 DAYS.

TO USE: Remove child resistant cap and replace with pump dispenser. Replace child resistant cap when not using product.

Active Ingredient:

Chlorpyrifos (0,0-Diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate)	0.25%
Inert Ingredients	99.75%
	<hr/> 100.00%

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Spray in well ventilated area.

FOR MAXIMUM EFFECTIVENESS IT IS IMPORTANT THAT THE DOG'S ENTIRE COAT BE THOROUGHLY WETTED RIGHT DOWN TO THE SKIN. APPLY AS FOLLOWS:

1. Apply to clean dry coat. Shampooing and drying prior to application are suggested. Do not repeat application for at least 30 days.
2. The coat should be thoroughly wetted down to the skin with this product. Spray against the lay of the hair, ruffling the coat on long haired dogs. Be sure to wet neck, chest, abdomen and legs thoroughly. Avoid spraying in eyes or on genitalia. Do not rinse dog.
4 mls require about 1.5 fl. oz. (50 squares) per 10 lbs. of body weight to thoroughly wet the coat.
3. Allow dog to air dry or blow dry. Do not towel dry as this will tend to remove the product. The coat will air dry quickly, usually in 10 to 15 minutes. When dry, any residual odor will disappear.
4. When dry, comb or brush dog as usual. The coat will feel natural to the touch with no stiffness or soapy residue.
5. If dog is thoroughly wetted with water (rain/swimming), product is still effective for 28 days against fleas. Wetting may prevent satisfactory control of ticks; however, the number of ticks will be substantially reduced during this period. Re-treat if necessary according to directions.

DO NOT USE ON CATS. KEEP OUT OF REACH OF CHILDREN.
WARNING: See back panel for additional precautionary statements.

8 FL. OZ. (237 ml)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

HAZARDS TO HUMANS: Spray in well ventilated area. Harmful if swallowed or inhaled. May cause eye irritation. Avoid contact with eyes, skin or clothing. Avoid contamination of food. Wash hands and skin exposed to product with soap and water after use.

HAZARDS TO ANIMALS: Do not use on cats. Do not use on nursing dogs or pups under 16 weeks of age. Avoid spraying in eyes. Do not use on sick or debilitated dogs. Chlorpyrifos is a cholinesterase inhibitor. Do not use this product simultaneously or within 30 days after treating with other cholinesterase inhibiting drugs or insecticides. In the event dog exhibits an adverse reaction from use of this polymer spray, it can be readily removed with soap and water.

Note to Physician/Veterinarian: Atropine only by injection is antidotal. Use only if symptoms of cholinesterase inhibition are present.

PHYSICAL OR CHEMICAL HAZARDS

Flammable: keep away from heat or open flame.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

STORAGE AND DISPOSAL

STORAGE: Store at room temperature.

DISPOSAL: Do not reuse empty container. Wrap bottle and put in trash.

EPA Reg. No. 8220-37
EPA Est. 70567X1

Product No. 15221
BT 35221

Distributed by Carter Wallace, Inc.
Granbury, NJ 08512-0187

U.S. Patent No. 4,808,615 © 1986, 1990 Carter Wallace, Inc.

16-02-21

16-02-21

