


BACK PANEL



L. E. M. A.

GENERAL USE PESTICIDE

(Contains 2,4-DB and 2,4-1)

KILLS WORMS, CHICKENS, CLOTHES

KILLS MANY OTHER LIVING BEINGS

KILLS POISON IVY, IRISH-RODENTS

ACTIVE INGREDIENTS:

Diethylamine salt of 2,4-dichlorophenoxyacetic acid* 12.00%

Diethylamine salt of dicamba (3,6-dichloro- α -
anisic acid)** 2.10%

Dimethylamine salt of related acid. 30%

INERT INGREDIENTS: 85.60%

100.00%

*Equivalent to 16.60% of 2,4-dichlorophenoxyacetic acid

**Equivalent to 2.00% of 3,6-Dichloro- α -anisic acid

CAUTION: KEEP OUT OF REACH OF CHILDREN

(See back panel for additional cautions)

PRODUCT NO. 296-
296-

USDA REG. NO. 1222-7

MAVIC[®] is a registered trademark of
Melsicol Chemical Corporation.

Manufactured for TOSI, A COUNTRY ENTERPRISES Co.

GENERAL PACKAGING & CHEMICAL CORP., BURBANK CITY, CALIFORNIA

BEST DOCUMENT AVAILABLE

8222-33

INSTRUCTIONS

FOR CONTROL OF INSECTS AND INJESTIVE PARASITES ON CROPS, FOREST AND INDUSTRIAL PLANTS, HORTICULTURE, AGRICULTURE, GARDENING, LANDSCAPING, FORESTRY, AND ALL FORMS OF CONSERVATION. FOR CONTROL OF INSECTS, SPIDER MITES, FLEA BEETLES, BORER BEETLES, BROWN MOTH, CATERPILLARS, WORMS, LEAFHOPERS, TANDEM FLIES, CHAFERS, GRASSHOPPERS, BIRDS, RODENTS, SQUIRRELS, CHIPMUNKS, FOXES, COONSKINS, BEAVERS, RABBIT, DEER, HOGS, HORSES, SHEEP, GOATS, CATTLE, PIGS, DOGS, CATS, BIRDS, ETC.

FOR INSECTS: Mix 1 part 10% DDT in 1 gallon of water, apply to 200 sq. ft. of area. If kept in 10 qts. of water, apply to 1,400 sq. ft. Repeat if necessary.

FOR BOXTOP WORM OR VINEWORM: Mix 2 parts 10% DDT in 1 gallon of water or 1 pt. in 5 gallons of water.

FOR MOLY LICE OR BIRD FLUZZ: Mix 10% DDT in 1 gallon of water or 1 pt. in 5 gallons of water.

FOR FOXES, BEAVERS, CHIPMUNKS, SQUIRRELS, CHIPMUNKS, FOXES, COONSKINS, BEAVERS, RABBIT, DEER, HOGS, HORSES, SHEEP, GOATS, CATTLE, PIGS, DOGS, CATS, BIRDS, ETC.

FOR INSECTS: Mix 1 part 10% DDT in 1 gallon of water or 1 pt. in 5 gallons of water. Repeat if necessary.

FOR INSECTS ON FRUIT TREES: Mix 10% DDT in 1 gallon of water or 1 pt. in 5 gallons of water. Apply when the new leaves are well developed, but before the flower buds are fully grown. Let leaves dry quickly. Repeat application as necessary.

C A U T I O N: AVOID CONTACT WITH SKIN, EYES AND CLOTHING. IN CASE OF CONTACT FLUSH WITH PLenty OF WATER; IF EYES GET RADIANT IRRITATION, DO NOT CONTAMINATE EYES. USE FOR PROTECTION OR DOMESTIC PURPOSES. IF OLD SPRAY DRIFT TO SUSCEPTIBLE PLANTS, SUCH AS: GARDEN VEGETABLES, FLOWER BEDS, SHRUBBERY AND OTHER ORNAMENTALS. COARSE SPRAYS ARE LESS LIKELY TO DRIFT. DO NOT APPLY ON OR NEAR DESIRABLE TREES AND SHRUBS OR IN LOCATIONS WHERE CHEMICAL MAY BE WASHED OR MOVED INTO CONTACT WITH THEIR ROOTS.

DO NOT USE SAME SPRAYER FOR OTHER PURPOSES UNLESS YOU thorougly clean spray equipment with suitable chemical cleaner. DO NOT STORE NEAR FERTILIZER, HERBICIDE, INSECTICIDE, FUMICIDES.

DO NOT DRINK CONTAINER. WASH THOROUGHLY AFTER WATER AND DETERGENT. DISCARD IN A SAFE PLACE.

DISCLAIMER: THE RECOMMENDATIONS FOR USE OF THIS PRODUCT ARE BASED UPON TESTS CONDUCTED BY THE MANUFACTURER. THE USE OF THIS PRODUCT SHOULD BE AT THE DISCRETION OF THE MANUFACTURER, OR RECOMMENDED, APPROVED OR PUBLISHED, BY ADEAR TO THE AIR, AIR POLLUTION, THE ENVIRONMENT, OR PUBLIC HEALTH, BASED IN ACCORDANCE WITH DIRECTIONS OF THE MANUFACTURER. THE MANUFACTURER ASSUMES NO RESPONSIBILITY, WHETHER FOR INJURY OR DAMAGE RESULTING FROM THE USE OF THIS PRODUCT OR IN COMBINATION WITH OTHER MATERIALS.

BEST DOCUMENT AVAILABLE

8