

FRONT PANEL

T E C

NET CONTENTS

## F R U I T

and

V E G E T A B L E

C O M B I N E D F I C I D E

TOP: Fruits: Grapes, Cane Berries, Strawberries, Melons, Vegetables.

## CONTENTS (%)

1,0-dimethyl-3-(3-methyl-1,2-benzodiazin-3(4H)-yl)methyl-	
Elaeophor® oil extract . . . . .	5.0%
Urgent Methyl Salicylate 92 U.R. . . . .	28.0%
Aromatic Petroleum Derivative Solvent . . . . .	28.0%
Einycone Dichloride. . . . .	28.0%
INERT INGREDIENTS. . . . .	11.0%


SOUTH-WESTERN Reg. by Chemagro Corp., Licensee U. S. Pat. 2,753,119 Canadian Pat. 552,660.

## W A R N I N G: KEEP OUT OF REACH OF CHILDREN.

See Back Panel For Additional Warnings and Antidotes  
Keep Away From Open Flame And Heat.

Product No. \_\_\_\_\_

U.S.D.A. REG. NO. \_\_\_\_\_



## BACK PANEL: READ ENTIRE LABEL. FOLLOW DIRECTIONS AND WARNINGS STRICTLY.

USE 2 tablespoons (1 oz.) per gallon of water for general spraying. For heavy mite infestations repeat spray after 5 days; on this repeat application the dosage can be reduced to 1 oz. in 1½ gallons of water.

WET ALL PART OF THE PLANT THOROUGHLY.

Repeat as needed except for limitations noted on crops below.

CROPS	PESTS	TIME AND LIMITATIONS
<u>APPLES</u>	aphids, mites, thrips, leaf-hoppers, <del>white flies</del> , scale, stink bugs, mealy bugs, scale, crawlers, codling moth, tortrix fruit worms, leaf-rollers.	Do not apply within 15 days of harvest. Make first application after bloom and 2nd two weeks later.
<u>PEARS</u>	aphids, mites, thrips, leaf-hoppers, mealy bugs, tarnished plant bugs, stink bugs, scale, crawlers, <del>white flies</del> , bud moth, tortrix worms, leafrollers, twig borers.	Do not apply within 21 days of harvest. Use not more than 4 times per season.
<u>QUINCE</u>	aphids, mites, thrips, leaf-hoppers, mealy bugs, tarnished plant bugs, stink bugs, scale, crawlers, <del>white flies</del> , bud moth, tortrix worms, leafrollers, twig borers.	Do not apply within 15 days of harvest. Use not more than 8 times per season.
<u>APRICOTS</u>	aphids, mites, thrips, leaf-hoppers, mealy bugs, tarnished plant bugs, stink bugs, scale, crawlers, <del>white flies</del> , bud moth, tortrix worms, leafrollers, twig borers.	Apply immediately after main bloom and 30 days before harvest. Use not over 2 times and not within 25 days of harvest.
<u>PLUMES</u>	leafhoppers, mites, grasshoppers, beetles.	Spray when new growth is 6 inches long and again at 10 inches, repeat shortly after each time but not more than 3 times per season.
<u>NECTARINES</u>	leafhoppers, mites, grasshoppers, beetles.	
<u>CHERRIES</u>	leafhoppers, mites, grasshoppers, beetles.	
<u>PLUMS</u>	leafhoppers, mites, grasshoppers, beetles.	
<u>PRUNES</u>	leafhoppers, mites, grasshoppers, beetles.	
<u>CITRUS</u>	aphids, mites, white flies, scale crawlers, tortrix, worms, tussock moth.	
<u>GRAPES</u>	leafhoppers, mites, grasshoppers, beetles.	

BACK PANEL CONTINUED:

CROPS	PESTS	TIME AND LIMITATIONS
BLACKBERRIES	Aphids, mites, leafhoppers,	Apply as needed, but not within 14 days of harvest.
BOYSENBERRIES	leaf rollers.	
LOGANBERRIES		
RASPBERRIES	Root borers and weevils.	Drench soil at base 7-10 days before bloom.
STRAWBERRIES	Aphids, mites, leaf rollers, 1. 7 tiers, root weevils.	Apply as needed, but not within 5 days of harvest. Drench soil for weevils.
FILBERTS	Aphids, mealybug, Leaf (Chionanthus.) roller, filbert worm	Do not apply within 30 days of harvest.
PECANS	Aphids, mites, webworm, pecan case bearer	Do not apply after husks split.
WALNUTS	Aphids, mites, codling moth, walnut husk fly.	Apply when nuts are size of peas and 6 weeks before harvest (not after husks split).
<b>ON VEGETABLES spray young plants when they have 4 or 5 true leaves. Repeat as needed within listed limitations.</b>		
ARTICHOKES	Plume moth	Not within 30 days of harvest.
BEANS	Aphids, mites, leafhoppers, leafrollers, worms, beetles.	Not within 7 days of harvest on snap beans. On dry beans spray not more than 4 times and not within 30 days of harvest.
BRUSSELS SPROUTS	Aphids, worms, loopers	Do not spray closer to harvest than 7 days Brussels Sprouts;
BROCCOLI	Root Maggot: Use 2 oz. per pt. water & drench soil with 1/2 cup around each transplant	15 days Broccoli & Cauliflower;
CABBAGE		21 days Cabbage.
CAULIFLOWER		
CUCUMBER	Beetles	Not over 3 times per season nor within 6 days of harvest.
LICORIS	Beetles, leafminers	Not within 14 days of harvest.
ONIONS (dry Only)	Thrips	Not within 28 days of harvest.
PEPPERS	Leafminer, flea beetles, European corn borer.	Not over 4 times per season nor within 5 days of harvest.
	Aphids, fleahoppers.	
POTATOES	Leafhoppers, spittle bugs, leafminers, tubeworms, Colorado potato beetles, European corn borer, tarnished plant bugs.	Do not apply within 7 days of harvest.
SPINACH	Aphids, leaf miners, mites.	Not within 14 days of harvest.
TOMATOES	Aphids, flea beetles, worms, leafhoppers, thrips, grasshoppers.	Thorough coverage when fruit is starting to set will reduce the need for later sprays.

**CAUTION:** Do not pasture animals in sprayed areas for 21 days after last application. Do not feed plant residues to animals.  
Do not apply to plants which are not well irrigated or when temperatures exceed 90°F. within 1 hour before or after spraying.

**H A R T H O L E:** May be Fatal if Swallowed, Inhaled or Absorbed Through the Skin. Do not get in eyes or on skin. Wear rubber gloves, protective clothing and goggles. Wash exposed skin with soap and water. If you feel unwell, seek medical attention immediately.

BACK PANEL CONTINUED:

where there may be danger of drift. Vacated areas should not be re-entered for at least two days. Do not store near feed and food products. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a Physician Immediately.

This product is toxic to fish and wildlife. Keep out of any body of water. When empty, wash container thoroughly and destroy. Do not transfer to unlabelled container.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established practice. The Buyer must assume all responsibility, including **injury or damage, resulting from its misuse as such, or in combination with other materials.**

Packed for TOWN & COUNTRY ENTERPRISES  
By GENERAL PACKAGING & CHEMICAL CORP., UNION CITY, CALIFORNIA