


TOWN & COUNTRY

T & C

SNAIL AND SLUG



ATTRACTS
AND KILLS!

12 PAGES PAGE

180

NET WT. 50 LBS

SNAIL & SLUG KILLER

4. MAINTAINING PELLETS

卷之三

1. *Chlorophytum comosum* (L.) Willd. (Fig. 1) (Plants from the Philippines)

THE TWENTIETH

WARNING

WARNING
Very fatal if swallowed, induce vomiting if swallowed. Avoid breathing dust and vapor. Avoid contact with skin, eyes or clothing. If material contacts the eyes get medical attention immediately. Wash hands thoroughly after handling.

Pets and domestic animals may be attracted to this bait. Care should be taken to keep these animals out of treated areas. Avoid contamination of feed, foodstuff's water supply, streams and ponds. Do not apply within 6 days of harvest to any food crop.

This feeding on treated areas may be killed.

Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of dust from areas treated.

PUBLICATIONS RECEIVED

Fording plants can also be treated with a thin line of bait for protection, but care should be exercised to avoid contaminating edible parts.

Night or evening feeding slugs and snails are two of the most common and destructive garden pests. Proper use of F & C SNAIL & SLUG KILLER will control these pests in your flowers, garden and lawn. METALdehyde is the killing agent in this slug bait and is more effective than most other slug killers with its 4% Metaldehyde content.

Pieces equipment and containers and dispose of wastes by burying in non-contaminated soil, away from water supplies. Containers should be disposed by burying with wastes or incinerating (burn out of smoke).

The recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use. It is the customer's responsibility if not used in accordance with directions or established practices, to furnish and assume all responsibility, including injury or damage, resulting from its use in conjunction with such, or in combination with other materials.

1

LEFT SIDE PAGE:

DIRECTIONS FOR USE

Apply 100 lbs. per acre or 12 lbs. per 1000 sq. ft. to control snails and slugs.
Sprinkle soil surface around plants. Do not contaminate edible parts. For use
on:

Cabbage	Cucumbers	Plants
Carrots	Egg Plants	Ferns
Beans	Endive	Gardens
Beets	Garlic	Grasses
Broccoli	Horseradish	Herbs
Brussel Sprouts	Kale	Horseradish
Cabbage	Kohlrabi	Iceberg
Cantaloupes	Leeks	Onions
Carrots	Lettuce	Spinach
Cauliflower	Melons	Squash
Celery	Mustard Greens	Sweet Potatoes
Collards	Okra	Swiss Chard
Corn	Onion	Tomatoes
Green Peas	Parsnips	Turnips
	Peas	Watermelons

STRAWBERRIES: Apply 100 lbs. per acre to control snails & slugs. Do not apply within 6 days of harvest.

The area to be treated should be damp. The method of applying will vary with the conditions; i.e. many small heaps of about 1 tablespoonful can be placed in a flower garden, a large garden may be covered by sprinkling material throughout the garden area, or the material may be spread in a thin line along a border of an adjacent area of infestation.

WARNING: KEEP OUT OF REACH OF CHILDREN.

Manufactured for TOWN & COUNTRY ENTERPRISES BY GENERAL PACKAGING & CHEMICAL CORP.
P. O. Box 825, Union City, California 94587, Area Code (415) 717-6300