



LIQUID SEVIN* SPRAY

FLOWABLE SUSPENSION
Insecticides for FLOWERS,
FRUITS, VEGETABLES
in Home & Garden

ACTIVE INGREDIENTS:

Carbaryl* (1-naphthyl methyl carbamate) 25%

INERT INGREDIENTS: 75%

Total 100%

*SEVIN, T.M. Reg. by Union Carbide.
U.S. Patents No. 2,903,478, 3,009,855

1 PINT

Product No. 14873
U.S.D.A. REG. NO. 8222-47

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS. TO IMPROVE FLOWABILITY FROM CONTAINER strike sharply with the palm of the hand or on a firm surface, then shake.

WHEN: Use at first sign of infestation or on schedules advised by State Extension Service Specialists. Repeat as necessary. On food crops do not use closer to harvest than shown in the chart below under "DAYS" before harvest and never closer than 30 days apart during the last month. On fruit trees for control of Codling Moth, Oriental Fruit Moth and Plum Curculio Apply at the end of plum blooming period (petal fall) and repeat at 10-14 day intervals until control is achieved.

HOW MUCH: Use 4 teaspoonfuls per gallon (2 oz. to 3 gal.) except as noted in chart. Never use more on tender foliage Use 2 tablespoonfuls (1 oz.) per gallon on extreme infestations.


CAUTION: SEVIN* may cause foliage injury when applied when plants are wet with rain or dew. SEVIN* does not control red spider mites. In areas and on plants where these pests are prevalent, combine or alternate with a good red spider control. DO NOT combine oil sprays on any plants except citrus or olives. Do not spray when temperatures exceed 90° F. within 3 hours before or after

CROP	Days Before Harvest	PESTS
FLOWERS, ORNAMENTALS		Aphid, Leafhopper, exposed Thrips, Young Scale Crawlers, Mealy Bug, Flea Beetle, Cucumber Beetle, Leaf Roller, Bud Moth, tarnished Plant Bug: Use 2 tablespoonfuls per gallon water.
LAWNS		Fall Armyworm, Sod Webworm, Ants, Fleas, Chinch Bug, Mosquitoes: Use 1 cup (8 oz.) to 8 gallons for each 500 square feet.
IN AND AROUND DOG, CAT AND CHICKEN QUARTERS		Fleas, Ticks, and Chicken Mites in housing quarters and sleeping areas: Use 3 oz. per gallon of water on each 500 square feet.
DEBRIS ACCUMULATION AROUND WALKWAYS AND FENCES		Ants, Cockroaches, Earwigs, Millipedes: Use 5 oz. per gallon of water
BEANS		Cucumber, Flea, Japanese and Mexican Bean Beetles, Cutworms: Use 2 tablespoonfuls per gallon of water.
CORN		Corn Ear Worm, Fall Armyworm, Loopers, Sap Beetles, European Corn Borer, Japanese Beetle: Use 2 tablespoonfuls per gallon of water. For corn Earworm: Apply to silks as soon as they appear, make 3-4 applications at 2-3 day intervals Reapply immediately after rain.
CUCUMBERS, MELON, SQUASH		Cucumber & Flea Beetles, Leafhoppers, Pickle Worms, Squash Bugs.
TOMATO, POTATO, PEPPER, EGGPLANT		Flea Beetle, Leafhopper, Lace Bug, Colorado Potato Beetle, Fall Armyworm, Stink Bugs and Tomato Fruit and Horn Worms: Use 1 oz. per gallon of water.
BROCCOLI, CABBAGE, BEETS	3	Armyworms, Cabbage Worm, young Cabbage Looper Larvae: Use 1 oz. per gal. water
LETTUCE, SPINACH	14	Small Armyworm, Leafhopper, Plant Bugs.
CITRUS	5	Cutworms, Leafroller, Tussock Moth, young Scale Crawlers.
CHERRY, PEACH, PLUM, PRUNE	1	Twig Borer, Leaf Roller, Cucumber Beetle, Scale Crawlers, Tarnished Plant Bug, Tent Caterpillar, Japanese Beetle, Codling Moth, Oriental Fruit Moth and Plum Curculio.
APRICOT, NECTARINE	3	
BLACKBERRY, BOYSENBERRY, RASPBERRY	7	Leaf Roller, Leaf Hopper, Raspberry Aphid, Japanese Beetles.
STRAWBERRY	1	Strawberry Leafroller and Weevil Meadow Spittlebug, Japanese Beetle.
GRAPE		Grape Leaf Folder, Leaf Hopper, Grape Berry Moth: Use 1 oz. per gal. of water
ALMOND, FILBERT, WALNUTS (before hulls split)		Scale Crawlers, Almond Twig Borer, Filbert Moth & Aphid, Codling Moth, Leaf Roller and (in California Walnut Aphid).
OLIVES		Scale Crawlers. Combine 2 tablespoonfuls SUMMER OIL per gallon of water.
APPLES, PEARS	1	Aphid, Bag Worm, Codling Moth, Apple Maggot, Leaf Roller, Pear Psylla, young Scale Crawlers. Apples sprayed within 4 weeks after bloom may result in fruit thinning.

CAUTION Harmful if inhaled or swallowed. Avoid prolonged or repeated breathing of spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Avoid contact with the eyes. Do not contaminate food, feed, water supplies, streams or ponds. This product is highly toxic to BEES exposed to direct treatment or residues on crops. Protective information may be obtained from your State Agricultural Experiment Station. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effect of such or the results to be obtained if not used in accordance with directions or established practice. The Buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

Packed for TOWN & COUNTRY ENTERPRISES,
Union City, California
By General Packaging & Chemical Corp.



ACCEPTED WITH COMMENTS