# Kennel-Septic

# **DISINFECTANT• DEODORIZER• SANITIZER**

# **ACTIVE INGREDIENTS:**

n-Alkyl (60%C1₄, 30%C16, 5%C12, 5%C18)	
dimethyl benzyl ammonium chlorides	. 5%
n-Alkyl (68%C <sub>12</sub> , 32%C <sub>14</sub> ) dimethyl	
ethylbenzyl ammonium chlorides	. 5%
Essential Oils	. 2%
INERT INGREDIENTS:	. 88%
	100%
	/0

Kennel-Septic is a concentrate and should be diluted before using.

# WARNING

Keep Out of Reach of Children. See other precautionary statements on other panel.

# NET CONTENTS: 1 GAL.

# ACCEPTED

# JUN 21972

UNDER THE PEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ED UNDER NOTE SUBJECT TO ATTACHED COMMENTS.

# USE DIRECTIONS:

### Household:

For sanitizing and disinfecting floors, walls and inanimate hard surfaces after prior cleaning to remove gross filth and dirt. Apply solution with mop or cloth. Sanitize with 1 ounce Kennel-Septic to 4 gallons of water. Disinfect with 3 ounces Kennel-Septic to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis.

### Kennels, Hospitals, Homes:

For disinfecting floors, walls, and other hard surfaces after prior cleaning to remove gross filth and dirt. In Homes, Kennels and Hospitals, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use Kennel-Septic at  $3\frac{1}{2}$  ounces per 5 gallons of water. At this level Kennel-Septic will be effective against Pseudomonas aeruginosa PRD-10.

### **Food Processing Equipment:**

For sanitization of food processing equipment and food utensils after prior cleaning and rinsing, dilute 1 ounce Kennel-Septic per 4 gallons of water to provide 200 ppm of active quaternary. At this level, No Potable Water Rinse is required.

### WARNING

Keep Out of Reach of Children. Do not get in eyes or on skin or clothing. Causes severe eye irritation. Causes skin irritation. In case of contact, wash immediately with plenty of water. For eyes get medical attention. Harmful if swallowed.

### FIRST AID

If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.


Rinse empty container thoroughly with water and discard.

## LAMBERT-KAY

Div. of Carter-Wallace, Inc. 3628 Crenshaw Boulevard Los Angeles, CA 90016

EPA Reg. No. 8220-18

226-1G-01



4 60%, C <sub>16</sub> 30%,	
8 5%) dimethyl	
onium chlorides	- 1.6%
12 68%, C <sub>14</sub> 32%)	
thylbenzyl ammo-	
ides	- 1.6%
oonate	- 2.0%
DIENTS	- 94.8%
dium nitrite	100.0%

### PHENOL COEFFICIENT (A.O.A.C.)Salmonella typhosa - 18 S. Aureus - 28 USE DILUTION CONFORM. (A.O.A.C.)Staphylococcus aureus 1:64 Salmonella choleraesuis 1:64 Trichophyton

### DIRECTIONS FOR USE

1:64

.

interdigitale

For routine cleaning and classific from swab, wipe or rinse surfaces (such as floors, walls, washstands, pails and bathroom fixtures) thoroughly using a 1:64 (2 oz./gal.) solution of BARCROBE. Disinfect small articles by immersion for at least 10 min. Allow surfaces to dry before use. For heavy cleaning and the cold disinfection of instruments: swab, wipe, rinse or immerse items in a 1:32 (4 oz./gal.) solution of BARCROBE. Remove excessive soil by a separate cleaning and allow a minimum contact time of 10 min. for disinfecting. For dismike finglasses and silverware: Immerse these items in a 1:64 (2 oz./gal.) solution of BARCROBE. operation with Follow a potable water rinse.

For cleaning and discritectingfood and beverage plant processing equipment: Immerse, swab, wipe or rinse these items in a 1:64 (2 oz./ gal.) solution of BARCROBE. Follow the cleaning and sanitizing operation with a potable water rinse. When disinfecting plastic or rubber surfaces, swab if possible. Limit exposure to the 10 min. period.