

BIOCIDE-T

AN ORGANO-METALLIC COMPLEX

ACCEPTED
JUN 30 1978
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 8249-2

WARNING

KEEP OUT OF REACH OF CHILDREN

Causes skin irritation. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. May be absorbed through skin. In case of contact, wash at once with plenty of soap and warm water. Remove contacted clothing. Wash clothing before reuse. If irritation persists, get medical attention. Do not take internally. Biocide T is not a food-stuff and should not come in contact with food. Avoid storage near food and feed products. FIRST AID: In case of swallowing or continued irritation after contact with skin, call a physician.

FISH AND/OR WILDLIFE: This product is toxic to fish. Do not discharge treated effluent where it will drain into lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of waste. Use only as directed on label.

CONTAINER DISPOSAL: Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

CAN

Net Weight.....40 lbs.
Net Volume.....4 gal., 3 qt., 1 oz.

DRUM

Net Weight.....440 lbs.
Net Volume.....52 gal., 1 qt., 13 oz.

INGREDIENTS

- a. ACTIVE INGREDIENTS: N-alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈) dimethyl benzyl ammonium chlorides..... 2.5%
Bis(tri-n-butyltin) oxide.... 0.5%
- b. INERT INGREDIENTS: including solubilizing and dispersing agents..... 97%

KEEP OUT OF REACH OF CHILDREN

WARNING

Causes skin and eye irritation. Harmful if swallowed or absorbed through skin. Avoid breathing spray mist. (See additional precautions and First-Aid information on left side panel)

This product contains 0.25 lbs. of active ingredients per gallon.

EPA REGISTRATION NO. 8249-2
EPA ESTABLISHMENT NO. 8249-TX-1

Net Contents: _____

DIRECTIONS FOR USE OF BIOCIDE T

Always add Biocide T at a point in the system when it will be uniformly mixed throughout the system.

INITIAL DOSE: To a tower visibly fouled, add 6.0 lbs. or 0.75 gal. per 1,000 gallons water in the system. (Shock treatment). Repeat treatment daily until biological control is achieved. Once growths are controlled, go on a continuous feed of Intermittent Shot Feed as a maintenance treatment.

MAINTENANCE DOSE:

1. Continuous Feed (generally for open systems): 0.06 gal. to 0.12 gal. or 0.5 to 1.0 lbs. per 1,000 gallons of make-up to maintain about 60-120 ppm in the system.
2. Intermittent Shot Feed: 0.06 gal. to 0.12 gal. or 0.5 to 1.0 lbs. per 1,000 gallons water in the system every 3 or 4 days or as needed to maintain control.

ALL QUANTITIES REFER TO BIOCIDE T - When Biocide T is used as directed, it will control the growth of algae and bacterial slime commonly found in cooling towers.

MECO

MECO

8249-2

06/30/78

MAINTENANCE ENGINEERING CORP.

P. O. BOX 1729

HOUSTON, TEXAS 77001

PHONE: 222-2351

CHEMICALS WITH ENGINEERING SERVICE