LORMEX MO. U. S. PAT. OFF.

ROOTING POWDER No. 8

Net weight

For rooting difficult to propagate plants such as evergreens and dormant leafless cuttings.

TYPICAL PLANTS ROOTED WITH THIS ROOTING POWDER

Azalea arborescens grandiflora, Azalea canescens, Azalea Roseum, Grape, Juniper vars., Lilac, Maple, Privet, Rhododendron vars.

part part of a state of

INERT INGREDIENT

Indole 3 Butyric Acid

0.8% 99.2%

DIRECTIONS:

- 1. THERE SHOULD BE AS LITTLE TIME AS POSSIBLE BETWEEN THE TIME OF MAKING CUTTINGS, TREATING WITH ROOTING POWDER AND PLANTING.
- 2. CUTTINGS SHOULD BE MOISTENED BEFORE TREATMENT AND THEN EXCESS MOISTURE SHAKEN OFF.
- 3. DIP BASAL ENDS OF CUTTINGS INTO HORMEX ROOTING POWDER.
- 4. SHAKE OFF EXCESS POWDER BY TAPPING CUTTINGS.
- 5. PLANT TREATED CUTTINGS IN ROOTING MEDIA BEING CAREFUL NOT TO RUB OFF ROOTING POWDER WHEN PLANTING.

INHIBITION OF ROOT FORMATION IS POSSIBLE BY THE USE OF TOO HIGH A CONCENTRATION OF ROOTING POWDER. IT IS ESSENTIAL FOR SUCCESSFUL PROPAGATION THAT THE PROPER STRENGTH ROOTING POWDER FOR THE SPECIES INVOLVED BE CHOSEN.

SINCE THIS MATERIAL IS USED UNDER VARYING CONDITIONS BEYOND OUR CONTROL NO WARRANTY OF EFFECTIVENESS IS MADE OR IMPLIED, NOR IS RESPONSIBILITY ASSUMED FOR ANY DETERIOUS RESULTS WHICH MAY BE OCCASIONED BY ITS USE.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Manufactured in U.S.A. by Brooker Chemical

P.O. Box 9335, North Hollywood, California 91606

USDA Reg. No. 8281-1

MC

Aza**u** Rhodi

DIRE

1,

3 6

3

4.


INHIE

ESSE1 CHOS

SINCE OR IM

CA

USD



Net weight

For rooting difficult to propagate plants such as evergreens and dormant leafless cuttings.

TYPICAL PLANTS ROOTED WITH THIS ROOTING POWDER

Azalea arborescens grandiflora, Azalea canescens, Azalea Roseum, Grape, Juniper vars., Lilac, Maple, Privet, Rhododendron vars.

> ACTIVE INGREDIENT INERT INGREDIENT

Indole 3 Butyric Acid

0.8% 99. 2%

DIRECTIONS:

- 1. THERE SHOULD BE AS LITTLE TIME AS POSSIBLE BETWEEN THETIME OF MAKING CUTTINGS, TREATING WITH ROOTING POWDER AND PLANTING.
- 2. CUTTINGS SHOULD BE MOISTENED BEFORE TREATMENT AND THEN EXCESS MOISTURE SHAKEN OFF.
- 3. DIP BASAL ENDS OF CUTTINGS INTO HORMEX ROOTING POWDER.
- 4. SHAKE OFF EXCESS POWDER BY TAPPING CUTTINGS.
- 5. PLANT TREATED CUTTINGS IN ROOTING MEDIA BEING CAREFUL NOT TO RUB OFF ROOTING POWDER WHEN PLANTING.

INHIBITION IF ROOT FORMATION IS POSSIBLE BY THE USE OF TOO HIGH A CONCENTRATION OF ROOTING POWDER. IT IS ESSENTIAL FOR SUCCESSFUL FROPAGATION THAT THE PROPER STRENGTH ROOTING POWDER FOR THE SPECIES INVOLVED BE CHOSEN.

SINCE THIS MATERIAL IS USED UNDER VARYING CONDITIONS BEYOND OUR CONTROL NO WARRANTY OF EFFECTIVENESS IS MADE OR IMPLIED, NOR IS RESPONSIBILITY ASSUMED FOR ANY DETERIOUS RESULTS WHICH MAY BE OCCASIONED BY ITS USE.

CAUTION

DE

KEEP OUT OF REACH OF CHILDREN

Manufactured in U.S.A. by Brooker Chemical

USDA Reg. No. 8281-1

P.O. Box 9335, North Hollywood, California 91606