## WRMEX ROOTING POWDER No. 16

Net weight

For rooting very difficult to propagate plants such as Juniperus torulosa and Thuja pryamidalis Sice

ACTIVE INGREDIENT INERT INGREDIENT

Indole 3 Butyric Acid 1.6%

98.4%

## DIRECTIONS:

- 1. THERE SHOULD BE AS LITTLE TIME AS POSSIBLE BETWEEN THE TIME OF MAKING CUTTINGS, TREATING WITH ROOTING POWDER AND PLANTING.
- 2. CUTTINGS SHOULD BE MOISTENED BEFORE TREATMENT AND THEN EXCESS MOISTURE SHAKEN OFF.
- 3. DIP BASAL ENDS OF CUTTINGS INTO HORMEX ROOTING POWDER.
- SHAKE OFF EXCESS POWDER BY TAPPING OUTTINGS.
- PLANT TREATED CUTTINGS IN ROOTING MEDIA BEING CAREFUL NOT TO RUB OFF ROOTING POWDER WHEN PLANTING.

THIS ROOTING POWDER IS VERY HIGH IN CONCENTRATION OF INDOLE 3 BUTYRIC ACID AND EXTREME CARE SHOULD BE OBSERVED IN ITS USE. IT HAS BEEN USED SUCCESSFULLY ON THE VARIETIES LISTED ABOVE. ITS USE ON OTHER VARIETIES OR DIFFERENT SPECIES SHOULD BE DONE WITH CARE AND ONLY ON AN EXPERIMENTAL BASIS UNTIL ITS EFFECT HAS BEEN PROVEN.

INHIBITION OF ROOT FORMATION IS POSSIBLE BY THE USE OF TOO HIGH A CONCENTRATION OF ROOTING POWDER. IT IS ESSENTIAL FOR SUCCESSFUL PROPAGATION THAT THE PROPER STRENGTH ROOTING POWDER FOR THE SPECIES INVOLVED BE CHOSEN.

SINCE THIS MATERIAL IS USED UNDER VARYING CONDITIONS BEYOND OUR CONTROL NO WARRANTY OF EFFECTIVENESS IS MADE OR IMPLIED, NOR IS RESPONSIBILITY ASSUMED FOR ANY DETERIOUS RESULTS WHICH MAY BE OCCASIONED BY ITS USE.

CAUTION KEEP OUT OF REACH OF CHILDREN

USDA Reg.No. 8281-2

Manufactured in U.S.A. by Brooker Chemical P.O. Box 9335, North Hollywood, California 91606

ACCEPTED I TO ATTACHED COMMENTS