

HORMEX
REG. U. S. PAT. OFF.

ROOTING POWDER No. 3

For rooting woody and semi-woody plants and easy to propagate evergreens.

TYPICAL PLANTS ROOTED WITH THIS ROOTING POWDER

*Arbor-vitae, Azalea dauricum, Azalea Mollis, Blueberry, Daphne, Elder, Ivy, Pyracantha,
Juniperus chinensis pfitzeriana, Juniperus sabina fastigiata, Sequoia gigantea.*


ACTIVE INGREDIENT

Indole 3 Butyric Acid

0.3%

INERT INGREDIENT

99.7%



DIRECTIONS:

1. THERE SHOULD BE AS LITTLE TIME AS POSSIBLE BETWEEN THE TIME OF MAKING CUTTINGS, TREATING WITH ROOTING POWDER AND PLANTING.
- 2A. DO NOT WET HAIRY OR PITHY CUTTINGS AS TOO MUCH ROOTING POWDER WILL ADHERE.
- 2B. CUTTINGS SHOULD BE MOISTENED BEFORE TREATMENT AND THEN EXCESS MOISTURE SHAKEN OFF.
3. DIP BASAL ENDS OF CUTTINGS INTO HORMEX ROOTING POWDER.
4. SHAKE OFF EXCESS POWDER BY TAPPING CUTTINGS.
5. PLANT TREATED CUTTINGS IN ROOTING MEDIA BEING CAREFUL NOT TO RUB OFF ROOTING POWDER WHEN PLANTING.

INHIBITION OF ROOT FORMATION IS POSSIBLE BY THE USE OF TOO HIGH A CONCENTRATION OF ROOTING POWDER. IT IS ESSENTIAL FOR SUCCESSFUL PROPAGATION THAT THE PROPER STRENGTH ROOTING POWDER FOR THE SPECIES INVOLVED BE CHOSEN.

SINCE THIS MATERIAL IS USED UNDER VARYING CONDITIONS BEYOND OUR CONTROL NO WARRANTY OF EFFECTIVENESS IS MADE OR IMPLIED, NOR IS RESPONSIBILITY ASSUMED FOR ANY DETERIOUS RESULTS WHICH MAY BE OCCASIONED BY ITS USE.

CAUTION KEEP OUT OF REACH OF CHILDREN

Manufactured in U.S.A. by Brooker Chemical
P.O. Box 9335, North Hollywood, California 91606

USDA Reg. No. 8281-3

ACCEPTED WITH COMMENTS

HORMEX
REG. U. S. PAT. OFF.

ROOTING POWDER No. 3

Net weight

For rooting woody and semi-woody plants and easy to propagate evergreens.

TYPICAL PLANTS ROOTED WITH THIS ROOTING POWDER

Arbor-vitae, Azalea dauricum, Azalea Mollis, Blueberry, Daphne, Elder, Ivy, Pyracantha, Juniperus chinensis pfitzeriana, Juniperus sabina fastigiata, Sequoia gigantea.

ACTIVE INGREDIENT	Indole 3 Butyric Acid	0.3%
INERT INGREDIENT		99.7%

DIRECTIONS:

1. THERE SHOULD BE AS LITTLE TIME AS POSSIBLE BETWEEN THE TIME OF MAKING CUTTINGS, TREATING WITH ROOTING POWDER AND PLANTING.
- 2A. DO NOT WET HAIRY OR PITHY CUTTINGS AS TOO MUCH ROOTING POWDER WILL ADHERE.
- 2B. CUTTINGS SHOULD BE MOISTENED BEFORE TREATMENT AND THEN EXCESS MOISTURE SHAKEN OFF.
3. DIP BASAL ENDS OF CUTTINGS INTO HORMEX ROOTING POWDER.
4. SHAKE OFF EXCESS POWDER BY TAPPING CUTTINGS.
5. PLANT TREATED CUTTINGS IN ROOTING MEDIA BEING CAREFUL NOT TO RUB OFF ROOTING POWDER WHEN PLANTING.

INHIBITION OF ROOT FORMATION IS POSSIBLE BY THE USE OF TOO HIGH A CONCENTRATION OF ROOTING POWDER. IT IS ESSENTIAL FOR SUCCESSFUL PROPAGATION THAT THE PROPER STRENGTH ROOTING POWDER FOR THE SPECIES INVOLVED BE CHOSEN.

SINCE THIS MATERIAL IS USED UNDER VARYING CONDITIONS BEYOND OUR CONTROL NO WARRANTY OF EFFECTIVENESS IS MADE OR IMPLIED, NOR IS RESPONSIBILITY ASSUMED FOR ANY DETERIOUS RESULTS WHICH MAY BE OCCASIONED BY ITS USE.

CAUTION KEEP OUT OF REACH OF CHILDREN

USDA Reg. No. 8281-3

Manufactured in U.S.A. by Brooker Chemical
P.O. Box 9335, North Hollywood, California 91606

ACCEPTED WITH COMMENTS