191

THIS IS AN EXACT COPY OF THE SILK SCREENED LABEL ON MAR-V-CIDE

IAR-V-CIDE

MAR-V-CIDE DISINFECTANT AND GERMICIDE

E.P.A. Reg. No. 08296 1 E.P.A. Est. No. C8296-MN-001 DANGER

(See side panel)

ONE GALLON (128 FL. OUNCES) 3.785 LITER

ONE-HALF GALLON (64 FL. OUNCES) 1.893 Liter

1 PINT (16 FL. OZ.) (.473 Liters)

DIRECTIONS

DISINFECTION:

Mix l tablespoon Mar-V-Cide in one quart of water; or make stock solution by mixing 2 ounces ($\frac{1}{4}$ cup) Mar-V-Cide in one gallon of water.

(Makes solution containing 1600 ppm active quaternary).

Completely immerse cleaned brushes, combs, razors, clipper heads, scissors, manicure instruments, etc. for 10 minutes (or as required by local health authorities). Change Mar-V-Cide solution every two weeks.

COLD DISINFECTION OF SURGICAL INSTRUMENTS: Clean thoroughly, rinse, and immerse in a solution of one ounce of Mar-V-Cide to one gallon of water for ten minutes.

FOR PROFESSIONAL USE ONLY

WILLIAM MARVY CO. 1540 St. Clair Ave. St. Paul, MN 55105 ACTIVE INGREDIENTS: n-alkyl (C12-40%, C14-50%,

C16-10% dimethyl benzyl ammonium chlorides 10%

INERT INGREDIENTS: 90% Phenol Coefficients: 50 (Salmonella typhosa) 75 (Staphylococcus aureus) (Std. A.O.A.C. Method).

Non-Flammable, non-volatile, odorless, noncorrosive in use dilution when used on metalic items.

DANGER

Keep out of reach of children.
Mar-V-Cide is a concentrated disinfectant. Do
not take internally. Avoid contamination of
food. Avoid contact with skin and eyes; in case
of contact, immediately flush thoroughly with
plenty of water for at least 15 minutes, and get
medical attention.

Do not mix with soaps or synthetic detergents, or other disinfectants.

Rinse empty container thoroughly with water before discarding.

FOR PROFESSIONAL USE ONLY

WILLIAM MARVY CO.

1540 St. Clair Ave., St. Paul, MN 55105

Hot Reviewed. Registrant claims to be in accordance with acregistration of Pesticide Product Guidance for Phase 2. Response, Page 2.12, Active to Inert Change in Status.