

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JAN 1 6 2009

Ms. Karen J. Larson Clarke Mosquito Control 110 E. Irving Park Rd Roselle, IL 60172

Subject:

Natular XRT, EPA Reg. No. 8329-85

Date of Registrant Submission: 10/23/08

Decision: 401983

Dear Ms. Larson:

The labeling referred to above, submitted in connection with registration under the Federal insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records. If you have any questions regarding this letter, please contact Samantha Hulkower at (703) 603-0683.

Sincerely,

Kimberly Nesci Product Manager 11

Insecticide Branch

Registration Division (7505P)

Enclosure: Copy of I

Copy of Label Stamped "Accepted"

JAN 16 2009

PRP 011609 / 8329-85

To be used in governmental mosquito control programs, by professional pest control operators, or in other mosquito or midge control operations.

Group INSECTICIDE

Active Ingredient: spinosad (a mixture of spinosyn A and spinosyn D)*

8.33%

Other ingredients Total

91.67% 100.00%

U.S. Patent No. 5,362,634 and 5,496,931 A Naturalyte® Insect Control product

Natular T30 is an 8.33% tablet, This product may absorb moisture; therefore, the weight of the tablet and percent by weight of active ingredient will vary with hydration.

Keep Out of Reach of Children **CAUTION**

EPA Reg. No. 8329-85

EPA Est.

Manufactured for Clarke Mosquito Control Products, Inc. 159 North Garden Avenue Roselle, IL 60172

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. Wear protective eyewear (such as goggles, face shield, or safety glasses).

	First Aid
If swallowed:	 Call a poison control center of doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.
If in eyes:	Hold eye open and rinse slowly and gently with warm water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
Have the produ	act container or label with you when calling a poison

control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is toxic to aquatic organisms. Non-target aquatic invertebrates may be killed in waters where this pesticide is used. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Read all Directions for Use carefully before applying.

General Information

Natular T30 is a Naturalyte® insect product for killing mosquito and midge larvae. This product's active ingredient, spinosad, is biologically derived from the fermentation of Saccharopolyspora spinosa, a naturally occurring soil organism. Natular T30 releases effective levels of spinosad for up to 30 days under typical environmental conditions.

Release of spinosad is affected by the dissolution of the Natular T30 tablet. If tablets become covered by obstructions such as debris, vegetation, or loose sediment as a result of high rainfall or flow, normal dispersion of the active ingredient can be inhibited. Water flow may increase the dissolution of the tablet, thus reducing the residual life of the tablet. Inspect areas of water flow to determine appropriate re-treatment intervals. To assure positive results, place Natular T30 tablets where they will not be swept away by flushing

General Use Precautions

Integrated Pest Management (IPM) Programs

Natular T30 is intended to kill mosquito and midge larvae.

Mosquitoes are best controlled when an IPM program is followed. Larval control efforts should be managed through habitat mapping, active adult and larval surveillance, and integrated with other control strategies such as source reduction, public education programs, harborage or barrier adult mosquito control applications, and targeted adulticide applications..

Insecticide Resistance Management (IRM)

contains Natular T30 contains a Group 5 insecticide. Insect biotypes with acquired resistance to Group 5 insecticides may eventually dominate the insect population if appropriate resistance management strategies are not followed. Currently, only spinetoram and spinosad active ingredients are classified as Group 5 insecticides. Resistance to other insecticide groups is not likely to impact the effectiveness of this product. Spinosad may be used in rotation with all other labeled products in a comprehensive IRM

To minimize the potential for resistance development, the following practices are recommended:

- · Base insecticide use on comprehensive IPM and IRM programs
- Do not use less than the labeled rates.
- Routinely evaluate applications for loss of effectiveness.
- Rotate with other labeled effective mosquito larvicides that have a different mode of action.
- In dormant rice fields, standing water within agricultural/crop sites, and permanent marine and freshwater sites, do not make more than 5 applications per year.
- Use insecticides with a different mode of action (different insecticide group) on adult mosquitoes so that both larvae and adults are not exposed to products with the same mode of
- Contact your local extension specialist, technical advisor, and/or Clarke Mosquito Control representative for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- · For further information or to report suspected resistance, you may contact your local Clarke Mosquito Control representative by calling 800-323-5727.

Proper application techniques help ensure adequate coverage and correct dosage necessary to obtain optimum kill of mosquito and midge larvae. Natular T30 tablets provide up to 30 days of residual kill. Natular T30 can be applied prior to flooding, on snow and ice in breeding sites prior to spring thaw, or at anytime after flooding in listed sites. Continue treating through the last brood of the season. Natular T30 tablets will be unaffected in dry down situations and will begin working again during subsequent wetting events until the tablet is exhausted. Note: Natular T30 has no effect on mosquitoes which have reached the pupal or adult stage prior to treatment.

Application Sites and Rates

Natular T30 tablets are designed to kill mosquitoes in small bodies of water. Do not apply to water intended for irrigation. Examples of application sites are:

Storm water drainage areas, sewers and catch basins, woodland pools, snow pools, roadside ditches, retention ponds, freshwater dredge spoils, tire tracks, rock holes, pot holes and similar areas subject to holding water.

Natural and manmade aquatic sites, fish ponds, ornamental ponds and fountains, other artificial water-holding containers, flooded crypts, transformer vaults, abandoned swimming pools, construction and other natural or manmade depressions.

Stream eddies, creek edges, detention ponds.

Freshwater swamps and marshes including mixed hardwood swamps, cattail marsh, common reed wetland, water hyacinth ponds, and similar freshwater areas with emergent vegetation.

Brackish water swamps and marshes, intertidal areas.

Sewage effluent, sewers, sewage lagoons, cesspools, oxidation ponds, septic ditches and tanks, animal waste lagoons and settling ponds, livestock runoff lagoons, wastewater impoundments associated with fruit and vegetable processing and similar areas.

Also for use in dormant rice fields (for application only during the interval between harvest and preparation of the field for the next cropping cycle) and in standing water within pastures/hay fields, rangeland, orchards, and citrus groves where mosquito breeding occurs. Do not apply to waters intended for irrigation.

For mosquito kill in non- or low-flow, shallow depressions (up to 2 feet in depth), treat on the basis of surface area placing 1 Natular T30 tablet per 100 sq ft.

For applications in storm water drainage areas, sewers and catch basins, place Natular T30 tablet into each catch basin.

For application sites connected by a water system, i.e., storm drains or catch basins, treat all of the water holding sites in the system to maximize the efficiency of the treatment program.

For application to small contained sites which may not be amenable to a rate of a single tablet per 100 sq ft, use 1 tablet per contained site (e.g., cesspools and septic tanks, transformer vaults, abandoned pools, and other small artificial water-holding containers).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool dry place in original container only.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Warran	t
--------	---

To the extent consistent with applicable law CLARKE MOSQUITO CONTROL PRODUCTS, INC. makes no warranty, express or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use/handling of this material when use and/or handling is contrary to label instructions.

Lot:	Net Weight:

Trademark of Dow AgroSciences LLC