

CROP and DISEASE	RATE of APPLICATION	METHOD OF APPLICATION
CABBAGE CABBAGE BROCCOLI BRUSSELS SPROUTS	150-300 lbs./acre	Row spacing: cabbage, broccoli, Brussels sprouts: 40 inch. Full row: 40 inch. Broadcast before transplanting. Mix into soil to a depth of 4-6". Lower rate for lighter soils.
Club Root <sup>Club Root</sup> (Plasmodiophora Brassicace)	100-200 lbs./acre	Row application in bands 12-15" wide before transplanting. Lower rate for lighter soils.
POTATOES (Phizoctonia solanii)	50-50 lbs./acre	Apply in 12-14" bands centered on the row as a preplanting treatment to all soil that covers and surrounds the seed piece and tuber forming zone. Rate based on 40-42" row spacing. To rot control 50 lbs. brassicol 20 per acre.
BULB'S IRIS Crown Rot Black Rot	500-1000 lbs./acre	Broadcast before planting. Mix thoroughly into soil to a depth of 6-7 inches. For black rot control and heavier soils, use the high rate.
	175-300 lbs./acre	Pre-plant application; 20-36 inch row spacing; at planting sides and edges of the open furrows have to be dusted before they are closed. Use the high rate for black rot control and heavier soils.
BULB'S IRIS Stem Rot Bulb Rot (Sclerotium rolfsii)	500-750 lbs./acre  30 lbs./acre	Broadcast before planting. Mix thoroughly into soil to a depth of 2-3".  Row application (30-40" row spacing) bands 12" wide, centered on row. Three additional applications at 15 day intervals.

U.S. Pat. No. 2,411,111  
 U.S. Pat. No. 2,411,112  
 U.S. Pat. No. 2,411,113  
 U.S. Pat. No. 2,411,114  
 U.S. Pat. No. 2,411,115  
 U.S. Pat. No. 2,411,116  
 U.S. Pat. No. 2,411,117  
 U.S. Pat. No. 2,411,118  
 U.S. Pat. No. 2,411,119  
 U.S. Pat. No. 2,411,120  
 U.S. Pat. No. 2,411,121  
 U.S. Pat. No. 2,411,122  
 U.S. Pat. No. 2,411,123  
 U.S. Pat. No. 2,411,124  
 U.S. Pat. No. 2,411,125  
 U.S. Pat. No. 2,411,126  
 U.S. Pat. No. 2,411,127  
 U.S. Pat. No. 2,411,128  
 U.S. Pat. No. 2,411,129  
 U.S. Pat. No. 2,411,130  
 U.S. Pat. No. 2,411,131  
 U.S. Pat. No. 2,411,132  
 U.S. Pat. No. 2,411,133  
 U.S. Pat. No. 2,411,134  
 U.S. Pat. No. 2,411,135  
 U.S. Pat. No. 2,411,136  
 U.S. Pat. No. 2,411,137  
 U.S. Pat. No. 2,411,138  
 U.S. Pat. No. 2,411,139  
 U.S. Pat. No. 2,411,140  
 U.S. Pat. No. 2,411,141  
 U.S. Pat. No. 2,411,142  
 U.S. Pat. No. 2,411,143  
 U.S. Pat. No. 2,411,144  
 U.S. Pat. No. 2,411,145  
 U.S. Pat. No. 2,411,146  
 U.S. Pat. No. 2,411,147  
 U.S. Pat. No. 2,411,148  
 U.S. Pat. No. 2,411,149  
 U.S. Pat. No. 2,411,150  
 U.S. Pat. No. 2,411,151  
 U.S. Pat. No. 2,411,152  
 U.S. Pat. No. 2,411,153  
 U.S. Pat. No. 2,411,154  
 U.S. Pat. No. 2,411,155  
 U.S. Pat. No. 2,411,156  
 U.S. Pat. No. 2,411,157  
 U.S. Pat. No. 2,411,158  
 U.S. Pat. No. 2,411,159  
 U.S. Pat. No. 2,411,160  
 U.S. Pat. No. 2,411,161  
 U.S. Pat. No. 2,411,162  
 U.S. Pat. No. 2,411,163  
 U.S. Pat. No. 2,411,164  
 U.S. Pat. No. 2,411,165  
 U.S. Pat. No. 2,411,166  
 U.S. Pat. No. 2,411,167  
 U.S. Pat. No. 2,411,168  
 U.S. Pat. No. 2,411,169  
 U.S. Pat. No. 2,411,170  
 U.S. Pat. No. 2,411,171  
 U.S. Pat. No. 2,411,172  
 U.S. Pat. No. 2,411,173  
 U.S. Pat. No. 2,411,174  
 U.S. Pat. No. 2,411,175  
 U.S. Pat. No. 2,411,176  
 U.S. Pat. No. 2,411,177  
 U.S. Pat. No. 2,411,178  
 U.S. Pat. No. 2,411,179  
 U.S. Pat. No. 2,411,180  
 U.S. Pat. No. 2,411,181  
 U.S. Pat. No. 2,411,182  
 U.S. Pat. No. 2,411,183  
 U.S. Pat. No. 2,411,184  
 U.S. Pat. No. 2,411,185  
 U.S. Pat. No. 2,411,186  
 U.S. Pat. No. 2,411,187  
 U.S. Pat. No. 2,411,188  
 U.S. Pat. No. 2,411,189  
 U.S. Pat. No. 2,411,190  
 U.S. Pat. No. 2,411,191  
 U.S. Pat. No. 2,411,192  
 U.S. Pat. No. 2,411,193  
 U.S. Pat. No. 2,411,194  
 U.S. Pat. No. 2,411,195  
 U.S. Pat. No. 2,411,196  
 U.S. Pat. No. 2,411,197  
 U.S. Pat. No. 2,411,198  
 U.S. Pat. No. 2,411,199  
 U.S. Pat. No. 2,411,200

For the control of various diseases, especially soil-borne diseases in vegetables, potatoes, cotton and flowers.

Caution: Keep out of reach of children  
 See back panel for additional cautions.

U.S. Pat. No. 2,411,140-149

It is a registered trademark of  
 Bayer AG, West Germany

U.S. Pat. No. 2,411,140-149  
 U.S. Pat. No. 2,411,150-159  
 U.S. Pat. No. 2,411,160-169  
 U.S. Pat. No. 2,411,170-179  
 U.S. Pat. No. 2,411,180-189  
 U.S. Pat. No. 2,411,190-199

ACCEPTED

340

CROP and DISEASE	RATE of APPLICATION	METHOD OF APPLICATION
GALENDULA LARKSPUR SNAPDRAGONS SWEET PEAS Stem Rot (Sclerotinia ssp.)	500 lbs./acre (1 lb. per 100 sq. ft.)	Broadcast before planting. Mix thoroughly into soil to a depth of 4".
<u>AZALLAS</u>		
Flower Blight (Ovulinia)	1000 lbs./acre (2.5 lbs. per 100 sq. ft.) Soil surface	Begin applications to soil surface when buds first show color. Repeat every 3,4 weeks covering soil surface thoroughly. Do not disturb treated surface.
CAMELLIAS <i>Camellias</i>		
Flower Blight (Sclerotinia ssp.)		
COTTON <i>Cotton</i>	10-25 lbs./acre	At planting Furrow application (42 inch row spacing). Dust over seeds and surrounding soil. Lower rate for lighter soils.
damping-off <i>Rhizoctonia solani</i>		
<u>GLADIOLUS <i>Gladiolus</i></u>		
neck or bulb dry Rot (Stromatinia or Sclerotinia)	600 lbs./acre  300 lbs./acre	Apply broadcast at planting. Thoroughly mix in top 4-6 inches of soil. Row treatment; Apply in 8-10" band centered on row at planting.

**CAUTION:**

This material may cause skin irritation. Wear protective clothing and approved respirator when loading the spreader or duster.

Avoid inhalation of the dust. Wash hands and face thoroughly after using. Avoid contamination of feed and food products.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Keep containers closed and dry. Do not reuse container. Destroy when empty.

**NOTICE TO BUYER:**

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith, or with established safe practice.