

file



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 2 1994

DIANE L. ROSSI
HOECHST CELANESE CORP.
C/O V A DOOR
RT 202-206 BOX 2500
SOMERVILLE, NJ 08876

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 9/15/93 Response to PR Notice 93-7
EPA Reg. No. 8340-14
PHASER INSECTICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canoa ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,


Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Diane L. Rosi
HOECHST CELANESE CORP
C/O V A DOOR
RT 202-206 BOX 2500
SOMERVILLE NJ 08876

Comment for: EPA Reg Nr.8340-14
PHASER INSECTICIDE

The following specific comments pertain to your WPS
labeling submission concerning the product
cited above:

Delete the crossed-out statements on your proposed label.
They are redundant statements or phrases.

41, 29

ACCEPTED
with COMMENTS
in EPA Letter Dated

Revised: ~~July 14, 1992~~
~~September 3, 1993~~

FEB 2 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
8340-14

PHASER[®] INSECTICIDE

For Agricultural or Commercial Use Only

ACTIVE INGREDIENT:

Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodio-
xathiepin-3-oxide).....33.70%*

INERT INGREDIENTS.....66.30%**

TOTAL: 100.00

* Equivalent to 3.0 pounds of active ingredient per gallon

** Contains xylene range aromatic solvents

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO



POISON



~~PRECAUTION AL USUARIO: Si usted no lee ingles, no use este producto hasta que
la etiqueta le haya sido explicada ampliamente.~~

~~Si Usted no entiende la etiqueta, busque a alguien para que se la explique a
Usted en detalle. (If you do not understand the label, find someone to
explain it to you in detail.)~~

EPA REG. NO. 8340-14

NET CONTENTS:

BEST AVAILABLE COPY

HOECHST CELANESE CORPORATION
Route 202-206
P.O. Box 2500
Somerville, NJ 08876-1258

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Vomiting should be induced only under the direct supervision of a physician. Do not induce vomiting or give anything by mouth to a person who is unconscious or convulsing.

If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on Skin: Remove contaminated clothing and wash skin with soap and water.

If in Eyes: Flush eyes with plenty of water. Call a physician immediately.

<p>For medical or environmental emergency, call 1-800-228-5635, ext. 132.</p>

Note to Physician: Endosulfan is a central nervous system stimulant absorbable by mouth, inhalation or through contact with skin. It may cause convulsions. There is no specific antidote. Diazepam I.V. is the drug of choice. Barbituric acid derivatives such as phenobarbital may be used additionally. A neuromuscular blocking agent may be used if convulsions persist. This type of drug may be used only if complete control of respiration can be maintained. Epinephrine derivatives are absolutely contraindicated.

This formulation contains petroleum hydrocarbons. Vomiting should be induced only under the direct supervision of a physician. Care should be taken to prevent aspiration because of the possibility of chemical pneumonia or pulmonary edema due to the organic solvent in the formulation.

PRECAUTIONARY STATEMENTS

DANGER

Hazards to Humans and Domestic Animals

Fatal if swallowed. May be fatal if inhaled or absorbed through skin. Do not breathe vapors or spray mist. Causes eye damage. Do not get in eyes, on skin, or on clothing. ~~Applicators must wear a long-sleeved shirt, trousers, unlined waterproof gloves made of natural rubber or neoprene, full-foot covering (such as shoes or boots), head covering (such as a hat), and a pesticide mask or respirator jointly approved by the Mine Safety and Health Administration (MSHA) and the National Institute for Occupational Safety and Health (NIOSH) under the provisions of 30 CFR Part 11 for protection against endosulfan. Mixers and/or loaders must wear a chemical resistant apron in addition to all of the protective equipment specified for applicators. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Wash thoroughly with soap and water after handling and before smoking or eating. Remove contaminated clothing and wash with soap and hot water before reuse.~~

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

Coveralls over short-sleeved shirt and short pants; chemical-resistant gloves, such as barrier laminate or viton 214 mils; chemical-resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical-resistant apron when cleaning equipment, mixing or loading; a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-140).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

BEST AVAILABLE COPY

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove shoes immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Due to the risk of runoff and drift, do not apply within a distance of 300 feet of lakes, ponds, streams and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label. This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

BEST AVAILABLE COPY

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Due to the risk of runoff and drift, do not apply within a distance of 300 feet of lakes, ponds, streams and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label. This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

BEST AVAILABLE COPY

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement of applicability, personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short-sleeved shirt and short pants; chemical-resistant gloves, such as barrier laminate or viton 214 mils; chemical-resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead exposure

~~Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.~~

~~Do not enter treated areas without protective clothing for 24 hours.~~

~~Protective clothing means, at least, a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall-type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.~~

~~Because certain states may require more restrictive re-entry intervals for various crops treated with this product, such as the 48-hour re-entry interval established by California, consult your State Department of Agriculture for further information.~~

BEST AVAILABLE COPY

~~Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated, and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:~~

DANGER

~~Area treated with endosulfan on (insert date). Do not enter without appropriate protective clothing for 24 hours (insert State Department of Agriculture's re-entry interval if more restrictive).~~

~~In case of accidental exposure to pesticide spray, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.~~

STORAGE AND DISPOSAL

Pesticide Storage

Do not store in or around the home.

Do not store below 20°F (-7°C).

Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or diluted material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill: Avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticides, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers.

BEST AVAILABLE COPY

GENERAL INSTRUCTIONS

Not for use or storage in or around the home.

Do not use in undiluted form.

Apply the listed amount per acre when insects first appear and repeat as required to maintain effective control. Use in sufficient water for thorough coverage of listed crops, unless otherwise noted. Coverage of upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 1 gallon of finished spray per acre on vegetable and field crops, unless otherwise noted, and a minimum of 20 gallons of finished spray per acre on fruit and nut trees and on vines. Where more than 2 quarts of this material are recommended, mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Phaser Insecticide used.

Do not plant root crops other than carrots, potatoes, sweet potatoes and sugar beets as follow-up crops.

Observe days interval between last application and harvest indicated by number in () following the crop.

INTEGRATED PEST MANAGEMENT (IPM)

Integrated Pest Management is the use of two or more methods of control to reduce damage by a pest or pests. These methods include: cultural practices, use of biological control agents such as predators and parasites, and the use of selective pesticides.

Phaser insecticide is the product of choice for IPM programs. Phaser used at specific rates will control certain pest species while allowing natural populations of predators and parasites to build to effective levels. When used in an IPM program, the potential for insect resistance to chemicals may also be greatly reduced. For maximum effectiveness the following is recommended:

1. Monitor fields to detect early infestation.
2. Apply Phaser when any pest species reaches the economic threshold for control and before significant crop injury occurs.
3. Monitor beneficial populations.
4. Repeat applications as needed within label directions.
5. If an integrated pest management program is desired, DO NOT tankmix Phaser insecticide with any broad spectrum insecticide.

ALFALFA (Grown for Forage) (21)

Insects Controlled	Rate of Application	Method of Application
Meadow Spittlebug	2/3 pint per acre	Apply 1 week after eggs begin to hatch or when first small masses of froth are produced in the crowns of the plant, which is usually when the alfalfa is about 4 to 6 inches tall. Apply in a minimum of 10 gallons of water per acre by ground or 2 gallons of water per acre by air.
<p>Do not graze or cut forage within 21 days of treatment. Do not apply more than once per cutting. Do not make more than 3 applications per year. Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.</p>		

APPLES (21)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphid) Apple Rust Mite Green Fruitworm Tarnished Plant Bug Tentiform Leafminer White Apple Leafhopper (first generation)	2/3 quart per 100 gallons or a maximum of 3-1/3 quarts per acre	Applications made at pink and/or petal fall provide best control of Tarnished Plant Bug and Green Fruitworm. For best control of first generation White Apple Leafhopper, apply when nymphs first appear on leaves. If necessary, prior to petal fall, use 1 quart per 100 gallons to control Apple Aphid and Rosy Apple Aphid. For control of Tentiform Leafminer, make first application as soon as moth flight begins. A second application should be made 10 days later.
<p>Do not feed pomace from treated apples to livestock. Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications during the fruiting period. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

APRICOTS (21), NECTARINES (21), PEACHES (21)

Insects Controlled	Rate of Application	Method of Application
Peachtree Borer Lesser Peachtree Borer	1 quart per 100 gallons - <u>West Coast</u> : 2/3 to 1 quart per 100 gallons - <u>Southeastern States</u> : 2 to 3-1/3 quarts per 100 gallons	Best control is obtained with a single application post-harvest during the first week of September. Spray all bark areas from ground level to lower scaffold limbs.
Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

APRICOTS (30), NECTARINES (30), PEACHES (30)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Black Cherry Aphid, Black Peach Aphid, Green Peach Aphid, Rusty Plum Aphid) Catfacing insects (Stink Bug type) Green Fruitworm Peach Silver Mite Peach Twig Borer	2/3 quart per 100 gallons or a maximum of 3-1/3 quarts per acre - <u>West Coast</u> : Do not use more than 4 quarts per acre.	Make applications when insects appear or feeding is noticed.
Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

ARTICHOKES (7)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Artichoke Aphid, Green Peach Aphid)	1 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. Apply in a minimum of 20 gallons of water per acre by ground and in a minimum of 10 gallons of water per acre by air.
Artichoke Plume Moth Whitefly	1-1/3 quarts per acre	
Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2-2/3 quarts) per acre per year.		

BARLEY, OATS, RYE, WHEAT

Insects Controlled	Rate of Application	Method of Application
Army Cutworm	2/3 quart per acre	For aerial application, apply in 2 gallons of diesel fuel oil per acre.
Cereal Leaf Beetle (Illinois, Indiana, Michigan and Ohio only)	1/3 to 2/3 quarts per acre	Apply when small larvae are readily found in the field. For aerial application, use a minimum of 1 to 2 gallons of water per acre.
Aphids	2/3 to 1 quart per acre	Make applications when insects appear or feeding is noticed.
Do not apply after heads begin to form. Do not feed treated forage to livestock. Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

BEANS, Succulent and Dry (except Lima Beans) (3) and SOUTHERN FIELD PEAS (Succulent type, including Black-eyed Peas, Crowder Peas and Southern Peas (3)

Insects Controlled	Rate of Application	Method of Application
Black Bean Aphid Bean Leaf Skeletonizer Cowpea Curculio Cucumber Beetle Flea Beetle Green Stink Bug Leafhopper Mexican Bean Beetle	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. For control of Cowpea Curculio, make 3 application at 5-day intervals starting when the pods are 1/2 inch long.
Aphids Armyworm Western Bean Cutworm Whitefly	1-1/3 quarts per acre	
Do not apply after buds are well formed. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

BLUEBERRIES

Insects Controlled	Rate of Application	Method of Application
Blueberry Bud Mite	2 quarts per 300 gallons	Apply immediately after harvest and repeat 6 to 8 weeks later.
Do not apply after buds are well formed. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

BROCCOLI (7), BRUSSEL SPROUTS (14), CABBAGE (7), CAULIFLOWER (14)

Insects Controlled	Rate of Application	Method of Application
Cabbage Aphid Cabbage Looper Cross-striped Cabbageworm Diamondback Moth (larvae) Flea Beetle Harlequin Bug Imported Cabbageworm Leafhopper Stink Bug	1 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Armyworm Cutworm Whitefly	1-1/3 quarts per acre	
Do not make more than 4 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

CARROTS (7)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Leafhopper	2/3 to 1-1/3 quarts per acre	Make application when insects appear or feeding is noticed.
Armyworm Flea Beetle Whitefly	1-1/3 quarts per acre	
Do not use tops for food or feed. Do not make more than 1 application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

CELERY (4)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Cabbage Looper Leafhopper	2/3 to 1-1/3 quarts per acre	Make application when insects appear or feeding is noticed.
Armyworm Flea Beetle Whitefly	1-1/3 quarts per acre	
Do not make more than 1 application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

CELERY (7)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid	2/3 quart per acre	Make applications when insects appear or feeding is noticed.
Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

CHERRIES (21)

Insects Controlled	Rate of Application	Method of Application
Peachtree Borer Lesser Peachtree Borer	1 quart per 100 gallons - <u>West Coast</u> : 2/3 to 1 quart per 100 gallons	Thoroughly wet all bark from ground to scaffold limbs.
Black Cherry Aphid Green Fruitworm Plum Rust (Nursery) Mite	2/3 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
<u>Pacific Northwest only</u> : Eyespotted Bud Moth Fruittree Leafroller		For Bud Moth control, apply at "popcorn" stage. For Leafroller control, apply during pre-pink stage of growth.
<u>Michigan only</u> : Mineola Moth	1-1/3 quarts per 100 gallons	Apply in the delayed dormant period.
<p>Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

CHERRY, PEACH, PLUM NURSERY STOCK DIP

Insects Controlled	Rate of Application	Method of Application
Peachtree Borer	2-2/3 quarts per 40 gallons	Mix thoroughly. Immerse trees so that the roots and crowns are covered well above the grafting bud scar.
<p>Wear rubber gloves during the dipping operation. Plant immediately or dry before returning stock to storage.</p>		

CITRUS (Non-Bearing Trees and Nursery Stock)

Insects Controlled	Rate of Application	Method of Application
Citrus Aphid	2/3 quart per 100 gallons or a maximum of 3-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
<p>Do not apply to bearing trees or trees that will bear fruit within 12 months. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

COLLARDS (21)

Insects Controlled	Rate of Application	Method of Application
Aphids Cabbage Looper Diamondback Moth (larvae) Fall Armyworm Flea Beetle Harlequin Bug Imported Cabbageworm Leafhopper	1 to 1-1/3 quarts per acre	Make application when insects appear or feeding is noticed.
Whitefly	1-1/3 quarts per acre	
<p>Do not make more than 1 application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.</p>		

COTTON

Insects Controlled	Rate of Application	Method of Application
Aphids	1/2 to 1 quart per acre	Make applications when insects appear or feeding is noticed. For control of Aphids, thorough coverage is important.
Boll Weevil	2/3 to 2 quarts per acre	
Bollworm Cabbage Looper Cotton Leaf-perforator Cotton Leafworm Fleahopper Lygus Bug Stink Bug Tobacco Budworm	1-1/3 to 2 quarts per acre	
Thrips	2 quarts per acre	
<p>Applications may be made using ground or aerial application equipment. The higher rate should be used under heavy pest pressure. Do not apply after bolls open. Do not graze dairy or meat animals in treated fields. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

CUCUMBERS (2), MELONS (2), PUMPKINS (2), SUMMER AND WINTER SQUASH (2)

Insects Controlled	Rate of Application	Method of Application
Aphids Cucumber Beetle Melonworm Pickleworm Rindworm (on watermelons) Squash Beetle Squash Bug Squash Vine Borer Striped Flea Beetle	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. For Squash Vine Borer control, apply weekly to flower buds, stems, and vines beginning when moths first appear.
Cabbage Looper Omniverous Leafroller Whitefly	1-1/3 quarts per acre	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

EGGPLANT (1)

Insects Controlled	Rate of Application	Method of Application
Blister Beetle Colorado Potato Beetle Flea Beetle Green Peach Aphid Green Stink Bug	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1-1/3 quarts per acre	
Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

GRAPES (7)

Insects Controlled	Rate of Application	Method of Application
Grape Leafhopper Grape Phylloxera (leaf form) Rose Chafer	2/3 quart per 100 gallons or 1-1/2 to 2 quarts per acre	Make applications when insects appear or feeding is noticed.
<p>Do not apply to Concord variety as severe injury is likely to occur. Ground application is preferred. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

KALE (21)

Insects Controlled	Rate of Application	Method of Application
Cabbage Flea Beetle Harlequin Bug Imported Cabbageworm	1 quart per acre	Make application when insects appear or feeding is noticed.
<p>Do not make more than 1 application per year. Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.</p>		

LETTUCE (14)

Insects Controlled	Rate of Application	Method of Application
Cabbage Looper Diamondback Moth (larvae) Green Peach Aphid Imported Cabbageworm Leafhopper	1 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Armyworm Whitefly	1-1/3 quarts per acre	
<p><u>On Head Lettuce:</u> Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. <u>On Leaf Lettuce:</u> Do not make more than 2 applications per year. Do not feed crop refuse to livestock. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

MUSTARD GREENS (21)

Insects Controlled	Rate of Application	Method of Application
Aphids Cabbage Looper Diamondback Moth (larvae) Fall Armyworm Flea Beetle Harlequin Bug Imported Cabbageworm Leafhopper	1 to 1-1/3 quarts per acre	Make application when insects appear or feeding is noticed.
Whitefly	1-1/3 quarts per acre	
<p>Do not make more than 1 application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.</p>		

PEARS (7)

Insects Controlled	Rate of Application	Method of Application
Green Fruitworm Tarnished Plant Bug	2/3 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Make applications at white bud or petal fall when insects appear or feeding is noticed.
Pear Psylla	3-1/3 quarts per acre	Apply in a minimum of 10 gallons of water for aerial application or in 300 gallons of water for dilute application. Apply when adults are first observed or nymphs are small and repeat to maintain control.
Pear Rust Mite Conspere Stink Bug (foliar treatment)	2/3 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. Stink Bugs must be wet by spray to obtain control.
Conspere Stink Bug (soil treatment)	2/3 quart per 100 gallons; 200 to 400 gallons per acre	Apply to orchard floor and around trees prior to bloom.
Pear Leaf Blister Mite	1/3 to 2/3 quart per 100 gallons	Apply as a post harvest or dormant treatment.
<p>Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p> <p>NOTE: Aerial application may not result in satisfactory control and should only be employed if impossible to apply by ground.</p>		

PEAS (Seed Crop Only) (1)

Insects Controlled	Rate of Application	Method of Application
Pea Aphid Pea Weevil	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
<p>Use only on peas to be harvested by combine. Do not feed treated vines or threshings to livestock or allow livestock to graze in treated fields. Do not make more than 2 applications per year. Do not exceed a maximum of 1.5 lbs. active ingredient (i.e., 2 quarts) per acre per year.</p>		

PECANS

Insects Controlled	Rate of Application	Method of Application
Black Pecan Aphid Pecan Nut Casebearer Spittlebug	1 quart to 100 gallons	Make applications when insects appear or feeding is noticed. For Casebearer, apply when eggs of first generation appear on the tips of the young nuts. Another application may be required after the second generation of eggs is deposited. For Spittlebug, apply when first leaves are half grown and repeat as required.
Pecan Leaf Phylloxera	2/3 to 1 quart per 100 gallons	Apply when nymphs appear and before they are enclosed in plant tissue. For high populations, use the higher rate.
<p>Do not apply after shuck split. Do not graze livestock on orchard crops or grasses in treated areas. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

PEPPERS (1)

Insects Controlled	Rate of Application	Method of Application
Flea Beetle Green Peach Aphid Hornworm Pepper Maggot	2/3 quart per acre	Make applications when insects appear or feeding is noticed.
Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

PEPPERS (4)

Insects Controlled	Rate of Application	Method of Application
Armyworm Flea Beetle Green Peach Aphid Hornworm Leafhopper Pepper Maggot	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1-1/3 quarts per acre	
Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2-2/3 quarts) per acre per year.		

PINEAPPLE (For Fresh Market Only) (7)

Insects Controlled	Rate of Application	Method of Application
Pineapple Fruit Mite	2 to 2-2/3 quarts per acre	Make applications when insects appear or feeding is noticed. Apply at intervals of 7 to 10 days, if necessary, particularly during the 40-day blooming period.
Do not feed forage or pineapple by-products to livestock. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

PLUMS (7), PRUNES (7)

Insects Controlled	Rate of Application	Method of Application
Fruittree Leafroller (Pacific Northwest Only)	2/3 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Apply during pre-pink stage of growth when insects appear or feeding is noticed.
Aphids (including Hop Aphid, Leafcurl Aphid, Thistle Aphid Plum Rust (Nursery) Mite		For control of Aphids, apply when Spray thoroughly to cover all bark areas from ground to scaffold limbs.
Peach Twig Borer	2/3 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Lesser Peachtree Borer	1 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Apply spray to thoroughly wet the trunk and main branches.
Peachtree Borer	1 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre <u>West Coast:</u> 2/3 to 1 quart per 100 gallons or 2-2/3 to 3-1/3 quarts per acre	Spray thoroughly to cover all bark area from ground to scaffold limbs.
Do not allow livestock to graze on orchard crops or grasses in treated areas. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

POTATOES (1)

Insects Controlled	Rate of Application	Method of Application
Aphids Armyworm Colorado Potato Beetle Green Stink Bug Leaffooted Bug Plant Bug Potato Flea Beetle Potato Leafhopper Potato Tuberworm Threelined Potato Beetle	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
European Corn Borer Potato Psyllid	1 to 1-1/3 quarts per acre	
False Chinch Bug Whitefly	1-1/3 quarts per acre	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

SAFFLOWER

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Whitefly	1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Do not apply after flower heads open. Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2-2/3 quarts) per acre per year.		

SPINACH (21)

Insects Controlled	Rate of Application	Method of Application
Armyworm Crown Mite Flea Beetle Green Peach Aphid Leafhopper	1 to 1-1/3 quarts per acre	Make application when insects appear or feeding is noticed.
Whitefly	1-1/3 quarts per acre	
Do not make more than 1 application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1-1/3 quarts) per acre per year.		

STRAWBERRIES (4)

Insects Controlled	Rate of Application	Method of Application
Meadow Spittlebug Strawberry Aphid Tarnished Plant Bug Whitefly	1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. Do not reapply within 15 days or more than twice during a 35-day period when fruit is present.
Cyclamen Mite	2-2/3 quarts per acre in 400 gallons	Make applications when insects appear or feeding is noticed. Thoroughly wet the foliage, stem and crown of the plant. For multiple applications, do not apply at intervals less than 35 days when fruit is present.
Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

STRAWBERRIES - Northwest Use Only

Insects Controlled	Rate of Application	Method of Application
Garden Symphylan (aids in reducing damage)	1-1/3 quarts per 100 gallons	Mix thoroughly. Dip entire plant. When immersing bundles of plants, make certain any trapped air is forced out to assure thorough wetting of the entire plant.
<p>Wear rubber gloves during the dipping operation. Drain and allow plants to dry before setting them out in the field.</p>		

SUGAR BEETS (30)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Meadow Spittlebug	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. Make one or two applications per year as needed for control.
Sugar Beet Webworm	1 to 1-1/3 quarts per acre	
Whitefly	1-1/3 quarts per acre	
<p>Do not feed treated tops to livestock. Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2-2/3 quarts) per acre per year.</p>		

SUNFLOWERS (1)

Insects Controlled	Rate of Application	Method of Application
Sunflower Moth	1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed. Make the first application at the onset of bloom. Allow 7 days between repeat applications.
<p>Do not feed treated forage to livestock. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

SWEET CORN (Fresh Vegetable Use Only) (1)

Insects Controlled	Rate of Application	Method of Application
Corn Leaf Aphid Whitefly	1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Corn Earworm	2 quarts per acre	Apply when silks first appear and continue until they begin to dry. Allow 5 days between applications.
<p>Do not apply to Sweet Corn to be processed. Do not feed treated forage or ensilage to livestock or allow livestock to graze in treated fields. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

SWEET POTATOES (1)

Insects Controlled	Rate of Application	Method of Application
Sweet Potato Flea Beetle Sweet Potato Weevil	2/3 quarts per acre	Make applications when insects appear or feeding is noticed. For Flea Beetle control, begin applications shortly after transplanting or as soon as flea beetles appear. For Sweet Potato Weevil control, apply in sufficient water for complete coverage. Applications may be made to transplants in the nursery and/or in the field. Repeat applications may be made as necessary. For Sweet Potato Weevil control in a clean-up program, apply at a rate of 2-2/3 quarts per acre to the soil under the sweet potato nursery beds and to the area immediately surrounding the nursery beds. Do not place the seed tubers directly on the ground treated with Phaser Insecticide.
Whitefly	1-1/3 quarts per acre	Make applications when insect activity or feeding is noticed.
Banded Cucumber Beetle (larvae) (aids in control) (South Central States and Puerto Rico only)	1-1/3 to 2-2/3 quarts per acre broadcast or 1/2 to 1 quart per acre on a 16-inch band over the row (48-inch row spacing)	Work into the soil to a depth of approximately 3 inches. Treatment should be made just prior to plant set.
<p>Do not feed cull potatoes to livestock or allow livestock to graze in treated fields.</p> <p>Do not make more than 3 applications per year.</p> <p>Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

TOBACCO (5)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Green Peach Aphid, Tobacco Aphid) Budworm Cabbage Looper Flea Beetle Hornworm	<u>Seed Bed:</u> 2/3 quart per 100 gallons	Make applications when insects appear or feeding is noticed. Apply approximately 6 gallons of finished spray per 100 square yards.
Green June Bug (larvae)	<u>Plant Bed:</u> 2/3 pint per 100 gallons	Make applications when insects appear or feeding is noticed. Drench at a rate of 1 gallon per square yard.
Aphids (including Green Peach Aphid, Tobacco Aphid) Budworm Cabbage Looper Flea Beetle Green June Bug (larvae) Hornworm	<u>Field:</u> 2/3 to 1-1/3 quarts per acre	Make applications when insect activity or feeding is noticed.
Stink Bug	<u>Field:</u> 2/3 quart per 100 gallons	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

TOMATOES (Field and Greenhouse) (2)

Insects Controlled	Rate of Application	Method of Application
Aphids Blister Beetle Colorado Potato Beetle Flea Beetle Tomato Hornworm Webworm (California only)	2/3 to 1-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Cabbage Looper Stink Bug	1 to 1-1/3 quarts per acre	
Tomato Fruitworm Tomato Russet Mite Whitefly Yellowstriped Armyworm	1-1/3 quarts per acre	
Whitefly	2/3 quart per 100 to 200 gallons of water	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

WALNUTS

Insects Controlled	Rate of Application	Method of Application
Walnut Aphid	2 to 3-1/3 quarts per acre	Make applications when insects appear or feeding is noticed.
Do not apply after husk split. Do not graze livestock on orchard crops or grasses in treated areas. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**COMMERCIALLY GROWN ORNAMENTALS
ORNAMENTAL PLANTS**

LEATHERLEAF FERN (Leather Holly Fern)

Insects Controlled	Rate of Application	Method of Application
Leatherleaf Fern Borer	2/3 quart per 100 gallons	Begin treatment when first larval feeding is observed in the mid-vein area at the base of the leaflets. Repeat at intervals of 2 to 3 weeks as necessary.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

ORNAMENTALS (Greenhouse and Out-of-Doors)

Insects Controlled	Rate of Application	Method of Application
Aphids Cyclamen Mite Rose Chafer Whitefly	2/3 quart per 100 gallons	Make applications when insects appear or feeding is noticed.
<p>On chrysanthemums, best results will be obtained if applied before plants flower.</p> <p>Do not apply to "Bonnafon Deluxe", "Fred Shoesmith", and "White Knight" chrysanthemums as injury may result.</p> <p>Do not use on Birch trees.</p> <p>Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.</p>		

ORNAMENTAL TREES AND SHRUBS

DOGWOOD, LILAC

Insects Controlled	Rate of Application	Method of Application
Dogwood Borer Lilac Borer	2/3 to 1-1/3 quarts per 100 gallons	Apply in early June and repeat in 10-14 days. Drench all bark areas down to the ground level.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**DOUGLAS FIR (Grown for Ornamentals, Nursery Stock or Christmas Trees)
Pacific Northwest Only**

Insects Controlled	Rate of Application	Method of Application
Aphids Cooley Spruce Gall Adelgid Douglas Fir Needle Midge	2/3 quart per 100 gallons	Make applications when insects appear or feeding is noticed. For control of Gall Adelgid, apply when white cottony tufts appear. For control of Needle Midge, apply in late April or early May just before buds open.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

PINES (Austrian, Jack, Red, Scotch, White)

Insects Controlled	Rate of Application	Method of Application
Zimmerman Pine Moth	1 quart per '00 gallons	Apply in mid-April and again, if necessary, in late fall. Thoroughly wet bark and main stem, especially where branches join main stem.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

SHADE TREES (except Birch), SHRUBS

Insects Controlled	Rate of Application	Method of Application
Aphids	2/3 quart per 100 gallons	Make applications when insects appear or feeding is noticed.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

SPRUCE (New England Area Only)

Insects Controlled	Rate of Application	Method of Application
Spruce Gall Aphid	2/3 quart per 100 gallons	Apply in late April or early May when aphids are present but before galls are formed.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

TAXUS

Insects Controlled	Rate of Application	Method of Application
Taxus Bud Mite	2/3 quart per 100 gallons	Make 3 to 5 applications beginning in mid-May. Thoroughly spray foliage, twigs and bark.
Black Vine Weevil	1-1/3 quarts per 100 gallons	Spray thoroughly and drench the soil under the plants. Apply when the weevils first appear and repeat in 14 days.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

IMPORTANT NOTICE: DISCLAIMER

Read "IMPORTANT NOTICE: DISCLAIMER" before buying or using. If terms are not acceptable, return at once unopened.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, or use of this product in a manner inconsistent with its labeling, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

HOECHST CELANESE CORPORATION warrants only that the product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to HOECHST CELANESE CORPORATION and user assumes the risk of any such use. HOECHST CELANESE CORPORATION MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING ANY IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR OF MERCHANTABILITY. In no case shall HOECHST CELANESE CORPORATION be liable for consequential, special, indirect or incidental damages resulting from the use or handling of this product. The foregoing conditions of sale and warranty can be varied only by an agreement in writing signed by a duly authorized representative of HOECHST CELANESE CORPORATION.

TH9215001 (1/92)

File:\msd\wp\dca\dir998.503