


APPLICATORS OR PERSONS UNDER THEIR DIRECT
SUPERVISION AND ONLY FOR THOSE USES COVERED
BY THE CERTIFIED APPLICATOR'S CERTIFICATION

supplemental label for

Hoelon
3EC HERBICIDE



For Early Postemergence Use For Wild Oats

IMPORTANT

Before using this product, read and carefully observe all applicable directions, restrictions and precautionary statements on the EPA approved Hoelon 3EC container and use directions labels.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application.

THIS PRODUCT IS RESTRICTED FOR USE ON THE FOLLOWING CROP IN THE FOLLOWING STATES:

LENTILS: Idaho and Washington

GENERAL INFORMATION: HOELON 3EC Herbicide is an emulsifiable concentrate for selective postemergence use in the control of the following:

WILD OATS.....Avena fatua

Mix HOELON 3EC in water and apply as a broadcast postemergence spray at the rates listed below

Crop	Stage of Growth (Annual Grassy Weeds)	Amount of HOELON 3EC per acre treated*	Equipment Notes
Lentils	1-4 leaves more than 4 leaves	2-2½ pints Do not apply	Thorough uniform spray coverage of weeds is essential. Ground applications should be in at least 10 gal. of water using a minimum of 40 psi and use of fan type or hollow cone nozzles is recommended. Only ground applications should be made on lentils. Do not make applications within 100 ft. of a lake, stream, pond, drainage basin, tidal marsh or estuary.

Special Notes

1. HOELON 3EC does NOT control broadleaf weeds or perennial grassy weeds.
2. HOELON 3EC is a slow acting product, therefore, visual results will not be evident immediately following application.
3. Do NOT apply more than one application of HOELON in a growing season.
4. Do NOT allow livestock to graze treated fields. Do NOT harvest forage, hay or straw from treated fields prior to crop harvest.
5. Do not tank mix with any broadleaf herbicide for use on lentils.

Spray Hoelon 3EC When the Majority of Wild Oats is in the Growth Stages Shown Below*

Sprout	Annual Grassy Weed Stage				More than 4 leaves
	1 leaf	2 leaf	3 leaf	4 leaf	
    					
	Use 2-2½ pints				Do not apply

BEST AVAILABLE COPY

*Apply the two pint rate only when the annual grassy weeds are in the 1-2 leaf stage of growth and growing conditions are normal.
Higher rates should be used (see rate chart above) when the majority of grassy weeds is not clearly in the specified stage of growth or if growth is retarded due to lack of moisture. Thorough spray coverage is essential.

