

CIDE

Vet Company **Hoechst**
NJ 08876 **Roussel**

ONE SHOT™

HERBICIDE

Active Ingredients: Cefazolin sodium
5-hydroxy-4-(hydroxyphenyl)-3-isoxazole
thiazolidine-2-thione-6-thiol
5-hydroxy-4-(hydroxyphenyl)-3-isoxazole
thiazolidine-2-thione-6-thiol
5-hydroxy-4-(hydroxyphenyl)-3-isoxazole
thiazolidine-2-thione-6-thiol

[illegible]

product may cause a 3rd term miscarriage. A 3rd term miscarriage is a miscarriage that occurs after the 28th week of pregnancy. It is a rare event, but it can be a serious complication of pregnancy. It is caused by a variety of factors, including infection, trauma, and hormonal imbalances. It is important to seek medical attention if you experience any of the following symptoms: vaginal bleeding, abdominal pain, or a change in your vaginal discharge. A 3rd term miscarriage is a medical emergency, and it is important to seek medical attention as soon as possible.

STATE OF NEW YORK
IN SENATE
JANUARY 11, 1966
REPORT
OF THE
COMMISSIONERS OF THE DEPARTMENT OF SOCIAL SERVICES
IN RESPONSE TO
RESOLUTION NO. 100, PASSED BY THE SENATE
MAY 11, 1965
AND
RESOLUTION NO. 101, PASSED BY THE SENATE
MAY 11, 1965
ALBANY: THE UNIVERSITY OF THE STATE OF NEW YORK PRESS, 1966
PUBLISHED BY AUTHORITY OF THE SENATE

Note to Physician:

Storage and Disposal:

Pesticide Disposal. The following information is provided for the safe storage and disposal of pesticides. For more information, contact your local extension agent or the National Pesticide Information Center at 1-800-858-7373.

Container Disposal: The following information is provided for the purpose of ensuring that the container is disposed of in a safe and secure manner. The container should be disposed of in a secure manner and should not be disposed of in a public area. The container should be disposed of in a secure manner and should not be disposed of in a public area. The container should be disposed of in a secure manner and should not be disposed of in a public area.

SEE ATTACHED FOR REFOR COMPLETE DIRECTIONS OF USE AND ADDITIONAL INFORMATION

ONE SHOT

HERBICIDE

- One twin container treats 10 acres.
- Remove this front panel for graduation marks to be used for applications of less than 10 acres.

NOT REVIEWED
In Accordance with FR Notice 82-2.
Based on Draft Labeling Dated

ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

Marketed By: Hoechst-Roussel Agri-Vet Company
Agricultural Chemicals • Somerville, NJ 08876



RESTRICTED USE PESTICIDE
For Retail Sale to and Use Only by Certified Applicators or Persons Under their Direct Supervision and Only for Those Uses Covered by the Certified Applicator's Certification

For Retail Sale to and Use Only by Certified Applicators or Persons Under their Direct Supervision and Only for Those Uses Covered by the Certified Applicator's Certification

HERBICIDE

NOT REVIEWED
In Accordance with FR Notice 82-24
Active Ingredients: ~~glyphosate~~
DRAFT LABELING DATED 11/1/82
ILLOXAN® 3 EC HERBICIDE
KEEP OUT OF REACH OF CHILDREN.
STORAGE AND DISPOSAL

Active Ingredients: ~~See Draft Labeling Paged~~
Inert Ingredients: ~~See Draft Labeling Paged~~

[illegible]

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

Inert Ingredients

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

SEE ATTACHED SECTION FOR COMPLETE DIRECTIONS OF USE AND ADDITIONAL INFORMATION

1. The first step is to identify the problem. In this case, the problem is that the system is not working properly.

SEE ATTACHED FOLDER FOR COMPLETE DIRECTIONS OF USE AND ADDITIONAL INFORMATION



ONE SHOT™

HERBICIDE
~~Not Registered~~

In Accordance with PR Notice 82-2,
Based on Draft Labeling Data
Important Instructions Enclosed

Manufactured by
Hoechst-Roussel Agri-Vet Company
Agricultural Chemicals Division
Somerville, N.J. 08876

Hoechst 
Roussel 

RESTRICTED USE PESTICIDE.

For sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

ONE SHOT

HERBICIDE

ONE SHOT is a selective herbicide for use in corn, soybeans, and other crops. It is effective against a wide range of weeds, including annual and perennial species. It is applied as a spray or granule, and its use is restricted to certified applicators or persons under their direct supervision.

ONE SHOT is a selective herbicide for use in corn, soybeans, and other crops. It is effective against a wide range of weeds, including annual and perennial species. It is applied as a spray or granule, and its use is restricted to certified applicators or persons under their direct supervision.

GENERAL INFORMATION

ONE SHOT is a selective herbicide for use in corn, soybeans, and other crops. It is effective against a wide range of weeds, including annual and perennial species. It is applied as a spray or granule, and its use is restricted to certified applicators or persons under their direct supervision.

ONE SHOT is a selective herbicide for use in corn, soybeans, and other crops. It is effective against a wide range of weeds, including annual and perennial species. It is applied as a spray or granule, and its use is restricted to certified applicators or persons under their direct supervision.

DIRECTIONS FOR USE IN SPRING WHEAT AND SPRING BARLEY

ONE SHOT is a selective herbicide for use in spring wheat and spring barley. It is effective against a wide range of weeds, including annual and perennial species. It is applied as a spray or granule, and its use is restricted to certified applicators or persons under their direct supervision.

MIXING INSTRUCTIONS

ONE SHOT is a selective herbicide for use in spring wheat and spring barley. It is effective against a wide range of weeds, including annual and perennial species. It is applied as a spray or granule, and its use is restricted to certified applicators or persons under their direct supervision.

1. The purpose of this document is to provide information regarding the application of the product to various surfaces. The information is intended to assist the user in selecting the appropriate application method and to ensure the proper use of the product.

APPLICATION INFORMATION:

Ground Application

For ground application, the product should be applied to the surface in a uniform, continuous layer. The application should be made in a direction perpendicular to the surface. The product should be applied in a layer that is approximately 1/8 inch thick. The application should be made in a uniform, continuous layer.

Aerial Application

For aerial application, the product should be applied to the surface in a uniform, continuous layer. The application should be made in a direction perpendicular to the surface. The product should be applied in a layer that is approximately 1/8 inch thick. The application should be made in a uniform, continuous layer.

CONTROL

The product should be applied to the surface in a uniform, continuous layer. The application should be made in a direction perpendicular to the surface. The product should be applied in a layer that is approximately 1/8 inch thick. The application should be made in a uniform, continuous layer.

CONTROL

The product should be applied to the surface in a uniform, continuous layer. The application should be made in a direction perpendicular to the surface. The product should be applied in a layer that is approximately 1/8 inch thick. The application should be made in a uniform, continuous layer.


The product should be applied to the surface in a uniform, continuous layer. The application should be made in a direction perpendicular to the surface. The product should be applied in a layer that is approximately 1/8 inch thick. The application should be made in a uniform, continuous layer.

Application Timing

SPECIAL NOTES:


Hoechst 
Roussel 

Hoechst-Roussel Agri-Vet Company



SUPPLEMENTAL LABELING

FOR PREPLANT INCORPORATED USE IN THE CONTROL OF CERTAIN BROMEGRASS SPECIES IN FALL PLANTED WHEAT, IN THE STATES OF COLORADO, IDAHO, KANSAS, MONTANA, NEBRASKA, OREGON, SOUTH DAKOTA, AND WASHINGTON



HOELON^R 3 EC HERBICIDE
(EPA REG. NO. 8340-20-54382)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

HOELON 3 EC is a selective herbicide for fall planted wheat (winter wheat) that when incorporated into the soil prior to planting will control Downy Brome (Bromus tectorum) and Ripgut Brome (Bromus rigidus) as germination occurs. HOELON 3 EC will not control established brome-grasses.

Prior to application, crop residues should be mixed into the soil and the resulting seedbed should be friable and free of clods.

APPLICATION INFORMATION

HOELON 3 EC should be applied at a rate of 2 2/3 - 3 1/3 pints per acre. Use the high rate when high densities of brome-grasses (more than 100 plants per square foot) are expected and/or conditions are very dry at the time of application. For ground applications, HOELON 3 EC should be diluted in at least 10 gallons of water and applied using a minimum of 40 psi. Aerial applications should be in at least 5 gallons of water.

Do not make aerial applications when wind is above 5 mph or within 10 ft. of a lake, stream, pond, drainage basin, tidal marsh or estuary.

INCORPORATION INFORMATION

Thorough incorporation to a depth of at least 2 inches but not greater than 3 inches of soil in the final seedbed within 48 hours of application is necessary to ensure effective weed control regardless of weather conditions. Incorporate twice over in opposite directions for uniform distribution of HOELON 3 EC. If the seedbed is clean, use a rod weeder followed by a flex tine harrow or shallow disc. If the seedbed is trashy, use a skew treader. This technique is effective in conventional and deep furrow plantings.

BEST AVAILABLE COPY

NOTE:

1. HOELON 3 EC does NOT control broadleaf weeds or perennial grassy weeds.
2. Do NOT tank mix HOELON 3 EC with any other herbicide for preplant incorporation.
3. Do NOT allow livestock to graze treated fields. Do NOT harvest forage, hay or straw from treated fields.
4. Do NOT apply more than one application of HOELON in a growing season.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

This labeling must be in the possession of the user
at the time of pesticide application.

Marketed by:
Hoechst-Roussel Agri-Vet Company
Agricultural Chemicals
Somerville, N.J. 08876

The name and logo HOECHST are registered trademarks of Hoechst AG
The name and logo ROUSSEL are registered trademarks of Roussel Uclaf S.A.

Hoechst 
Roussel 

