

**RESTRICTED USE PESTICIDE**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by certified applicator's certification.

**ACCEPTED****MAY 19 1980**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

supplemental label for

**Hoelon**  
3EC HERBICIDE

Dry Peas (Field Peas) Including  
Austrian Winter Peas

**For Early Postemergence Use For Wild Oats****IMPORTANT**

Before using this product, read and carefully observe all applicable directions, restrictions and precautionary statements on the EPA approved Hoelon 3EC container and use directions labels

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application.

THIS PRODUCT IS RESTRICTED FOR USE ON THE FOLLOWING CROP IN THE FOLLOWING STATES  
DRY PEAS (FIELD PEAS) INCLUDING AUSTRIAN WINTER PEAS Idaho, Oregon and Washington

**GENERAL INFORMATION:** HOELON 3EC Herbicide is an emulsifiable concentrate for selective post-emergence use in the control of the following  
WILD OATS - *Avena fatua*







Dilute HOELON 3EC in water and apply as a broadcast  
postemergence spray at the rates listed below:

Crop	Stage of Growth (Annual Grassy Weeds)	Amount of HOELON 3EC per acre treated*	Equipment Notes
Dry Peas (Field Peas) Including Austrian Winter Peas	1-4 leaves	2-2½ pints	Thorough uniform spray coverage of weeds is essential. Ground applications should be in at least 10 gal. of water using a minimum of 40 psi, and use of fan type or hollow cone nozzles is recommended. Only ground applications should be made on dry peas. Do not make applications within 100 ft. of lake, stream, pond, drainage basin, tidal marsh or estuary.
	More than 4 leaves	(Do not apply)	

**SPECIAL NOTES:**

- 1 HOELON 3EC does NOT control broadleaf weeds or perennial grassy weeds
- 2 HOELON 3EC is a slow-acting product, therefore, visual results will not be evident immediately following application.
- 3 Do NOT apply more than one application of HOELON in a growing season ✓
- 4 Do NOT allow livestock to graze treated fields. Do NOT harvest pea vine forage or hay from treated fields for livestock feed
- 5 Do NOT tank mix with any broadleaf herbicide for use on dry peas

Spray HOELON 3EC When the Majority of Wild Oats  
Is in the Growth Stages Shown Below\*

Sprout	1 leaf	2 leaf	3 leaf	4 leaf	More than 4 leaves
					
	Use 2 pints		Use 2½ pints		Do not apply

\*Apply the two pint rate only when the annual grassy weeds are in the 1-2 leaf stage of growth and growing conditions are normal. Higher rates should be used (see rate chart above) when the majority of grassy weeds is not clearly in the specified stage of growth or if growth is retarded due to lack of moisture. Thorough spray coverage is essential.

EPA REG. NO. 8340-20-54382

Marketed by  
**Hoechst-Roussel Agri-Vet Company**  
Agricultural Chemicals  
Somerville, N.J. 08876

**Hoechst**   
**Roussel** 

The name and logo of HOECHST are registered trademarks of Hoechst AG.  
The name and logo of ROUSSEL are registered trademarks of Roussel Uclaf.

7186-10118-15-861