

ma3 8340-20

LOF2

SEP 28 1988

Dr. Bert Volger
Hoechst Celanese Corporation
Route 202-206 North
Somerville, NJ 08876

Dear Dr. Volger:

Subject: Tank Mix - Preplant - Glean
Hoelon 3EC Herbicide
EPA Registration No. 8340-20
Your Submission Dated September 7, 1988

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

For clarity, move the tank mix directions (paragraph 2) out of "Special Notes," and place in the Directions for Use. Refer to the Preplant Supplemental labeling accepted by this office on July 18, 1986.

Sincerely yours,

Richard F. Mountfort *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosure

BEST AVAILABLE COPY

51078:I:Mountfort:LMF-09:KENCO:09/26/88:10/06/88:aw:VO:aw

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

SEP 28 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended for the pesticide
EPA Reg. No.

SUPPLEMENTAL LABELING

HOELON^R 3EC HERBICIDE
(EPA Reg. No. 8340-20-54382)

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified
Applicators or persons under their direct
supervision and only for those uses covered by
the certified applicator's certification.

DIRECTIONS FOR USE FOR TANK MIXES

IMPORTANT

Before using these products, read and carefully observe all
applicable directions, restrictions, and precautionary statements
on the EPA approved Hoelon 3EC Herbicide Label and the Glean
Herbicide Label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner
inconsistent with its labeling. This labeling must be in the
possession of the user at the time of application.

General Notes:

1. HOELON^R 3EC Herbicide does not control broadleaf weeds or
perennial grassy weeds.
2. DO NOT allow livestock to graze treated fields. DO NOT
harvest forage, hay or straw from treated fields prior
to grain harvest.
3. HOELON^R 3EC Herbicide can be tankmixed with the broadleaf
herbicides listed on this label for broad spectrum weed
control.
4. DO NOT apply phenoxy, phenoxy-type, or Banvel^R Herbicide
within 5 days of a HOELON^R 3EC Herbicide application, or
reduced grassy weed control will occur.
5. DO NOT apply HOELON^R 3EC Herbicide through any type of
irrigation system.

PRE-PLANT INCORPORATED SPECIAL NOTES

1. DO NOT apply HOELON^R 3EC Herbicide pre-plant incorporated to barley, as barley damage will occur.
2. HOELON^R 3EC Herbicide may be tank mixed with Glean^R Herbicide for control of annual broadleaf weeds. HOELON^R 3EC Herbicide should be used at the 2.0 to 2 2/3 pint per acre rate while Glean^R should be used at 1/3 to 1/2 oz. per acre. Refer to the EPA-approved Glean^R label for crop rotation restrictions and weed species controlled at the above-mentioned rates. Prior to the use of Glean^R determine the soil pH by laboratory analysis using 1:1 soil:water suspension. Do not use Glean^R on soils with a pH greater than 7.9.
3. DO NOT tank mix HOELON^R 3EC Herbicide with any other herbicides for pre-plant incorporation unless specifically recommended in this label.
4. HOELON^R 3EC Herbicide may be tank mixed with liquid fertilizers. Follow label recommendations concerning rates and incorporation. Before mixing HOELON^R 3EC Herbicide with fertilizer, perform a compatibility test in a quart jar. If the materials do not readily mix, do not tank mix HOELON^R 3EC Herbicide with the fertilizer.

BEST AVAILABLE COPY

SUPPLEMENTAL LABELING

8340-20
HOELON^R 3EC HERBICIDE
(EPA Reg. No. 8340-20-54382)

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

DIRECTIONS FOR USE FOR TANK MIXES

IMPORTANT

Before using these products, read and carefully observe all applicable directions, restrictions, and precautionary statements on the EPA approved Hoelon 3EC Herbicide Label and the Glean Herbicide Label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application.

General Notes:

1. HOELON^R 3EC Herbicide does not control broadleaf weeds or perennial grassy weeds.
2. DO NOT allow livestock to graze treated fields. DO NOT harvest forage, hay or straw from treated fields prior to grain harvest.
3. HOELON^R 3EC Herbicide can be tankmixed with the broadleaf herbicides listed on this label for broad spectrum weed control.
4. DO NOT apply phenoxy, phenoxy-type, or Banvel^R Herbicide within 5 days of a HOELON^R 3EC Herbicide application, or reduced grassy weed control will occur.
5. DO NOT apply HOELON^R 3EC Herbicide through any type of irrigation system.

PRE-PLANT INCORPORATED SPECIAL NOTES

1. DO NOT apply HOELON^R 3EC Herbicide pre-plant incorporated to barley, as barley damage will occur.
2. HOELON^R 3EC Herbicide may be tank mixed with Glean^R Herbicide for control of annual broadleaf weeds. HOELON^R 3EC Herbicide should be used at the 2.0 to 2 2/3 pint per acre rate while Glean^R should be used at 1/3 to 1/2 oz. per acre. Refer to the EPA-approved Glean^R label for crop rotation restrictions and weed species controlled at the above-mentioned rates. Prior to the use of Glean^R determine the soil pH by laboratory analysis using 1:1 soil:water suspension. Do not use Glean^R on soils with a pH greater than 7.9.
3. DO NOT tank mix HOELON^R 3EC Herbicide with any other herbicides for pre-plant incorporation unless specifically recommended in this label.
4. HOELON^R 3EC Herbicide may be tank mixed with liquid fertilizers. Follow label recommendations concerning rates and incorporation. Before mixing HOELON^R 3EC Herbicide with fertilizer, perform a compatibility test in a quart jar. If the materials do not readily mix, do not tank mix HOELON^R 3EC Herbicide with the fertilizer.