

PH 23

8340-40

10725

14 JAN 1993

Mr. Victor A. Dorr
Hoechst Celanese Corporation
Route 202-205, P.O. Box 2500
Somerville, NJ 08876-1258

Dear Mr. Dorr:

Subject: Bugle[®] Herbicide
EPA Registration No. 8340-40
Application Dated November 19, 1992, Request To
Amend Registration To Revise Use Directions and
Disclaimers and Your Letter Dated December 19, 1992
Re: Revised Disclaimer Statement

The proposed amendments reflected in the labeling submitted have been reviewed and are acceptable for registration under Section 3 of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended provided that you:

1. Submit required data as described as required for conditional registration of the subject product on September 27, 1991.
2. Submit five (5) printed copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of this product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (H-7505C)

Enclosure

E.Wilson:Diskette #10001a:01-12-93

BEST AVAILABLE COPY

Revised: December 19, 1992

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

BUGLE™ HERBICIDE

FOR THE SELECTIVE POSTEMERGENCE CONTROL OF MONOCOT ANNUAL AND PERENNIAL GRASSES IN WHEAT, SOYBEANS, COTTON, PEANUTS AND ACREAGE CONSERVATION RESERVE (SET-ASIDE)

ACTIVE INGREDIENT:

fenoxaprop-ethyl: (±)-ethyl 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoate..... 9.76%*

INERT INGREDIENTS:..... 90.24%**

TOTAL 100.00%

* Equivalent to 0.79 pound of active ingredient per gallon
** Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN.

DANGER

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS.
SEE ATTACHED FOLDER FOR COMPLETE DIRECTIONS OF USE AND ADDITIONAL INFORMATION.

EPA Reg. No. 8340-40

NET CONTENTS:

HOECHST CELANESE CORPORATION

Route 202-206
P.O. Box 2500
Somerville, NJ 08876-1258

14 JAN 1993
8340-40

EPA EST. NO.

.....
.....
.....
.....
.....



PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin.

Do not get in eyes, on skin, or on clothing. Wear eye goggles or face shield and impermeable rubber gloves (such as neoprene or PVC) while mixing or loading. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift. Do not contaminate arable land and/or water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not store over 100°F or below 10°F. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Empty containers should be triple rinsed into the spray tank during the spray operation. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

7/25


SOYBEANS, COTTON AND PEANUTS

APPLICATION INFORMATION:

A. **Ground Application:** Broadcast - Refer to the Rate and Grass Recommendation Chart for proper application rates. Bugle™ Herbicide should be applied in a minimum of 10 gallons of water per broadcast acre. Flat fan or hollow cone nozzles are recommended. Use a minimum pressure of 40 pounds per square inch. Under dense weed/crop canopies, high spray pressure is very important for obtaining thorough coverage; therefore, use higher spray pressure under these conditions. Spot Treatment - Bugle™ Herbicide may be applied for the control of grasses through knapsack sprayers or high-volume equipment utilizing handguns or other suitable nozzle arrangements in a 0.74% v/v solution with water (e.g., 1 quart per 34 gallons of water). Apply to actively growing grasses. Apply to the foliage of grasses on a spray-to-wet basis. DO NOT spray to the point of runoff. The spray gallonage should not exceed 25 gallons per acre. Spray coverage should be uniform and complete.

B. **Air Application:** Calibrate the spray equipment prior to use. Bugle™ Herbicide should be applied in a minimum of 5 gallons of water per broadcast acre. To get uniform spray coverage, use nozzles to provide 200-350 micron size droplets. DO NOT use raindrop nozzles. Aerial applications with this product should be made at a maximum height of 10 feet above the crop with low drift nozzles at a maximum pressure of 40 psi and wind speed not exceeding 10 mph to assure accurate application within the target area. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.




PRECAUTIONS TO BE TAKEN TO AVOID SPRAY DRIFT

Do not allow spray from aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including the following:

- Do not spray if wind speeds are or become excessive. Do not spray if the wind is 10 mph or greater. If sensitive crops or plants are down wind, extreme caution must be used under all conditions. Do not spray if winds are gusty.
- Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive crops are located nearby.
- Drift from aerial applications of the herbicide is likely to result in damage to sensitive plants adjacent to the treatment site. This damage can occur at levels below the concentrations which can be detected with chemical analysis.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
- Use the lowest number of nozzles practical with the largest possible orifice size within the specified range to obtain the minimum 5 gallon per acre spray volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.
- Reduce the volume of spray mixture by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce spray drift.
- Apply as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

TIMING OF APPLICATION:

Bugle™ Herbicide will control grasses at most growth stages; but for optimum grass control, applications should be made during periods of rapid growth. Follow the recommendations for time of application listed on the Rate and Grass Recommendation Chart. Earlier applications of Bugle™ Herbicide (before all grasses have emerged) could result in late flushes of monocot grassy weeds in the treated area.



90405

RATE AND GRASS RECOMMENDATION CHART


For use only in the following states: Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, Missouri, New Mexico, North Carolina, Oklahoma, South Carolina, Tennessee and Texas.

Recommended Rate of Bugle™ Herbicide		
Monocot Grassy Weeds Annual Grasses	Maximum Height	Maximum Height
	0.5 pt./acre (8.0 fl. oz./acre)	0.6 pt./acre (9.6 fl. oz./acre)
Giant Foxtail (<u>Setaria faberii</u>)	3"	6"
Green Foxtail (<u>Setaria viridis</u>)	3"	6"
Volunteer Corn (<u>Zea mays</u>)	10"	24"
Wild Proso Millet (<u>Panicum miliaceum</u>)	6"	10"
Johnsongrass, Seedling (<u>Sorghum halepense</u>)	6"	12"
Wild Cane/Shattercane (<u>Sorghum bicolor</u>)	6"	12"
	1.0 pt./acre (16.0 fl. oz./acre)	1.2 pts/acre (19.2 fl. oz./acre)
Barnyardgrass (<u>Echinochloa crus-galli</u>)	3"	6"
Broadleaf Signalgrass (<u>Brachiaria platyphylla</u>)	3"	6"
Fall Panicum (<u>Panicum dichotomiflorum</u>)	3"	6"
Bristle Foxtail (<u>Setaria verticillata</u>)	3"	6"
Purple Foxtail (<u>Setaria viridis</u> <u>robusta purpurea</u>)	3"	6"
Robust Foxtail (<u>Setaria viridis</u> <u>robusta alba</u>)	3"	6"
Sandbur (<u>Cenchrus incertus</u>)	3"	6"
Yellow Foxtail (<u>Setaria lutescens</u>)	2"	6"

RATE AND GRASS RECOMMENDATION CHART


For use only in the following states: Colorado, Connecticut, Delaware, Illinois, Indiana, Iowa, Kansas, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri (excluding Ripley, Butler, Stoddard, Scott, Mississippi, New Madrid, Pemiscot, and Dunklin Counties), Nebraska, New Hampshire, New Jersey, New York, North Dakota, Ohio, Pennsylvania, Rhode Island, South Dakota, Vermont, Virginia, West Virginia, and Wisconsin.

Recommended Rate of Bugle™ Herbicide		
Monocot Grassy Weeds Annual Grasses	Maximum Height	Maximum Height
	0.4 pt./acre (6.4 fl.oz./acre)	0.6 pt./acre (9.6 fl.oz./acre)
Giant Foxtail (<u>Setaria faberii</u>)	3"	8"
Green Foxtail (<u>Setaria viridis</u>)	3"	8"
Volunteer Corn (<u>Zea mays</u>)	10"	24"
Wild Proso Millet (<u>Panicum miliaceum</u>)	6"	10"
Johnsongrass, Seedling (<u>Sorghum halepense</u>)	6"	12"
Wild Cane/Shattercane (<u>Sorghum bicolor</u>)	6"	12"
	0.6 pt./acre (9.6 fl.oz./acre)	0.8 pt./acre (12.8 fl.oz./acre)
Barnyard grass (<u>Echinochloa crus-galli</u>)	3"	6"
Broadleaf Signalgrass (<u>Brachiaria platyphylla</u>)	3"	8"
Fall Panicum (<u>Panicum dichotomiflorum</u>)	3"	8"
Bristle Foxtail (<u>Setaria verticillata</u>)	3"	8"
Purple Foxtail (<u>Setaria viridis robusta purpurea</u>)	3"	8"
Robust Foxtail (<u>Setaria viridis robusta alba</u>)	3"	8"
Sandbur (<u>Cenchrus incertus</u>)	3"	8"
Yellow Foxtail (<u>Setaria lutescens</u>)	2"	8"



Recommended Rate of Bugle™ Herbicide		
Monocot Grassy Weeds Annual Grasses	Maximum Height	Maximum Height
	0.8 pt./acre (12.8 fl.oz./acre)	1.1 pts./acre (17.6 fl.oz./acre)
Jungle Rice (<u>Echinochloa colonum</u>)	3"	8"
Southwestern Cupgrass (<u>Eriochloa gracilis</u>)	3"	8"
Sprangletop (<u>Leptochloa filiformis</u>)	3"	8"
Wild oats (<u>Avena fatua</u>)	3"	8"
Witchgrass (<u>Panicum capillare</u>)	3"	8"
Wooly Cupgrass (<u>Eriochloa villosa</u>)	3"	8"
Large Crabgrass (<u>Digitaria sanguinalis</u>)	2"	8"
Smooth Crabgrass (<u>Digitaria ischaemum</u>)	2"	8"
Goosegrass (<u>Eleusine indica</u>)	2"	9"
Itchgrass (<u>Rottboellia exaltata</u>)	2"	8"
Texas Panicum (<u>Panicum Texanum</u>)	2"	8"
Wirestem muhly (<u>Muhlenbergia frondosa</u>)	2"	8"
Perrennial Grasses	1.0 pt./acre (16.0 fl.oz./acre)	
Johnsongrass from rhizomes (<u>Sorghum halepense</u>)	20"	
Johnsongrass from rhizomes (<u>Sorghum halepense</u>) (Second application if needed)	0.5 pt./acre (8.0 fl. oz./acre)	
(A timely cultivation may override the necessity for a second application.)	20"	

BEST AVAILABLE COPY



Water Volume and Spray Pressure

Ground Equipment: For the tank mix, use a minimum of 20 gallons per acre of total spray solution and a minimum pressure of 40 PSI. Use standard high pressure hollow cone or flat fan nozzles. Do not use flood nozzles.

Aerial Equipment: For tank mixes, use a minimum of 5 gallons per acre of total spray solution and a minimum pressure of 40 PSI.

Mixing: Fill the spray tank half full with water while the agitator is running. Add the recommended amount of Bugle™ Herbicide followed by the tank mix component. Then add the remaining amount of water.

Tank Mix with Basagran Herbicide

Bugle™ Herbicide should be applied at a rate of 1.0 to 1.5 pints per acre (except when specific grassy weeds are small and actively growing as shown in the following special low rate tank mix table when lower rates may be used) and Basagran Herbicide at a rate of 1.5 to 2.0 pints per acre. The choice of rates of each product and additives is dependent on the weed size and weed spectrum present. Refer to the Basagran Herbicide label to identify the proper rate and additives for control of the species and size of the broadleaf weeds present.

The Bugle™ Herbicide rates for tank mixing with Basagran Herbicide are 1.0 pint per acre when the annual grassy weeds are 1 to 3 inches tall, and 1.5 pints per acre when the annual grassy weeds are 4 to 6 inches tall or less than 2 tillers (except when the special low tank mix rates are applicable). DO NOT use this tank mix if the annual grassy weeds have developed more than 2 tillers or are larger than 6 inches tall. For the control of shattercane 6 to 12 inches tall, volunteer corn 10 to 24 inches tall, and broadleaf weeds that are on the Basagran Herbicide label, tank mix Bugle™ Herbicide at a rate of 1.0 pint per acre with Basagran Herbicide at 1.5 to 2.0 pints per acre. DO NOT use this tank mix to control rhizome Johnsongrass.

Sequential applications (instead of a tank mix application) of Bugle™ Herbicide and Basagran Herbicide may be necessary if the stages of the grass and broadleaf weeds are not within tank mix label recommendations at the same time.

111
222
333
444
555
666
777
888
999
000

150405

Special Low Rates for Tank Mixing Bugle™ Herbicide with Basagran Herbicide when Specific Annual Grassy Weeds are Actively Growing and at the Stage of Growth Listed Below

Species	Maximum Height	Bugle Rate When Tank Mixed with Basagran **	
		Pts./A	Fl.Oz./A
Giant Foxtail	1-3"	0.6	9.6
Green Foxtail	1-2"	0.6	9.6
Seedling Johnsongrass	2-6"	0.6	9.6
Shattercane	4-10"	0.6	9.6
Volunteer Corn	2-10"	0.6	9.6
Wild Proso Millet	2-6"	0.6	9.6
Barnyardgrass	1-3"	0.8	12.8
Sandbur	1-3"	0.8	12.8
Wild Oats	1-3"	1.1	17.6
Wirestem Muhly	1-3"	1.1	17.6

* These rates are recommended only for the annual grass species listed in the above chart and when the grasses are small and actively growing.

Tank Mix with Blazer Herbicide

Bugle™ Herbicide should be tank mixed at a rate of 1.5 pints per acre. Blazer Herbicide should be tank mixed at a rate of 1.5 to 2.0 pints per acre. In no instances should crop oil concentrate or a surfactant be used with this tank mix. This tank mix should not be used for the control of rhizome Johnsongrass.

The stage of growth of both the annual grassy weeds and the broadleaf weeds should conform to the directions on each product label. The tank mix of Bugle™ Herbicide plus Blazer Herbicide should not be applied after the annual grasses have begun tillering. Whenever the grass and broadleaf weeds are not in the proper stage of growth according to this tank mix label, a sequential application should be utilized. When Bugle™ Herbicide is applied first, a waiting period of 3 days is necessary before applying Blazer Herbicide. When Blazer Herbicide is applied first, a waiting period of 7 days is necessary before applying Bugle™ Herbicide.

Special Note: The mixture of Bugle™ Herbicide plus Blazer Herbicide may only suppress velvetleaf, as additives cannot be used with this tank mix.

Tank Mix with Reflex 2LC Herbicide (Soybeans Only)

METHOD 1

Tank Mix Application


Bugle™ Herbicide and Reflex 2LC Herbicide

(Annual grasses and broadleaf weeds are at the proper stage of growth for treatment as per the respective labels.)

A tank mix of Bugle™ Herbicide and Reflex 2LC Herbicide may be applied at the recommended rates and growth stages to susceptible annual grass and broadleaf weed species in a manner consistent with respective labels. Bugle™ Herbicide should be applied at a rate of 1.0 - 1.5 pints per acre and Reflex 2LC Herbicide at 1.0 - 1.5 pints per acre. The choice of rates for Bugle™ Herbicide is dependent on the weed size and weed spectrum present. The Bugle™ Herbicide rate for tank mixing with Reflex 2LC Herbicide is 1.0 pint per acre when annual grassy weeds are 1-3 inches tall and 1.5 pints per acre when annual grassy weeds are 4-6 inches tall or less than 2 tillers. For the control of shattercane 6-12 inches tall, tank mix Bugle™ Herbicide at a rate of 1.0 pint per acre with Reflex 2LC Herbicide. The choice of rates for Reflex 2LC Herbicide is dependent on the weed size, weed spectrum and geographical locations. Refer to the Reflex 2LC Herbicide label for specific rates, geographical restrictions, and for a list of weeds controlled.

Use crop oil concentrate in the tank mix at 1% v/v (1 quart per 25 gallons of spray solution.)

- * DO NOT use this tank mix if perennial grasses such as rhizome Johnsongrass are the predominant grass species to be controlled.
- * DO NOT use this tank mix if the annual grassy weeds have developed more than 2 tillers or are larger than 6 inches tall as reduced annual grass control will occur.



17925

METHOD 2

Sequential Application

Bugle™ Herbicide followed by Reflex 2LC Herbicide

(Annual and/or perennial grass weeds are at the proper growth stage for treatment, prior to broadleaf weed treatment.)

Apply Bugle™ Herbicide to annual and/or perennial grass weeds at the recommended rate and growth stage listed on this label.

When treating annual grass weeds, allow at least 3 days, and when treating perennial grass weeds allow at least 5 days to elapse prior to a Reflex 2LC Herbicide application. After the appropriate time interval has elapsed, apply Reflex 2LC Herbicide with an approved adjuvant to actively growing weeds at the recommended rate and growth stage. Refer to the Reflex 2LC Herbicide label for specific rates, geographical restrictions, and for a list of weeds controlled.

METHOD 3

Sequential Application

Reflex 2LC Herbicide followed by Bugle™ Herbicide

(Broadleaf weeds are at the proper growth stage for treatment, prior to annual and/or perennial grass weed treatment.)

Apply Reflex 2LC Herbicide with an approved adjuvant to susceptible broadleaf weeds at the recommended rate and growth stage listed on the Reflex 2LC Herbicide label. Refer to the Reflex 2LC Herbicide label for specific rates, geographical restrictions, and for a list of weeds controlled.

A sequential application of Bugle™ Herbicide may be made following a Reflex 2LC Herbicide application when annual or perennial grass weeds resume active growth indicated by the development of a new leaf. Follow the recommended rates and growth stages listed on the Bugle™ Herbicide label.

SPECIAL NOTE: Tank mix applications can result in increases in crop injury as compared to either product used alone.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

BEST AVAILABLE COPY

180725

Tank Mix with Pinnacle Herbicide (Soybeans Only)

Bugle™ Herbicide may be tank mixed with Pinnacle Herbicide for broader spectrum weed control. The application rate for Pinnacle Herbicide is 0.25 ounce per acre. Refer to the rate chart below for the Bugle™ Herbicide use rates. This tank mix application should be made to grasses in the 2-leaf to 2-tiller stage of growth. Sequential applications may be necessary if the stages of growth of the grass and broadleaf weeds are not within the recommended timing of application. It is recommended that a non-ionic surfactant at a rate of 0.125-0.250% v/v be added to this tank mix. DO NOT exceed 1 pint of crop oil concentrate per acre. Refer to the Pinnacle Herbicide label for additional information.

Species	Maximum Height	Bugle™ Herbicide Rates (Pts./A) When Tank Mixed with Pinnacle Herbicide (0.25 oz.)
Giant Foxtail	5"	0.7 pt.
Volunteer Corn	24"	0.7
Seedling Johnsongrass	10"	0.7
Green Foxtail	4"	0.7
WildProso Millet	10"	1.0
Shattercane	12"	1.0
Barnyardgrass	8"	1.0
Wooly Cupgrass	6"	1.0
Yellow Foxtail	4"	1.0
Sandbur	6"	1.0

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

BEST AVAILABLE COPY

Tank Mix with Pinnacle Herbicide Plus Classic Herbicide (Soybeans Only)

Bugle™ Herbicide may be tank mixed with Pinnacle Herbicide plus Classic Herbicide for broader spectrum weed control. The application rate for Pinnacle Herbicide is 0.25 ounce per acre plus Classic Herbicide at 0.25 ounce per acre. Refer to the rate chart below for the Bugle™ Herbicide use rates. This tank mix application should be made to grasses in the 2-leaf to 2-tiller stage of growth. Sequential applications may be necessary if the stages of growth of the grass and broadleaf weeds are not within the recommended timing of application. Refer to the Pinnacle Herbicide and Classic Herbicide labels for additional information.

Species	Maximum Height	Bugle™ Herbicide Rates (Pts./A) When Tank Mixed with Pinnacle Herbicide (0.25 oz.)
Giant Foxtail	5"	1.0 pt.
Volunteer Corn	24"	1.0
Seedling Johnsongrass	10"	1.0
Green Foxtail	4"	1.0
Wild Proso Millet	10"	1.1
Shattercane	12"	1.1
Barnyardgrass	8"	1.2
Wooly Cupgrass	6"	1.2
Yellow Foxtail	4"	1.2
Sandbur	6"	1.2

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

20 1/25

Tank Mix with Pursuit Herbicide (Soybeans only)

A tank mix of Bugle™ Herbicide and Pursuit Herbicide may be applied for annual grass and broadleaf weed control in soybeans. The recommended rate for Pursuit Herbicide is 4 oz./A when the broadleaf weeds are actively growing and before they exceed a height of 3 inches. Refer to the following rate chart for the Bugle™ Herbicide rates recommended for this tank mix. It is recommended that an EPA approved nonionic surfactant at a rate of 0.25% v/v be added to this tank mix or add 28% nitrogen solution at 1-2 quarts per acre with 1 quart of COC per acre.

Species	Maximum Height	Bugle™ Herbicide Rate When Tank Mixed with Pursuit Herbicide	
		Pts./acre	Fl. Oz./acre
Giant Foxtail	6"	0.6	9.6
Volunteer corn	24"	0.6	9.6
Wild Proso Millet	10"	0.6	9.6
Seedling Johnson Grass	10"	0.6	9.6
Shattercane	12"	0.6	9.6
Green Foxtail	6"	0.8	13.0
Barnyardgrass	6"	1.0	16.0
Sandbur	6"	1.0	16.0
Wild Oats	6"	1.2	19.2
Wirestem Muhly	6"	1.2	19.2

When the annual grassy weed species and the broadleaf weeds are not in the proper growth stage for this tank mix treatment, a sequential application of Bugle™ Herbicide and Pursuit Herbicide is recommended.

Tank Mix with Fusilade 2000 Herbicide

For improved control of Johnsongrass and/or annual grass, Bugle™ Herbicide may be tank mixed with Fusilade 2000 Herbicide. When rhizome Johnsongrass is the predominant grassy weed to be controlled, Bugle™ Herbicide should be applied at a rate of 7.0 fl.oz. per acre and Fusilade at 12.0 fl.oz. per acre. When annual grasses are the predominant grassy weeds, Bugle™ Herbicide should be applied at a rate of 8.2 fl.oz. per acre and Fusilade 2000 at 9.6 fl.oz. per acre. Tank mix applications are to be used only if both annual grass and rhizome Johnsongrass are at the proper stage of growth as specified on each respective label.

For all ground applications, use crop oil concentrate in the tank mix at 1% v/v (1 quart per 25 gallons of spray solution) or a nonionic surfactant at 1/4%-1/2% v/v. For aerial applications, use crop oil concentrate at 1 pint per acre.

2/24/25


The tank mix of Bugle™ Herbicide plus Fusilade 2000 Herbicide may be applied in combination with Reflex 2LC, Basagran, or Blazer Herbicides as described on this and the Fusilade 2000 Herbicide labels. If there are any differences in labeling, the most restrictive labeling applies. DO NOT tank mix with Reflex 2LC, Basagran, or Blazer Herbicides when rhizome Johnsongrass is the predominant weed problem.

SPECIAL NOTES FOR SOYBEANS, COTTON AND PEANUTS

- 1. Annual ryegrass (Lolium sp.), quackgrass (Agropyron repens) and Bermudagrass (Cynodon dactylon) are not controlled by Bugle™ Herbicide applied alone.
- 2. Rainfall within one hour of an application may cause a reduction in grass control.
- 3. If a new flush of grass occurs, either a timely cultivation or a second application of Bugle™ Herbicide may be necessary.

DO NOT make more than two applications of Bugle™ Herbicide per growing season and DO NOT apply more than 1.5 pints per acre per growing season.

- 4. DO NOT cultivate within four days before or after a Bugle™ Herbicide application.
- 5. ALWAYS clean sprayer thoroughly before and after any pesticide application.
- 6. As a spot treatment, apply Bugle™ Herbicide in a 0.74% v/v solution with water (e.g., 1 quart per 34 gallons of water). (See instructions for use in the Ground Application Section.)
- 7. Bugle™ Herbicide is not phytotoxic to soybeans, cotton or peanuts at any growth stage. But for best results, it should be applied according to the development of the annual grassy weeds as noted in the rate and grass recommendation chart.
- 8. DO NOT graze or feed treated forage, hay, straw, or vines.
- 9. Application of Bugle™ Herbicide to grasses under stress (e.g., drought), may result in reduced control. Foxtail that have rolled leaves (onion leaf) should not be sprayed until soil moisture improves.
- 10. DO NOT plant any rotational crop in a Bugle™ Herbicide treated field for 30 days after application.



BEST AVAILABLE COPY

20935

- 11. DO NOT apply Bugle™ Herbicide less than 90 days before harvesting soybeans, 60 days before harvesting peanuts, and 40 days before harvesting cotton.
- 12. DO NOT apply this product through any irrigation system.
- 13. The application of any pesticide (other than those listed on this label) made within 7 days of the Bugle™ Herbicide application causing stress to the target grass may reduce the effectiveness of the Bugle™ Herbicide application.
- 14. Read and follow restrictions and limitations on the Basagran Herbicide, Reflex 2LC Herbicide, Blazer Herbicide, Pinnacle Herbicide, Classic Herbicide, Pursuit Herbicide, and Fusilade 2000 Herbicide labels as applicable. The most restrictive labeling applies in tank mixes.
- 15. When spraying no-till soybeans or dense weed stands, use 20 gallons of spray solution per acre.


WHEAT

Bugle™ Herbicide is a postemergence herbicide for the control of green and yellow foxtail (pigeongrass), volunteer and wild millet species, and wild oats in hard red spring wheat and winter wheat. When applying to wheat, Bugle™ Herbicide MUST be tank mixed with MCPA Herbicide. The rate of MCPA Herbicide (4 lb./gal. formulation) is 12 fl.oz. per acre. Failure to tank mix Bugle™ Herbicide with MCPA Herbicide will cause injury to the wheat. DO NOT apply to durum wheat or barley.

APPLICATION INFORMATION

Bugle™ Herbicide, when applied as recommended in the following Rate Recommendation Chart, controls the listed annual grassy weeds.

Ground Application: DO NOT apply when winds are gusty, or when conditions will favor movement of spray particles off the desired spray target. To avoid drift and insure consistent weed control, apply Bugle™ Herbicide with the spray boom as low as possible while maintaining a uniform spray pattern. Apply a minimum of 10 gallons of spray solution per acre, using a spray pressure of 30-40 psi with flat fan nozzle tips. DO NOT apply with hollow cone type insecticide or other nozzles that produce a fine droplet spray. Under dense weed/crop canopies, high spray volume (15-20 gpa) is very important for obtaining thorough coverage.



RATE RECOMMENDATION CHART					
Products	Green Foxtail	Foxtail Millets	Yellow Foxtail	Proso Millet	Wild Oats
	Fluid Ounces of Bugle™ Herbicide Per Acre				
Bugle plus MCPA	6	6	6	6	15.5
Bugle plus MCPA plus Banvel®	6	6	6	6	*
Bugle plus MCPA plus Buctril®	6	7.6	7.6	7.6	
Bugle plus MCPA plus Ally®	6	6	7.6	7.6	*
Bugle plus MCPA plus Harmony Extra®	5	6	7.6	7.6	15.5
Bugle plus MCPA plus Express®	6	6	7.6	7.6	15.5

* These tank mixes are not recommended for wild oat control.

TIMING OF APPLICATION

The time of application is determined by the growth stage of the wheat and the grassy weed species. Apply Bugle™ Herbicide to wheat from the 3-leaf stage to the end of tillering stage of growth. DO NOT spray after wheat jointing begins. Bugle™ Herbicide will control susceptible grassy weeds from the 2-leaf stage to the 2-tiller stage of growth. Applications should be made to young, actively growing weeds. Bugle™ Herbicide has no effect via the soil and will only control emerged grassy weeds.

TANK MIX COMBINATIONS

Bugle™ Herbicide may be tank mixed with several broadleaf herbicides for annual grass and broadleaf weed control in hard red spring wheat and winter wheat only. Refer to the Rate Recommendation Chart for the approved broadleaf tank mix partners and the correct rate of Bugle™ Herbicide. The maximum use rates for the broadleaf herbicides are as follows:

- Banvel Herbicide 2.0 fl.oz./A
- Banvel SGF Herbicide 4.0 fl.oz./A
- Harmony Extra 75DF 0.3 oz./A
- Express 60DF 0.17 oz./A
- Ally 6CDF 0.1 oz./A
- *Buctril 2EC 16 fl.oz./A

* The addition of Buctril 2EC Herbicide to Bugle™ Herbicide may cause temporary wheat yellowing and/or stunting.

Refer to the specific product labels for all use and precautionary statements.

2/10/25

MOISTURE EFFECTS ON ANNUAL GRASSY WEED CONTROL

Foxtail (pigeongrass) can be controlled well over a wide range of soil moisture conditions. However, foxtail under drought stress exhibits rolled leaves ("onion leaf") and should not be sprayed as poor control may result.

NOTES FOR HARD RED SPRING WHEAT AND WINTER WHEAT

1. DO NOT apply to the following crops: durum wheat, barley, rye or oats.
2. Rainfall within 1 hour of application may cause a reduction in weed control.
3. DO NOT apply more than 1 application of Bugle™ Herbicide in a growing season to hard red spring wheat or winter wheat.
4. DO NOT apply Bugle™ Herbicide within 70 days of harvesting wheat.
5. DO NOT apply this product through any type of irrigation system.
6. DO NOT apply this product aerially.
7. DO NOT tank mix Bugle™ Herbicide with other broadleaf herbicides, surfactants, or liquid fertilizers unless specifically recommended on this label.
8. Cool, wet conditions at the time of application may result in temporary stunting or chlorosis of the wheat.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

25 405

ACREAGE CONSERVATION RESERVE (SET-ASIDE)

Bugle™ Herbicide may be used to control annual grassy weeds in acreage conservation reserve (set-aside) acres. This acreage is often seeded to the following cover crops: clover, alfalfa, tall fescue, bromegrass, and ryegrass. Special note: Timothy and orchardgrass are sensitive to Bugle™ Herbicide. The cover crops listed above have excellent tolerance to Bugle™ Herbicide at 12-16 fl.oz. per acre. Select the proper rate from the Rate and Grass Recommendation Chart found in the Soybean Section of this label.

SPECIAL NOTES:

1. DO NOT harvest or graze cover crops treated with Bugle™ Herbicide.
2. DO NOT apply to cover crops such as oats, sorghum, sudangrass, and Timothy as injury may occur.

Ally® is a registered trademark of E.I. duPont de Nemours & Co., Inc.

Banvel® and Banvel® SGF are registered trademarks of Sandoz Crop Protection Corporation.

Basagran® is a registered trademark of BASF AG.

Blazer® is a registered trademark of BASF Corporation.

Buctril® is a registered trademark of Rhone-Poulenc Inc.

Classic® is a registered trademark of E.I. duPont de Nemours & Co., Inc.

Express® is a registered trademark of E.I. duPont de Nemours & Co., Inc.

Fusilade® 2000 is a registered trademark of ICI Americas Inc.

Harmony Extra® is a registered trademark of E.I. duPont de Nemours & Co., Inc.

Pinnacle® is a registered trademark of E.I. duPont de Nemours & Co., Inc.

Pursuit® Herbicide is a registered trademark of American Cyanamid Co.

Reflex® 2LC is a registered trademark of ICI Americas Inc.

File: f:\group\regscil\vd1219.923
 Date: December 19, 1992
 Edition No. 3

