

CAUTION

HARMFUL IF SWALLOWED. AVOID BREATHING SPRAY MIST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. KEEP OUT OF REACH OF CHILDREN AND PETS.

NOTE TO PHYSICIANS: Malathion, upon use, may cause cholinesterase inhibition.

ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Observe use limitations given on the label for specific crops. Do not apply during bloom when honey bees or other pollinators are active. Avoid drifts to adjoining food and forage crops or crops in bloom.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.


DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult state agricultural authorities for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

DECONTAMINATE AND DESTROY USED CONTAINERS: Dispose of containers by washing with detergent and water thoroughly. Then sell to copperage company. Do not reuse empty container. Destroy by perforating or brushing. Bury or discard in a safe place.

Avoid contamination of milk, milk equipment and water. Avoid contamination of feed and food products, also drinking fountains and feed troughs. Remove lactating animals from buildings before treating. Also remove calves under one month of age before treating.



MALATHION 57-E

EMULSIFIABLE LIQUID - AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS

Malathion (O,O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate)	57%
Aromatic Petroleum Solvent	37%
INERT INGREDIENTS	6%
	100%

APPLY THIS PRODUCT ONLY AS SPECIFIED ON LABEL

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR FURTHER CAUTIONS

NET CONTENTS GALLONS
1 5 30 55

USDA REG. NO. 8343-3

GABRIEL CHEMICALS, LTD., Robbinsville, New Jersey

Unless otherwise in water to cover one insect. First appear

ASPARAGUS As

BEANS Me

BEETS Ag

CELERY Ap

CORN CROPS (Br

Do

CORN Ag

CUCURBITS (C

Do

EGGPLANT Ap

Do

LETTUCE Ap

Do

ONIONS Thr

Do

PEAS Pec

Do

PEPPERS Ap

Do

POTATOES Ap

Ap

SPINACH Do

Do

TOMATOES Ap

Do

ALFALFA Alf

Lea

CLOVER Ap

Do

TOBACCO Ap

Do

DIRECTIONS FOR USE

Unless otherwise indicated, dosages on this label are given in pints of Malathion 57 in sufficient water to cover one acre. Thorough full coverage applications should be made beginning when insects first appear

		Dosage per acre
ASPARAGUS:	Asparagus Beetles Do not apply within 1 day of harvest	2 pts.
BEANS:	Mexican Bean Beetle, Spider Mites Do not apply within 1 day of harvest.	2 pts.
BEETS:	Aphids Do not apply within 7 days of harvest	2-3 pts.
CELERY:	Aphids, Spider Mites Do not apply within 7 days of harvest.	2 pts.
COLE CROPS:	(Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kale, Mustard Turnip): Aphids, Cabbage Looper, Imported Cabbage Worm Do not apply within 3 days of harvest on Broccoli and turnip; within 7 days of harvest on other cole crops.	2-3 pts.
CORN:	Aphids, Corn Rootworm Adults, Sap Beetles, Thrips Injury may occur in the whorl and silk stage. Do not apply within 5 days of harvest for forage use.	1½ pts.
CUCURBITS:	(Cucumber, Melon, Squash): Aphids, Spider Mites Do not apply to Cucurbits unless plants are dry. Do not apply within 1 day of harvest	2 pts.
EGGPLANT:	Aphids, Spider Mites Lace Bugs Do not apply within 3 days of harvest.	2 pts. 3 pts.
LETTUCE:	Aphids, Leafhoppers Do not apply within 14 days of harvest on leaf lettuce and within 7 days of harvest on head lettuce.	3 pts.
ONIONS:	Thrips Do not apply within 3 days of harvest on green onions.	3 pts.
PEAS:	Pea Aphid Do not apply within 3 days of harvest	2 pts.
PEPPERS:	Aphids Do not apply within 3 days of harvest.	2 pts.
POTATOES:	Aphids, Leafhoppers	2 pts.
SPINACH:	Aphids Do not apply within 7 days of harvest.	2 pts.
TOMATOES:	Aphids, Spider Mites Drosophila Flies Do not apply within 1 day of harvest.	2 pts. 2½ pts.
ALFALFA:	Alfalfa Weevil Larvae, Aphids, Grasshoppers, Potato Leafhopper, Spider Mites, Spittlebug	1 qt.
CLOVER:	Aphids, Grasshoppers, Leafhoppers Do not apply to Alfalfa or Clover or bloom or within 7 days of harvest or pasturing.	1 qt.
TOBACCO:	Aphids Do not apply to tobacco within 7 days of cutting.	1-1½ pts.

ORNAMENTALS:

Aphids, European Pine Shoot Moth, Four-lined Leaf Bug, Japanese Beetle Adult, Mealybugs, Potato Leafhopper, Rose Leafhopper, Scuffy Scale, Spider Mites, Tarnished Plant Bug, Thrips, Whitefly - use 1½ pts. per 100 gal., dilute spray (1½ teaspoonfuls per gal.).

Azalea Scale, Bagworms, Birch Leafminer, Boxwood Leafminer, Fletcher Scale, Florida-red Scale, Juniper Scale, Magnolia Scale, Oak Kermes, Pine Leaf Scale, Tent Caterpillar - use 2 pts. per 100 gal., dilute spray (2 teaspoonfuls per gal.).


For Fletcher, Juniper and Oak Kermes Scale, apply when scale crawlers have settled on foliage.

CAUTION:

Avoid use on certain ferns including Boston, Maidenhair and Pteris, as well as on some species of Crassula.

For Aerial Application, Use a minimum of 3 to 10 gallons of water per acre consistent with crop growth and good coverage.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.



E

CIDE

..... 57%
..... 37%
..... 6%
100%

EL

ONS

oy