PYBUTOX FRUIT FLY DUST

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

ACTIVE INGREDIENTS:

| Pyrethrins | 0.1% |
|--------------------------------|--------|
| Technical Piperonyl Butoxide * | 1.0% |
| INERT INGREDIENTS | 98.9% |
| TOTAL | 100.0% |

*Equivalent to .8% (butylcarbity1) (6-propylpiperony1) ether and .2% related compounds.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

DIRECTIONS FOR USE

This material, when applied with hand or power dusting equipment, will kill adult drosophila present at time of application and when thoroughly applied to the products listed below will greatly reduce egg deposition.

FRUITS: Apples, Blackberries, Boysenberries, Currants, Dewberries, Grapes, Loganberries, Peaches, Pears, Plums, Prunes (fresh).

VEGETABLES: Melons, Tomatoes - Dust containers thoroughly as soon after picking as possible. Also thoroughly dust loads after they have been assembled. Crops that are held for 20 hours after harvest should receive a second application. Consult local Agricultural Authorities for specific information as to timing, number and rates of applications which will vary with local conditions.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

ACCEPTED

SEP 141970

UNDER THE FEDERAL INSECTICIES FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NOTE TO ATTACHED COMMENTS.

USDA REG. # 8343-24

NET CONTENTS 50 LBS.

