

[8343-35]

Agrisect



LIVESTOCK SPRAY

ACTIVE INGREDIENTS:

Petroleum Distillates	99.74%
Piperonyl Butoxide Technical *	0.19%
Pyrethrins	0.07%
TOTAL	100.00%

*Equivalent to 0.152% (butylcarbityl) (6-propylpiperonyl) ether and 0.038% related compounds.

HARMFUL IF SWALLOWED

THIS PRODUCT IS TOXIC TO FISH. THIS PRODUCT IS TOXIC TO BIRDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT STORE NEAR OR CONTAMINATE FEED OR FOOD-STUFFS.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

The directions on this label are supplied by us gratuitously and are believed to be reliable and of value but are in no way guaranteed. To obtain maximum protection against flies it is essential to employ strict sanitary measures in and around the barn to destroy breeding places. Always use a good sprayer that atomizes the spray to a fine mist. Hold spray about 2 ft. away from the animal while spraying. Agrisect(R) Brand Livestock Spray is a top quality dual purpose formulation. It is designed for use as a spray to protect dairy and beef animals from the attacks of biting flies. It is also a very effective space spray and will provide fast knockdown and kill of house and other flies in the barn and milk room.

DIRECTIONS FOR USE

AS AN ANIMAL SPRAY: To protect animal against the attack of horse flies, stable flies, hornflies and mosquitoes and gnats, apply just enough spray to wet the ends of the hair but not enough to soak the hide. One or two ounces is enough to treat an adult cow. Apply once daily, preferably in the morning. Repeat at daily intervals or as necessary to give continued protection.

AS A SPACE SPRAY: To clear enclosed areas of flies, mosquitoes and gnats, spray toward the ceilings and corners until the area is filled with a light mist. For best results, close all windows and doors during spraying and keep them closed for 10 to 15 minutes after spraying. Exclude animals at the time of spraying. Do not remain in treated areas. Ventillate the area after treatment is completed.

ON HORSES: Spray lightly with a fine mist over the back, along sides, under the belly and on the legs, using two ounces per animal.

NET CONTENTS GALS.

EPA REG. NO. 8343 - 35



ROBBINSVILLE, NEW JERSEY