



MALATHION 2 - METHOXYCHLOR 2 E.C.

MULTISIFIABLE LIQUID AGRICULTURAL INSECTICIDE  
(contains 2# Malathion & 2# Methoxychlor/gallon)

ACTIVE INGREDIENTS:

Malathion *	23.8%
Methoxychlor Technical **	26.3%
Xylene	44.2%
INERT INGREDIENTS	5.7%
TOTAL	100.0%

\*O,O-dimethyl dithiophosphate of diethyl mercap-  
tosuccinate.  
\*\*Equivalent to 23.1% 2,2-bis(p-methoxyphenyl)-  
1,1,1-trichloroethane and 3.2% other isomers  
and related compounds.

**CAUTION** KEEP OUT OF THE REACH OF CHILDREN

HAZARDOUS IF SWALLOWED OR ABSORBED THROUGH SKIN.  
AVOID BREATHING SPRAY MIST. AVOID CONTACT WITH SKIN,  
CLOTHING AND CLOTHING. KEEP OUT OF REACH OF CHILDREN  
AND DOMESTIC ANIMALS. WASH THOROUGHLY WITH SOAP AND  
WATER AFTER USING. AVOID CONTAMINATION OF FEED AND  
POODSTUFFS. NOTE TO PHYSICIAN: MALATHION, UPON  
REPEATED, PROLONGED OR CARELESS USE, MAY CAUSE  
ACETYLCHOLINESTERASE INHIBITION. ATROPINE IS ANTIDOTAL.  
This product is highly toxic to bees exposed to  
direct treatment or residues on crops. Protective  
information may be obtained from your Cooperative  
Agricultural Extension Service. To protect bees do  
not apply when crops are in bloom. Remove or cover  
bees if spraying is necessary.  
Do not apply when weather conditions favor drift  
or spray from areas treated.

NET CONTENTS GALLONS

This product is toxic to fish. Keep out of lakes, ponds  
and streams.  
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. DO NOT  
STORE BELOW 32 deg. F.

Observe use limitations given on label for specific  
crops. Apply this product only as specified on label.  
The Federal Food & Drug Administration has set offi-  
cial tolerances for certain pesticide residues. In  
order that residues on food and forage crops will not  
exceed established tolerances use only at recommended  
rates and do not apply closer to harvest than speci-  
fied. Consult your local State Agricultural Extension  
Service for additional information as the timing, rate  
and number of applications needed will vary with lo-  
cal conditions.  
Rinse equipment and containers and dispose of wastes  
by burying in non-crop lands away from water supplies  
Containers should be disposed of by punching holes in  
them and burying with wastes or by burning (keep out  
of smoke).

DIRECTIONS FOR USE

ALFALFA, CLOVER: To control Alfalfa Caterpillars, Al-  
falfa Weevils (Larvae), Fall Armyworms, Flea Beetles,  
Japanese Beetles, Leafhoppers, Lygus Bugs, Spittle-Bugs,  
apply at rate of 2-3 qts./acre. Make full coverage  
application on foliage. Repeat as infestations warrant  
Do not treat during bloom. Make no applications within  
7 days of harvest.  
SOYBEANS: To control Japanese Beetles, Mexican Bean  
Beetles, Soybean Caterpillars, Velvetbean Caterpillars,  
apply at rate of 2-3 qts./acre. Make full coverage  
application on foliage. Repeat as infestations warrant  
Do not treat during bloom. Make no applications within  
7 days of harvest.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING  
OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH  
DIRECTIONS GIVEN HEREWITH.

USDA REG. # 8343-43