




MALATHION 4-PYRETHRUM 0.1 DUST



ACTIVE INGREDIENTS:	
MALATHION *	4.0%
PYRETHRINS	0.1%
INERT INGREDIENTS	<u>95.9%</u>
TOTAL	100.0%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

CAUTION **KEEP OUT OF THE REACH OF CHILDREN**

HARMFUL IF SWALLOWED. AVOID BREATHING DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS.

NOTE TO PHYSICIAN: MALATHION, UPON REPEATED, PROLONGED OR CARELESS USE, MAY CAUSE CHOLINESTERASE INHIBITION. ATROPINE IS ANTIDOTAL.

DIRECTIONS FOR USE

For control of Phorid Fly and Sciarid Fly on Mushroom Crops, dust should be applied at the rate of 2 lbs. per 60 ft. single house or 1 lb. per 1000 sq.ft. of bed surface. Make thorough application as soon after picking as possible. Repeat applications as necessary, usually twice a week. Do not apply within 24 hours of harvest.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

NET CONTENTS 50 LBS.

USDA REG. NO. 3343-50

CABOT CHEMICALS LTD.

ROBINSVILLE, NEW JERSEY