

Agri-scent BRAND 8343-51

ACCEPTED
 DEC 21 1972
 FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 U.S. DEPARTMENT OF AGRICULTURE
 REGISTERED UNDER NO. 8343-51
 SUBJECT TO ATTACHED COMMENTS.

HEPTACHLOR 2 E

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

DIRECTIONS FOR USE

ACTIVE INGREDIENTS:

Heptachlor*	23.4%
Related Compounds	9.1%
Aromatic Petroleum Solvent	60.5%
INERT INGREDIENTS:	7.0%
TOTAL	100.0%

CORN, SOYBEANS: To control Grape Colapsis, Green June Beetle larvae, Japanese Beetle larvae, Corn Seed Maggot, White Grubs, Wireworms, apply at rate of 1 to 1½ gal. (2-3 lbs. actual) per acre prior to or at time of planting.

*Heptachlorotetrahydro-4,7-methanoindene *Ornamental*
 Quantities needed for 1 pound actual Heptachlor:
 Heptachlor 2 E 2 Quarts

TURF INSECT CONTROL: To control Asiatic Garden Beetle larvae, European Chafer larvae, Japanese Beetle larvae, White Fringed Beetle larvae, White Grubs, Wireworms, apply at rate of 1 gal. (2 lbs. actual) per acre and work into the top few inches of the soil. To control Green June Beetle, apply at rate of 1½ gal. (3 lbs. actual) per acre and wash into the top few inches of soil.

WARNING KEEP OUT OF THE REACH OF CHILDREN

WARNING

MAY BE FATAL IF SWALLOWED. DO NOT GET ON SKIN. IN CASE OF SPILAGE ON SKIN, WASH IMMEDIATELY WITH SOAP AND WATER. DO NOT BREATHE VAPOR, FUMES OR SPRAY MIST. DO NOT CONTAMINATE FEED AND FOODSTUFFS. KEEP ALL PERSONS AND BENEFICIAL ANIMALS OUT OF OPERATING AREAS. VACATED AREAS SHOULD NOT BE REENTERED FOR AN HOUR AFTER SPRAYING.

ANTIDOTE

EXTERNAL: IN CASE OF SPILAGE ON SKIN, WASH PROMPTLY WITH SOAP AND WATER.

INTERNAL: IF SWALLOWED, EMETIC OF 1 TABLESPOONFUL MUSTARD IN A TUMBLER OF WATER. CALL A PHYSICIAN.

This product will kill fish and wildlife.
 Keep out of any body of water.
 Birds feeding on treated areas may be killed.
 Apply this product only as specified on this label.

DISPOSAL OF WASTES

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of smoke).

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HERewith.

USDA REG. # 8343-51



ROBBINSVILLE, NEW JERSEY

Agriseet®
BRAND

CHLORDANE 10% DUST

ACCEPTED

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
UNDER NO. 8343-55

Jan-4-1973

DIRECTIONS FOR USE

LAWNS: For Japanese Beetle Grubs, White Grubs and Chafer Beetle Grubs apply 1 lb. to 700 sq.ft. of lawn area and water thoroughly to carry insecticide into soil. For Lawn Moths, Sod Webworms and Mole Crickets apply 1 lb. to 700 sq.ft. of lawn area and water area until thoroughly wet.

KILLS ANTS, BEETLE GRUBS, CUTWORMS AND LAWN AND GARDEN SOIL INSECTS.

ACTIVE INGREDIENTS:

Chlordane Technical*	10%
INERT INGREDIENTS	90%
TOTAL	100%

*Equivalent to 6% Octachloro-4,7-Methanotetrahydroindane and 4% related compounds.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED. CONTACT WITH SKIN CAN CAUSE TOXIC SYMPTOMS. AVOID BREATHING DUST. IN CASE OF CONTACT WITH SKIN, WASH WITH SOAP AND WATER. KEEP CHILDREN AND PETS OFF TREATED AREA UNTIL MATERIAL IS WASHED INTO THE SOIL BY SPRINKLING OR RAIN AND TREATED AREA IS COMPLETELY DRY.

AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT CONTAMINATE DRINKING WATER OR USE NEAR FISH PONDS.

DO NOT FEED TREATED FORAGE (PEA VINE) TO DAIRY OR BEEF CATTLE. DISPOSE OF CONTAINER PROMPTLY WHEN EMPTY.

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER.

BIRDS FEEDING ON TREATED AREAS MAY BE HARMED.

FLOWERS AND ORNAMENTALS: Dust all plant surfaces in the early morning or evening when air is still. Use to control Ants, Leaf Miners, Leaf Rollers, Thrips, Plant Bugs, Leaf Beetles, Lace Bugs, Japanese Beetles (adults) and certain Caterpillars. For Taxus Weevils treat lower branches and soil around base of plants. Use 1 lb. to 700 sq.ft. of soil around plants for Earwigs, Crickets and Mole Crickets. Repeat as necessary.

VEGETABLE GARDEN: Work into top few inches of soil around peppers and celery at time of planting, using 6 oz. per 1000 sq.ft. (25 lbs./acre) for Wire Worm. For Flea Beetles and Hornworms on tomatoes, Flea Beetles on beans, black-eye peas, eggplant and cabbage, Cucumber Beetles, Squash Bugs and Squash Vine Borers on cucumbers, melons and squash, dust 4-6 oz. per 1000 sq.ft. Repeat as necessary. To avoid occasional injury to cucumbers, melons and squash do not apply when plants are wet with dew. Do not apply to any of the above plants after fruit, pods or edible parts begin to form.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HERewith.

EPA REG. NO. 8343-55

NET WEIGHT 50 LBS.



ROBBINSVILLE, NEW JERSEY