



HEPTACHLOR 5% GRANULAR

FEB 5 1977

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. BIRDS FEEDING ON TREATED AREAS MAY BE KILLED. KEEP OUT OF ANY BODY OF WATER.

DIRECTIONS FOR USE

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL

FOR TURF INSECT CONTROL: For the control of ants, Asiatic Garden Beetle Larvae, European Chafer Larvae, Japanese Beetle Larvae, White Fringed Beetle Larvae, White Grubs and Wireworms apply 40 lbs of Heptachlor 5% Granular per acre. Wash into the top few inches of soil. For the control of Green June Beetle apply 60 lbs of Heptachlor 5% Granular per acre. Wash into the top few inches of soil. Use only on ornamental turf. Birds feeding on treated areas may be killed.

Consult your State Experiment Stations for local recommendations.

DO NOT RE-USE EMPTY CONTAINER. DESTROY BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREON.

ACTIVE INGREDIENTS:

Heptachlor*.....	5.00%
Related Compounds.....	1.94%
INERT INGREDIENTS.....	93.06%
<hr/>	
TOTAL	100.00%

*Heptachlorotetrahydro-4,7-methanoindene

WARNING KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED. AVOID INHALATION. IRRITATED OR PROLONGED CONTACT WITH SKIN CAN CAUSE TOXIC SYMPTOMS. IN CASE OF CONTACT WITH SKIN WASH WITH SOAP AND WATER. AVOID CONTAMINATION OF FEED AND EGGS/PUFFS. IF SWALLOWED, GULP 1 TABLESPOON SALT IN A QUARTER OF WATER. CALL A PHYSICIAN.

DO NOT USE THIS PRODUCT ON DAIRY ANIMALS, IN ANY PARTS OF OR FEED OR FORAGE TO BE FED TO ANY ANIMALS OR OTHER ANIMALS BEING BREEDING OR BEARING. DO NOT USE IN OR AROUND BIRD FEEDERS. DO NOT USE ON HEAVILY SEEDED TREES. KEEP FEEDERS AND TREES OUT OF TREATED AREAS OR TURF WILL BECOME UNUSABLE AND WASHED AWAY SOON BY WINDING OR RAIN.

net contents 50 pounds

USDA REG. # 8313-63

Aspirin (R) BRAND

Ornamental Turf

ROBINSVILLE, NEW JERSEY