

# AGRISECT

CUTWORM BAIT  
CONTAINS 5% TOXAPHENE

ACCEPTED

MAY 3 1 1967

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 8343-62 SUBJECT  
TO ATTACHED COMMENTS.

## INGREDIENTS

ACTIVE INGREDIENT:

\* TOXAPHENE..... 5%

INERT INGREDIENTS\*\* .....95%

\* TECHNICAL CHLORINATED CAMPHENE (CHLORINE CONTENT.....67% TO 69%)

\*\* APPLY POMACE

CUTWORM BAIT IS AN IMPREGNATED INSECTICIDE BAIT CONTAINING 5% CHLORINATED CAMPHENE. IT IS DESIGNED FOR THE FOLLOWING USAGE: NOTE CAUTIONS AND READ DIRECTIONS BEFORE USING.

CUTWORMS: TOBACCO, TOMATOES, BEANS - DISTRIBUTE BAIT WHEN FIRST SIGNS OF CUTWORMS ARE NOTED. SCATTER BAIT ALONG ROWS OF PLANTS OR BROADCAST UNIFORMLY IN FIELD. BAITING IN LATE AFTERNOON IS PREFERRED. FOR BROADCAST APPLICATION. USE 30 TO 40 LBS. PER ACRE.

## CAUTION

KEEP OUT OF THE REACH OF CHILDREN

CHLORINATED CAMPHENE IS TOXIC AND IS ABSORBED THROUGH THE SKIN IN THE DRY OR WET FORM. AVOID SKIN CONTACT. WASH SKIN THOROUGHLY WITH SOAP AND WATER IMMEDIATELY AFTER CONTACT OR USE. KEEP CLOTHING FREE FROM RESIDUES. REMOVE AND WASH CONTAMINATED CLOTHING. AVOID INHALING DUST OR GETTING IT IN THE EYES. USE RESPIRATOR AND GOGGLES IF NECESSARY FOR PROTECTION. KEEP AWAY FROM DOMESTIC ANIMALS. HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

DO NOT USE ON CUCURBITS. DO NOT USE ON YOUNG PLANTS. DO NOT USE ON FORAGE CROPS TO BE FED TO DAIRY ANIMALS OR TO ANIMALS BEING FINISHED FOR SLAUGHTER. DO NOT APPLY TO FOOD OR FEED CROPS UNDER CONDITIONS RESULTING IN EXCESSIVE RESIDUE AT HARVEST OR FORAGE TIME. DO NOT APPLY TO GREEN OR SNAP BEANS AFTER PODS BEGIN TO FORM. DO NOT APPLY TO TOMATOES AFTER THE FRUIT BEGINS TO FORM.

TOXIC TO FISH AND WILDLIFE. DO NOT CONTAMINATE STREAMS, LAKES OR PONDS WITH THIS DUST. BAIT EXPOSED ON SOIL SURFACE MAY BE HAZARDOUS TO BIRDS AND WILDLIFE.

NET WEIGHT

POUNDS

**AGRISECT**  
CHEMICAL CORPORATION

U.S.D.A. REG. NO. 8343-

ROBBINSVILLE, NEW JERSEY

• NEW YORK, NEW YORK