

AGRISPECT

NEMAGON 10 G
FOR USE AS A SOIL FUMIGANT FOR CONTROL OF ROOT-LESION AND ROOT-KNOT NEMATODES IN THE SOIL.

ACTIVE INGREDIENTS:

1,2-Dibromo-3-Chloropropane.... 9.5%
Other Halogenated C₃ Compounds. 0.5% *
INERT INGREDIENTS..... 90.0%
TOTAL.....100.0%

*Equivalent to 10% Namagon Soil Fumigant.

GENERAL DIRECTIONS

Nemagon is a soil fumigant material for the control of nematodes in the soil to decrease root or plant damages. Use this material to control Pin, Root Knot, Meadow, Dagger, Stem and bulb, Ring and Stubby root nematodes.

Land to be treated for Nematodes should be plowed to a depth of 8 to 10 inches 4 weeks before treatment to allow plant material to partially decompose. A plow-down application of fertilizer high in Nitrogen will speed up the decomposition of the organic matter.

Apply material when soil temperature at 6 inches is 40° to 20°F. Soil should be in good tilth and contain adequate moisture for good seed germination. Extremely wet or dry soils should not be treated.

10% Nemagon Granules may be mixed with fertilizer to facilitate application.

Use only on mineral soils.

PREPLANTING. Over-all or broadcast treatment, use grain drill or other means to apply in bands approximately 12 inches apart to a depth of 6 to 8 inches. Seal the chisel channels with pressure roller or flood with water. Draggings or cultipacking soil behind this application will help to seal gases in the soil.

PREPLANTING ROW TREATMENT: Apply in rows to be treated to a depth of 6 to 8 inches. Seal chisel channels with a press wheel or drag. Adjust chisels to same spacing as planter.

FOR ALL PREPLANTING TREATMENTS WAIT 7 TO 14 DAYS BETWEEN APPLICATION AND PLANTING. For deep rooted plants wait 4 to 6 weeks before planting treated area.

ESTABLISHED PLANTS: Apply as bands 12 inches apart 6 to 8 inches deep as close to plants as possible. For treatment of trees, treat area where feeder roots are found. Seal chisel channels.

When treating growing annuals, apply as a band 4 inches deep as close to plants as possible without disturbing or damaging the roots.

VEGETABLE CROPS

Snap Beans, Carrots, Celery, Parsnips, Radishes and Turnips, Cabbage, Cauliflower, Broccoli, Brussels Sprouts, etc.

Preplanting—Over-all or Broadcast Treatment: Apply 170 to 240 lbs. per acre.

Preplanting—Row Treatment: Apply 85 to 120 lbs. per acre for 24 inch rows. Apply 45 to 60 lbs. per acre for 48 inch rows.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

STORAGE PRECAUTIONS: DO NOT STORE NEAR PLANTS, SEEDS OR FERTILIZERS. STORE LEFTOVER GRANULES IN ORIGINAL CONTAINER OR IN GLASS CONTAINER WITH TIGHT-FITTING LID TO PREVENT EVAPORATION OF THE NEMAGON SCIL.

CAUTION KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED. May cause irritation of skin, eyes, nose and throat. Avoid contact with skin, eyes or clothing. Avoid prolonged breathing of vapors or dust. Wash thoroughly with soap and water after handling and before eating or smoking. Keep children and pets off treated areas until this material has been washed into soil. Birds feeding in treated areas may be harmed.

This product toxic to fish and wildlife. Keep out of lakes, streams and ponds.

THIS PESTICIDE IS TO BE SOLD ONLY IN THIS ORIGINAL UNBROKEN PACKAGE.

Lettuce: Preplanting. Over-all treatment. Apply 160 to 170 lbs. per acre.

Wait 7 to 14 days before planting. Do not feed vines of Beans to Livestock.

Tomatoes (Transplants): Preplanting Over-all treatment. Allow at least 14 days between treatment and planting. Apply 100 to 130 lbs. per acre.

FRUIT AND VINE CROPS

Strawberries: Allow at least 300 days between treatment and harvest.

Preplanting or postplanting over-all or broadcast treatment. Apply 170 to 240 lbs. per acre. Apply 85 to 120 lbs. per acre for 24 inch rows.

Row application. Apply in soil 6 inches deep and 6 inches from each side of the plant growing in the row.

Cucumbers, Cantalope, Honeydew Melons, Muskmelons and Summer Squash. Preplanting or at time of planting application to mineral soils. For preplanting wait 7 to 14 days before planting. Apply 170 to 240 lbs. per acre. No post-planting applications.

TURF AND LAWNS

Established Grasses: Bermuda Grass, Chewink, Fescue, Colonial Bent, Kentucky Bluegrass and Zoysia. Apply

500 lbs. of Nemagon 10G per acre or 12 lbs. per 1000 sq. ft. May be applied with a fertilizer spreader or shaker can. Wet down thoroughly with overhead irrigation immediately after application. Keep children and pets off treated areas until this material is washed into the soil.

ORNAMENTALS: Apply as outlined under Preplanting: Over-all or broadcast treatment.

Rate: Apply at rate of 430 to 850 lbs/acre. Wait 7 to 14 days before planting to:

| | | |
|------------|--------------------|--------------|
| Agave | Funkia | Pansy |
| Anchusa | Gypsophillum | Peony |
| Arbutus | Heliosia | Philodendron |
| Aspidistra | Heliborus | Poppy |
| Azalea | Hibiscus | Primula |
| Boxwood | Holly | Rudbeckia |
| Coreopsis | Lavandula | Schaffneria |
| Daisy | Lily-of-the-Valley | Sedum |
| Dianthus | Maple | Taxus |
| Dicentra | Maranta | Tritoma |
| Elm | Oak | Violet |
| | Pachysandra | |

Rate: Apply at rate of 170 lbs. per acre to:

| | | |
|---------------|------------|--------|
| Chrysanthemum | Gardenias | Roses |
| Delphinium | Gladiolus | Spirea |
| Dwarf Rose | Periwinkle | Vinca |
| Forsythia | Phlox | |

NET CONTENTS LBS.

USDA REG. NO. 8343-56

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 8343-56

JUN 1 1968

ACCEPTED

AGRISPECT
CHEMICAL CORPORATION

ROBBINSVILLE, NEW JERSEY

NEW YORK, NEW YORK