

8343-79

**ACCEPTED**

Jul 2 1968

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR REGISTERED PERSONS ASSIGNED  
TO THIS PRODUCT SUBJECT  
TO ATTACHED COMMENTS.

# AGRISECT

PARATHION 2# E.C.

**ACTIVE INGREDIENTS:**

Parathion*.....	25.42%
Xylene.....	69.58%
<b>INERT INGREDIENTS.....</b>	<b>5.00%</b>
<b>TOTAL.....</b>	<b>100.00%</b>

\*O,O-diethyl O-p-nitrophenyl thiophosphate  
Contains 2# Parathion/U.S.gallon at 70°F.



**DANGER**

POISON

KEEP OUT OF THE REACH OF CHILDREN

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. If spilled on skin wash immediately with soap and warm water. Do not breathe vapors or spray mist. Wear a mask or respirator of a type passed by the USDA for Parathion protection. Do not contaminate feed or food-stuffs. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Keep all persons out of the operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residues have dissipated.

This product toxic to fish and wildlife. Keep out of lakes, streams and ponds. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burning with wastes.

**FLAMMABLE: DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. KEEP CONTAINER CLOSED.**

**ANTIDOTE-INTERNAL:** Give tablespoon of salt in glass of warm water. Repeat until vomit fluid clear. Keep victim prone and quiet. **CALL PHYSICIAN IMMEDIATELY.**  
**EXTERNAL:** If skin contacted wash immediately with soap and water. **NOTE TO PHYSICIAN:** Therapeutic doses of atropine appear effective as antidote.

DIRECTIONS FOR USE

**SMALL GRAINS SUCH AS WHEAT, OATS, BARLEY:** To control Greenbug (aphids) apply 1 pt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. If infestation persists repeat treatment 10-14 days later. Do not apply within 15 days of harvest.

**VETCH AND ALFALFA:** To control aphids apply 1 pt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. Apply at first sign of infestation. Do not apply within 15 days of harvest.

**PEAS:** To control aphids apply 1 pt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. Apply at first sign of infestation. Do not apply within 15 days of harvest.

**CORN:** To control European Corn Borer apply 1 qt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. Apply after 20 egg masses are counted in 100 stalk samples and first eggs are in blackhead stage. Repeat at 5-7 day intervals. Do not apply within 12 days of picking or harvest as forage.

**BEANS:** To control Mexican Bean Beetle, Leafhoppers, Aphids, Red Spider, Armyworms, Leaf Rollers and Leaf Miners, apply 1 pt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. Repeat application as needed. Do not apply within 21 days of harvest.

**CELERY:** To control Aphids and Celery Worms use 1 pt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. Do not apply within 21 days of harvest.

**COTTON:** To control Cotton Leafworms, Aphids, Fleahoppers, Garden Webworm, Red Spider Mites, Thrips apply 1 pt/acre in 2 gals spray from airplane or in 5-10 gals spray in power sprayer. Apply when infestation first appears, repeat as necessary. Do not apply within 15 days of hand picking.

ROBBINSVILLE, NEW JERSEY      NEW YORK, NEW YORK

NET CONTENTS                      GALLONS

USDA REG. NO. 8343-

AGRISECT  
CHEMICAL CORPORATION