

AGRISECT

GUTHION 2% DUST
Contains 2% Guthion *

NET WEIGHT POUNDS
STORE IN A COOL DRY PLACE

GUARANTEE

*Reg. U.S. and Canadian Pat Offs. by Farbenfabriken Bayer A.G., Chemarg Corporation licensee

ACTIVE INGREDIENT:

O,O-Dimethyl S-(4-oxo-1,2,3-benzotriazin-4-ylmethyl) phosphorodithioate..... 2%

INERT INGREDIENTS..... 98%

100%

U.S. Patent No. 2,758,115--Canadian Patent No. 552,660 Effective for purpose claimed

RECOMMENDED APPLICATIONS

CROP	INSECT	POUNDS GUTHION 2%	REMARKS
<u>FRUIT</u>			
<u>Blueberries</u>	<u>Blueberry maggot, Fruitworms, Leaf miner, Lecanium scale, Plum curculio</u>	30	Do not apply more than twice per season. Note: Higher rates (up to 45 lbs. per acre) may be applied by ground or air as necessary but not within 14 days of harvest. -AIRPLANE APPLICATION ONLY: Apply specified dosage per acre. A second application may be made up to within 4 hrs of harvest. -WITH GROUND EQUIPMENT, second application can be made within 7 days of harvest.
<u>Cranberries</u>	<u>Cranberry fruitworm, Fireworms, Sparganothis sulfureana, Tipworms</u>	30 to 45	Apply specified dosage per acre. Repeat as necessary but not within 21 days of harvest.
<u>VEGETABLES</u>			
<u>Beans (Snap and Dry)</u>	<u>Leaf roller, Mites, Leafhoppers, Mexican bean beetle</u>	22 to 30	Apply specified dosage per acre. Repeat as necessary. Do not apply within 30 days of harvest for dry beans nor within 7 days of harvest for snap beans. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.
<u>Broccoli, Brussels Sprouts, Cabbager, Cauliflower</u>	<u>Aphids, Cabbage looper, Imported cabbageworm, Diamondback moth</u>	30 to 37	Apply specified dosage per acre. Repeat as necessary. Do not apply within 7 days of harvest for Brussels sprouts, 15 days of harvest for broccoli and cauliflower, not within 21 days of harvest for cabbage.
<u>Potatoes</u>	<u>Colorado potato beetle, Leaf miners, Tubeworm, Aphids, European corn borer, Flea beetle, Leafhoppers, Spittlebugs, Tarnished plant bug</u>	22 22 to 30 30 30 to 37	Apply specified dosage per acre. Repeat as necessary. Do not apply within 7 days of harvest.

AGRISECT
CHEMICAL CORPORATION

ACCEPTED
Comments
DEC 9 1968
FOR THE FEDERAL DEPARTMENT OF AGRICULTURE AND DOMESTIC AFFAIRS
FOR ECONOMIC POISON
REGISTERED UNDER NO. 8249-93

see continued sheet for additional recommended applications and Warning statement etc.

ROBBINSVILLE, NEW JERSEY • NEW YORK, NEW YORK

AGRISECT

GUTHION 2% DUST (continued)

CROP	INSECT	POUNDS GUTHION 2%	REMARKS
Tomatoes	Colorado potato beetle		Apply specified dosage per acre. Repeat as necessary.
	Green stink bug	22	
	<u>Drosophila</u>		
	Leaf miners	22 to 30	
	Aphids, European corn borer, Flea beetles		
	Hornworm, Leafhoppers, Thrips	30 to 37	
	Fruitworm, Tuberworm	37	
ORNAMENTALS			
Ornamentals, Nursery Plants, Forest & Shade Trees	Aphids, Lace bugs, Leafhoppers, Mites, Thrips	30	Apply specified dosage per acre. Dust all foliage surfaces including the underside of leaves for complete coverage. Repeat as necessary.
	European pine shoot moth Nantucker pine tip moth	27 to 37	Apply specified dosage per acre. Time applications to coincide with moth flight. Thorough coverage is necessary for best results.
Do not allow children or pets on treated area until material has been washed into soil and treated area is dry.			

RESTRICTIONS


Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues. Do not graze livestock in treated orchards or groves for 21 days after treatment. Do not treat food crops grown in the green house. Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on the label.

WARNING

KEEP OUT OF THE REACH OF CHILDREN

May be fatal if swallowed, inhaled, or absorbed through skin. Wear natural rubber gloves, protective clothing, and goggles. In case of contact wash immediately with soap and water. Do not breathe dust. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for GUTHION protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Do not store near feed and food products. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

TO PHYSICIAN - Atropine sulfate is antidotal.



AGRISECT
CHEMICAL CORPORATION

ROBBINSVILLE, NEW JERSEY • NEW YORK, NEW YORK