




CHLORDANE 25% W.P.



ACTIVE INGREDIENT	
Technical Chlordane.....	25%
INERT INGREDIENTS.....	75%
TOTAL	100%

\*Equivalent to 15% octa-chloro-4,7-methanotetrahydroindane and 10% related compounds.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust or spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not use on humans or household pets. Keep children and pets off treated areas until insecticide has been washed into soil by sprinkling or rain and area is dry. Do not store with foodstuffs. Do not reuse empty container. Destroy by burying or burning. Stay out of smoke or fumes.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

For listed Soil-Inhabiting Insects make application in sufficient water for thorough coverage and wash or work into top few inches of soil immediately. For grubbs of Japanese Beetle in Federal/State Quarantine Programs, use 40 lbs. per acre (1 lb./1,000 sq. ft.). Make application any time when ground is not frozen or soggy. Spring or Fall treatments are preferred. Treatment will kill high percentage of full-grown grubs at time of application and will control annual broods for 2 to 3 years.

For Black Vine Weevil Control in Nursery Plantings of Japanese Yew in Michigan, use 4 pounds per 100 gallons of water as a spray. Spray foliage and ground beneath branches. As pre-planting treatment, use 40 pounds per acre, spray and work into top 3 inches of soil.

NOTICE: BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

EPA Reg. No. 8343-87

EPA Est. 904-NJ-2

NET WEIGHT: POUNDS



ROBBINSVILLE, NEW JERSEY 08691