



DIRECTIONS FOR USE

USE THIS PRODUCT ONLY AS DIRECTED ON THIS LABEL

DURSBAN* ICG

Granular Insecticide for Control of Mosquito Larvae. To Be Applied Only by Trained Personnel of Public Health Organizations, Mosquito Abatement Districts, Public Mosquito Control Programs and Professional Pest Control Operators.

ACTIVE INGREDIENT:

| | |
|---|------|
| Chlorpyrifos [0,0-Diethyl 0-3,5,6-trichloro-2-pyridyl phosphorothioate] | 1% |
| INERT INGREDIENTS | 99% |
| TOTAL | 100% |

*DURSBAN(R) - Trademark of the Dow Chemical Company, U.S. Patent No. 3,244,586

CAUTION KEEP OUT OF THE REACH OF CHILDREN

MAY BE HARMFUL IF SWALLOWED. AVOID EYE CONTACT. WASH THOROUGHLY AFTER HANDLING. WASH CONTAMINATED CLOTHING BEFORE REUSE. FLUSH CONTAMINATED EYES WITH PLENTY OF WATER AND GET MEDICAL ATTENTION IF IRRITATION PERSISTS. KEEP AWAY FROM FOOD, FEEDSTUFFS AND DOMESTIC WATER SUPPLIES.

NOTE TO PHYSICIAN: ACTIVE INGREDIENT IS A CHOLINESTERASE INHIBITOR. TREAT SYMPTOMATICALLY. ATROPINE IS AN ANTIDOTE.

THIS PRODUCT IS TOXIC TO FISH, WILDLIFE AND BIRDS. FISH, SHRIMP AND CRAB MAY BE KILLED AT APPLICATION RATES RECOMMENDED ON THIS LABEL. DO NOT APPLY WHERE THESE ARE IMPORTANT RESOURCES. CONSULT YOUR STATE FISH AND GAME DEPARTMENT BEFORE USING THIS PRODUCT.

Use DURSBAN ICG granular insecticide to control mosquito larvae in non-crop areas that serve as breeding sites for these pests. Apply using a hand operated or ground vehicular mounted gravity or rotary type spreader or seeder, or with aerial application equipment that will provide uniform distribution of the granules.

Flooded Area Treatment: Apply uniformly over flood water impoundments whenever mosquito larvae are present. Use DURSBAN ICG at the rate of 2.5 to 5 lbs. per acre. This rate is equivalent to 0.025 to 0.05 lbs. of active ingredient per acre.


Pre-Hatch Treatment: Many mosquito breeding areas are not covered by water during certain months of the year, yet these areas contain mosquito eggs from which larvae will hatch and grow to maturity when flooding occurs. To control these larvae uniformly apply DURSBAN ICG at the rate of 10 lbs. per acre over the known breeding sites prior to the time flooding is expected. This rate is equivalent to 0.1 lbs. of active ingredient per acre.

DISPOSE OF EMPTY CONTAINERS AND WASTES BY BURN OR BURYING IN NON-CROP LANDS AWAY FROM DOMESTIC WATER SUPPLIES.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

NET WEIGHT LBS.

EPA REG. NO. 8343-96



ROBBINSVILLE, NEW JERSEY



DIRECTIONS FOR USE

USE THIS PRODUCT ONLY AS DIRECTED ON THIS LABEL.

DURSBAN* ICG

Granular Insecticide for Control of Mosquito Larvae. To Be Applied Only by Trained Personnel of Public Health Organizations, Mosquito Abatement Districts, Public Mosquito Control Programs and Professional Pest Control Operators.

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-Diethyl 0-3,5,6-trichloro-2-pyridyl phosphorothioate]

INERT INGREDIENTS

TOTAL

1%
99%
100%

*DURSBAN (R) - Trademark of the Dow Chemical Company, U.S. Patent No. 3,244,586

CAUTION KEEP OUT OF THE REACH OF CHILDREN

MAY BE HARMFUL IF SWALLOWED. AVOID EYE CONTACT. WASH THOROUGHLY AFTER HANDLING. WASH CONTAMINATED CLOTHING BEFORE REUSE. FLUSH CONTAMINATED EYES WITH PLENTY OF WATER AND GET MEDICAL ATTENTION IF IRRITATION PERSISTS. KEEP AWAY FROM FOOD, FEEDSTUFFS AND DOMESTIC WATER SUPPLIES.

NOTE TO PHYSICIAN: ACTIVE INGREDIENT IS A CHOLINESTERASE INHIBITOR. TREAT SYMPTOMATICALLY. ATROPINE IS AN ANTIDOTE.

THIS PRODUCT IS TOXIC TO FISH, WILDLIFE AND BIRDS. FISH, SHRIMP AND CRAB MAY BE KILLED AT APPLICATION RATES RECOMMENDED ON THIS LABEL. DO NOT APPLY WHERE THESE ARE IMPORTANT RESOURCES. CONSULT YOUR STATE FISH AND GAME DEPARTMENT BEFORE USING THIS PRODUCT.

Use DURSBAN ICG granular insecticide to control mosquito larvae in non-crop areas that serve as breeding sites for these pests. Apply using any hand operated or ground vehicular mounted gravity or rotary type spreader or seeder, or with aerial application equipment that will provide uniform distribution of the granules.
Flooded Area Treatment: Apply uniformly over flood water impoundments whenever mosquito larvae are present. Use DURSBAN ICG at the rate of 2.5 to 5 lbs. per acre. This rate is equivalent to 0.025 to 0.05 lbs. of active ingredient per acre.
Pre-Hatch Treatment: Many mosquito breeding areas are not covered by water during certain months of the year, yet these areas contain mosquito eggs from which larvae will hatch and grow to maturity when flooding occurs. To control these larvae uniformly apply DURSBAN ICG at the rate of 10 lbs. per acre over the known breeding sites prior to the time flooding is expected. This rate is equivalent to 0.1 lbs. of active ingredient per acre.

DISPOSE OF EMPTY CONTAINERS AND WASTES BY BURNING OR BURYING IN NON-CROP LANDS AWAY FROM DOMESTIC WATER SUPPLIES.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

NET WEIGHT LBS.

EPA REG. NO. 8343-96



ROBINNVILLE, NEW JERSEY

5-72
8343-9