



CAPTAN 7% DUST - WETTABLE POWDER

ACTIVE INGREDIENT:		
Captan *	.....	7%
INERT INGREDIENTS	.....	93%
	TOTAL .....	100%

ACCEPTED

*May 16, 1973*

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR FEDERAL POISON REGISTER-  
ED MATERIAL NO. 8343-107

\*N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide

**CAUTION**

**KEEP OUT OF THE REACH OF CHILDREN**

AVOID INHALATION OF DUST OR SPRAY MIST. AVOID CONTACT WITH SKIN. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING (STAY AWAY FROM SMOKE OR FUMES).

DIRECTIONS FOR USE

TURF, LAWN: Brown Patch, Leafspot, Damp-Off, Root Rot: 14 lbs. per 100 gals. of water. Apply 10 gals. to each 1000 sq.ft. of turf. Apply prior to infection. Repeat at 7-10 day intervals if necessary.

Do not graze treated areas. Do not feed clippings to livestock.

ORNAMENTALS, ROSES: Black Spot; CARNATIONS: Rust and Leaf Spot; CHRYSANTHEMUMS: Flower Blight: 14 lbs. per 100 gals. of water. Apply at first sign of diseases, cover upper and lower leaf surfaces thoroughly. Repeat every 7-10 days.

SOIL AND GREENHOUSE BENCH TREATMENTS: Preplanting for Damp-Off and Root Rot Diseases on Seedlings or Transplants of ~~Roses and~~ Other Shrubs and Trees), many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds. Apply as a dust at 2 lbs. per 1000 sq.ft. or spray at 14 lbs. per 100 gals. of water, using 15 gals. to each 1000 sq.ft. Cultivate into upper 3-4 inches of soil before planting.

*transplant*

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

NET WEIGHT 50 LBS.

EPA REG. NO. 8343-107



ROBBINSVILLE, NEW JERSEY