



NET WEIGHT

Toxamite Powder

50 LBS.

ACCEPTED
comments
 APR 18 1967
 UNDER THE FEDERAL
 FUNGICIDE AND
 FOR ECONOMIC
 ED UNDER NO. 8344-1
 TO ATTACHED COMMENT

Contains- **SEVIN** *

For Control of Poultry Pests on Chickens, Turkeys, Ducks, Geese, Game Birds and Pigeons

Active Ingredient: carbaryl (1-naphthyl N-methylcarbamate) 5% by wt.
 Inert Ingredient (s) 95% by wt.

USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS

DIRECTIONS FOR USE

Litter treatment: Apply 1 lb. per 40 sq. ft. to control Northern fowl Mite, chicken mite, lice, fleas and bed bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

Hand application to birds: Use one pound to treat 100 birds by means of a shaker can, squeeze bottle, or hand duster to control Northern fowl mite, chicken mite, lice and fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and fleas is a supplement to litter treatment for control of these pests.

Dust-bath-boxes: Use 2-1/2 pounds per box for each 50 birds to control Northern fowl mite, chicken mite, lice and fleas. This is a supplement to litter treatment for chicken mite and flea control. On ranges, 2 natural wallows may be substituted for each dust-bath-box.

Do Not apply within 7 days of slaughter.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dust. Do Not Take Internally
Skin Contact With Any Insecticide May be Harmful. Avoid Unnecessary Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles
KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

WARRANTY

The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, of toxicity to plants and to laboratory animals and of residue remaining on food crops.

USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS

DIRECTIONS FOR USE

Litter treatment: Apply 1 lb. per 40 sq. ft. to control Northern fowl Mite, chicken mite, lice, fleas and bed bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

Hand application to birds: Use one pound to treat 100 birds by means of a shaker can, squeeze bottle, or hand duster to control Northern fowl mite, chicken mite, lice and fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and fleas is a supplement to litter treatment for control of these pests.

Dust-bath-boxes: Use 2-1/2 pounds per box for each 50 birds to control Northern fowl mite, chicken mite, lice and fleas. This is a supplement to litter treatment for chicken mite and flea control. On ranges, 2 natural wallows may be substituted for each dust-bath-box.

Do Not apply within 7 days of slaughter.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dust. Do Not Take Internally. Skin Contact With Any Insecticide May be Harmful. Avoid Unnecessary Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidote a.l. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

WARRANTY

The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, of toxicity to plants and to laboratory animals and of residue remaining on food crops.

The manufacturer makes no other express or implied guarantee, warranty of representation whatever of a similar or different sort, and none is to be inferred by the purchaser or user. No modification of this paragraph is authorized, except by specific reference to this paragraph in writing by an employee of the manufacturer.

*Trade mark of Union Carbide Corporation.



USA Registration No. 6344-1

PROGRESS CHEMICAL CO.

CANTON, GEORGIA

