

Net wt.

SANITIZING RINSE FOR COMMERCIAL EGGS

Use a dilution that will give 50 p.p.m. to 60 p.p.m. available chlorine. Eggs should be sprayed with a fine mist of sanitizing solution maintained at a temperature 5° F. higher than wash solution at the final spray section of the washer. Allow eggs to dry before packing.

FOR USE WITH PROPORTIONERS THAT METER 1 OZ. PER GALLON

For 50 p.p.m. available chlorine at final rinse:
Use 6.4 oz. Sanitizer per gallon water for stock solution.
For 100 p.p.m. available chlorine at final rinse:
Use 12.8 oz. Sanitizer per gallon water for stock solution.

Make up stock solutions fresh daily.

Hot water may be necessary to get sanitizer powder into solution.

PCC CHLORINE SANITIZER POWDER

FOR USE IN SANITIZATION APPLICATIONS IN EGG PROCESSING PLANTS, POULTRY PROCESSING PLANTS, FINAL RINSE SPRAY SANITIZING OF COMMERCIAL EGGS AND OTHER APPLICATIONS WHERE CHLORINE SANITIZATION IS NEEDED.

ACTIVE INGREDIENTS:

Potassium dichloro-s-triazinetrione 22%

INERT INGREDIENTS

78%
100%


CAUTION: KEEP OUT OF REACH OF CHILDREN Harmful if swallowed. Do not take internally. If accidentally swallowed, induce vomiting and call a physician. Prolonged contact may irritate skin, eyes or nose.

USDA Registration No. 8344-6
MANUFACTURED BY



PROGRESS CHEMICAL COMPANY, INC.

CANTON, GEORGIA 30114



25 lbs.

DILUTION GUIDE

PARTS PER MILLION AVAILABLE CHLORINE

| | |
|------------------------|-------------|
| 1 oz. in 20 gal. water | 50 p.p.m. |
| 1 oz. in 10 gal. water | 100 p.p.m. |
| 1 oz. in 5 gal. water | 200 p.p.m. |
| 1 oz. in 1 gal. water | 1000 p.p.m. |

DIRECTIONS

PLANT-WALLS, FLOORS AND ETC.

Sanitize previously cleaned surfaces using a 1000 p.p.m. solution allowing not less than a 2 minute exposure time. (1 oz. PCC Sanitizer Powder to 1 gallon water).

EQUIPMENT

Sanitize previously cleaned equipment using a 100 p.p.m. solution allowing not less than a 2 minute exposure time. Equipment need not be rinsed before use. (1 oz. PCC Sanitizer Powder to 5 gal. water)



COMMERCIAL EGGS

from 10 a.m. to 60 p.p.m.

are sprayed with a

solution maintained at a

temperature solution at the

Allow eggs to dry

USE THAT METER

Final rinse :

with water for stock

Final rinse :

with water for stock

Refresh daily.

Use get sanitizer

← PCC CHLORINE SANITIZER POWDER

FOR USE IN SANITIZATION APPLICATIONS IN EGG PROCESSING PLANTS, POULTRY PROCESSING PLANTS, FINAL RINSE SANITIZING OF COMMERCIAL EGGS AND OTHER APPLICATIONS WHERE CHLORINE SANITIZATION IS NEEDED.

ACTIVE INGREDIENTS:

Potassium dichloro-s-triazinetriene 22

INERT INGREDIENTS 78

100

CAUTION: KEEP OUT OF REACH OF CHILDREN
If swallowed. Do not take internally. If accidentally swallowed, induce vomiting and call a physician. Prolonged contact may irritate eyes or nose.

USDA Registration No. 8344-6

MANUFACTURED BY



PROGRESS CHEMICAL COMPANY, INC.

CANTON, GEORGIA 30114

ACCEPTED

Feb. 6, 1968

REGISTERED UNDER THE FEDERAL INSECTICIDE ACT
FOR USE AS AN INSECTICIDE
DURING THE MONTH OF FEBRUARY 1968