

DIRECTIONS FOR USE

To be used as a dust without dilution in conventional ground or air dusters. Dust should be applied when the wind velocity is less than 5 miles per hour. The best hours for dusting are usually between 5:00 P.M. and 9:00 A.M. Apply on foliage when needed unless otherwise noted in special instructions and repeat at 7 to 10 day intervals if necessary.

CROP & INSECTS	RATE PER ACRE	INTERVAL (Last Application to Harvest)
COTTON —Thrips, Flea Beetles, Leafworm, Striped Blister Beetle, Boll Weevil, Bollworm, Fall Armyworm, Leaf Perforator, Leaf Beetles, Leafhoppers, Tarnished Plant Bug, Lygus & Other Plant Bugs, Grasshoppers	5-10 lbs. 10-20 lbs.	0 day
Pink Bollworm Salt-marsh Caterpillar, Stink Bug	15-25 lbs. 20 lbs.	
Do not pasture dairy animals or animals being finished for slaughter in cotton fields or feed gin waste.		
FIELD & SWEET CORN —Corn Earworm	10-20 lbs.	7 days [or forage]
Apply when first silks appear and repeat at 2 to 3 days intervals during silking period. Direct dusters to ear zone to insure complete coverage of each silk. On sweet corn for food it can be applied up to and including the day of harvest.		
Corn Earworm Adults, European Corn Borer, Fall Armyworms, Sap Beetles, Japanese Beetles, Flea Beetles, Leafhoppers	10-20 lbs.	
SUGAR BEETS —Leafhoppers, Armyworms, Webworms, Grasshoppers	10-20 lbs.	14 days
FRUITS, ALFALFA, CLOVERS, COWPEAS, SOYBEANS —Blister Beetles, Mexican Bean Beetle	5-10 lbs.	
No time limitations.		
Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvetbean Caterpillar, Green Cloverworm, Armyworm, Webworm, Corn Earworm, Stink Bugs, Grasshoppers	10 lbs. 10-15 lbs.	
TOBACCO (Field) —Harlequin, Bollworm, Flea Beetle, June Beetle, Japanese Beetle	10-20 lbs.	3 days before pruning or cutting
(Plant Beds)—Tobacco Flea Beetle, Green June Beetle Grubs 1 lb. per 100 sq. yds. 3 days before transplanting.		
RICE, FORAGE GRASS PASTURE —Thrips, Stink Bug, Armyworm, Grasshoppers	10-15 lbs.	0 day 14 day (Rice)
SORGHUM —Armyworms, Webworms, Corn Earworm, Stink Bug, Grasshoppers	10-20 lbs.	21 days (Grain) 0 day Forage
Sorghum Midge	15 lbs.	
Apply 3 or 4 days after heads have emerged from boot.		
ASPARAGUS (On Spears) —Asparagus Beetle	10-20 lbs.	1 day
Do not repeat application within 3 days.		
(On Ferns & Brush)—Asparagus Beetle, Apache Cicada	20-40 lbs.	
Post-harvest application		
BEANS —Lygus Bug, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Beetle, Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms	10-20 lbs.	0 day
Do not feed forage to dairy animals or animals being finished for slaughter.		
CUCUMBERS, SQUASH, CANTALOUPE, WATERMELONS —Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Cucumber Beetles, Fleabeetles	10 lbs.	0 day

Net Weight 50 Lbs.

P C C

10% SEVIN DUST INSECTICIDE

CONTAINS SEVIN CARBARYL INSECTICIDE

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE WARNING ON SIDE PANEL

ACTIVE INGREDIENTS

Carbaryl (1-Naphthyl N-methylcarbamate)..... 10% by Wt.

INERT INGREDIENTS..... 90% by Wt.

TOTAL..... 100% by Wt.

NOTICE TO BUYER

Seller warrants that the within product will comply with the description of the goods contained on the label but Buyer's remedy shall be limited exclusively to the remedy of rescission. The warranty is expressly in lieu of all other warranties and conditions express or implied, all liability for negligence and all other obligations or liabilities on Seller's part or on the part of anyone in Seller's chain of title. If unwilling to accept the product under these conditions, Buyer may return it for full credit.

B

Registered trademark of Union Carbide Corporation for carbaryl insecticide.

PROGRESS CHEMICAL COMPANY, INC.

MASTON, GEORGIA 30113

POTATOES, TOMATOES, SPOONPLANTS, PEPPERS —Fleabeetles, Cucumber Beetles, Tobacco Root Worm, Fall Armyworms, Corn Earworm, Tomato Leafworm, Tobacco Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Tomato Pinworm, Leafhoppers, Lygus Bug	10-20 lbs.	0 day
LETTUCE —Fleabeetles, Stink Bug, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Cabbage Caterpillars, Spittlebug, Lygus Bug	10-20 lbs.	3 days (Head) 14 day (Leaf)
CABBAGE —Flea Beetles, Leafhoppers, Harlequin Bug, Six-spotted Leafhopper (Aster Yellow Vector), Corn Earworm, Armyworms, Cabbage Caterpillars, Spittlebug, Lygus & Stink Bug	10-20 lbs.	
No time limitations.		
CABBAGE BROCCOLI, BRUSSEL SPROUTS, CABBAGEFLOWER, KALE —Fleabeetles, Harlequin Bug	5-10 lbs.	3 days
Corn Earworm, Armyworms, Cabbage Caterpillars	10-20 lbs.	
ORCHARD —Corn Earworm, Fleabeetles, Japanese Beetles, Stink Bug	10-20 lbs.	0 day
CRANBERRIES —Leafhoppers, Stink Bug, Bee Chaffer, Cutworms, Fleaworms, Cutworms, Japanese Beetle	15-40 lbs.	1 day
STRAWBERRIES —Fleabeetles, Lygus Bug, Spittlebug, Leafhopper, Strawberry Leaf Miner, Strawberry Weevil	10-20 lbs.	1 day
BLUEBERRY —Blueberry Maggot, Cherry Fruitworm, Grape Berry Fruitworm, European Fruit Lecanium, Japanese Beetle	15-20 lbs.	0 day
GRAPES —Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Miner, Japanese Beetle, Juno Beetle, Red-tailed Vine Weevil	20-30 lbs.	0 day
CITRUS (Non-bearing) —Soft Brown Scale	40-60 lbs.	
Apply to non-bearing citrus trees or to mature trees with no fruit present.		
ORNAMENTALS —Aster, Vase Anemone, Aster, Black, Rosewood, Camellia, Chrysanthemum, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose, Rose of Sharon, Yew, Zinnia, Apple, Begonia, Blister Beetle, Emerald Bug, Leaf Miner (Syrphid), Leaf Miner, Fleabeetles, Japanese Beetle, Leafhoppers, Leaf Miner, Orange Tortrix, Periodical Cicada, Plant Bug, Tort Caterpillar, Wasp (Exposed)		
Apply to completely cover foliage and repeat as necessary.		

CAUTION

Harmful if inhaled or swallowed. Do not take internally. Avoid prolonged breathing of dust. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothes. Change to clean clothing daily. Avoid storage near feed or food products.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS
NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

USDA REG. NO. 8344-7