

PCC
5% SEVIN®
DUST

PCC
5% SEVIN®
DUST

PCC
5% SEVIN®
DUST

PCC
5% SEVIN® DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS, WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

PCC 5% SEVIN DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. DO NOT use with any other chemical material. May be applied up to harvest except where noted.

DIRECTIONS FOR USE

BEANS—For control of Mexican bean beetles, leafhoppers, flea beetles, apply approximately 1/2 lb./1000 sq. ft. DO NOT feed treated bean vines to dairy or meat animals.

LETTUCE—For control of lygus and stink bugs, apply approximately 1 lb./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce. To avoid possible injury to foliage, do not apply when foliage is wet or when excessive humidity or rain is expected during the next two days.

CORN—Use approximately 3/4 lb./1000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese beetles, Sp. beetles, leafhoppers and European corn borer. For leaf beetles and grasshoppers in whorls (budworm damage) and for foliage feeders, make one or more applications beginning when damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing and treatments for home or small garden use. Apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For leaf beetles, grasshoppers and saws, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Reapply immediately after rain. Allow 7 days before harvest or grazing. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce the populations.

POTATOES, TOMATOES, EGGPLANTS and PEPPERS—For control of Colorado potato beetles, leafhoppers, and flea beetles apply approximately 1 lb./1000 sq. ft. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bug and lygus bugs, apply approximately 3/4 lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH—For control of pickleworm, melonworm, cucumber beetle, leaf beetles, flea beetles, apply approximately 1/2 lb./1000 sq. ft. To avoid possible injury to foliage, do not apply when foliage is wet or when excessive humidity or rain is expected during the next two days.

GRAPES—For control of leafhoppers, Japanese beetles, grape berry moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES—For control of meadow spittlebug, leafhopper, apply approximately 1 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS—For control of Japanese beetle, periodical cicada, aphid, tent caterpillars, lace bugs, boxwood leaf miner, apply approximately 1-1 1/4 lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dust is applied. In the presence of high humidity, DO NOT use on Boston Ivy.

DOGS, CATS—For control of fleas and brown dog ticks, apply liberally and rub into their feet, legs, chest, sleeping quarters and infested areas weekly. DO NOT treat with dust for more than 4 weeks old. DO NOT apply more than once weekly. For brown dog ticks, apply liberally on legs, feet and between toes. Comb out dead ticks a few hours after treatment. Rub dust into eyes. Force dust into cracks and crevices. Sprinkle lightly on furniture, floors and rugs. Rinse with a vacuum cleaner after an hour or two.

POULTRY—Chickens, Partridges, Pheasants and Pigeons. Northern fowl mite, chicken red mite, fleas and bed bugs in houses or quarters. Apply 1 lb./40 sq. ft. of floor and litter and to walls and interior surfaces. DO NOT apply directly to eggs or nest litter. Do not house birds in treated houses or areas within 7 days of slaughter.

For Northern Fowl mite and lice on the birds, as a supplement to treatment of premises, apply 1 lb./100 birds. Direct dust to vent and fluff areas and distribute throughout feathers. DO NOT repeat within 4 weeks. DO NOT APPLY within 7 days of slaughter.

DO NOT CONTAMINATE FEED, DRINKING WATER, FEEDING AND WATERING RECEPTACLES

SPECIAL NOTICE

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity several days following application. Does not control spider mites, but is compatible with most commonly used miticides.

*Trademark of Union Carbide Corporation.

INGREDIENTS

ACTIVE INGREDIENT	
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS	95%
TOTAL	100%

*Trademark Union Carbide Corp. for 1-naphthyl N-methylcarbamate

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on this Label

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Services. Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

EPA REG. NO. 8344-14 EPA EST. NO. 8344-GA-1

NET. WT. 4 POUNDS

MANUFACTURED BY

PROGRESS CHEMICAL CO., INC., CANTON, GEORGIA 30114

WARRANTY

The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, or toxicity to plants and to laboratory animals and of residue remaining on food crops.

The manufacturer makes no other express or implied guarantee, warranty of representation whatever of a similar or different sort, and none is to be inferred by the purchaser or user. No modification of this paragraph is authorized, except by specific reference to this paragraph in writing by an employee of the manufacturer.

Front

Side

Back

Photo Reduced is 6/10 of the original