

PCC 5% SEVIN DUST

INGREDIENTS

ACTIVE INGREDIENT	
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS	95%
TOTAL	100%

©Trademark Union Carbide Corp. for 1-naphthyl N-methylcarbamate

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on this Label

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidote. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Services. Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

EPA REG. NO. 8344-14

E.P.A. EST. NO. 8344-GA-1

NET WT. 50 POUNDS

MANUFACTURED BY
PROGRESS CHEMICAL CO., INC., CANTON, GEORGIA 30114

OLD LABEL

PCC
5% SEVIN
DUST

ACCEPTED
NOV 13 1984
9344-14

WARRANTY

The manufacturer guarantees and warrants that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated and that the directions, warnings and other statements on this label are reasonably and fairly based upon responsible testing of effectiveness, of toxicity to plants and to laboratory animals and of residue remaining on food crops.

The manufacturer makes no other express or implied guarantee, warranty of representation whatever of a similar or different sort, and none is to be inferred by the purchaser or user. No modification of this paragraph is authorized, except by specific reference to this paragraph in writing by an employee of the manufacturer.

PCC
5% SEVIN DUST

SEE FRONT LABEL FOR STATEMENT OF APPROVED USES, DIRECTIONS, WARNINGS AND PRECAUTIONS, AND AN EXPLANATION OF THE TESTING AND OTHER INFORMATION.

GENERAL INFORMATION

PCC 5% SEVIN DUST contains a wide range of insect pests. At first signs of insect infestation, particularly before damage has become serious, apply liberally to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. DO NOT use with lime or other alkaline material. May be applied as a dormant spray where noted.

DIRECTIONS FOR USE

APPLES—For control of European leaf beetles, leafhoppers, the leafminer, apply approximately 1 lb./1000 sq. ft. DO NOT feed treated trees close to day or night activities.

APRICOTS—For control of hoppers and other bugs, apply approximately 1 lb./1000 sq. ft. DO NOT apply to fruit within 7 days of harvest or within 14 days to leaf fall. To avoid possible injury to foliage, do not apply when foliage is wet or when excessive humidity or rain is expected during the next two days.

BERRIES—For approximately 1 lb./1000 sq. ft. for cane curculionids, gall midges, the leafminer, Japanese beetles, the leafhopper, leafhoppers and European saw flies. For cane borer and larvae in which European damage and for foliage beetles, make one or more applications to entire plant beginning when damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For hoppers or other pests not apply liberally into the whorls where plants are about three high. Repeat as needed, always 1 week before harvest. For insects attacking soil and stems, make 2 to 3 dry intervals starting when fruit falls, covering and maintaining until after leaf fall. These or more applications may be required depending on severity of infestation. Reapply immediately prior to June 7 days before harvest or grading of large applications of this product to the level required of care during the period stated and severely reduce bee populations.

BROCCOLI, TOMATOES, CUCUMBERS AND PEPPERS—For control of Colorado potato beetles, leafhoppers, and the leafminer, apply approximately 1 lb./1000 sq. ft. for control of tomato fruitworms, leaf miners, tomato hornworms, European corn borer, stalk bug and hoppers, apply approximately 1 lb./1000 sq. ft.

CUCUMBERS, CUCUMBER MELONS—For control of cucurbitworms, melonworms, cucumber beetles, squash bugs, the leafminer, apply approximately 1 lb./1000 sq. ft. To avoid possible injury to foliage, do not apply when foliage is wet or when excessive humidity or rain is expected during the next two days.

CURRIERS—For control of leafhoppers, Japanese beetles, grape berry moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES—For control of meadow spittlebug, leafhopper apply approximately 1 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

BEANS, ASPARAGUS, BRASSICAE VEGETABLES AND OTHER CUCURBITACEAE—For control of Japanese beetles, spotted alfalfa beetle, leaf caterpillars, leaf bugs, beanweb leaf miner, apply approximately 1-1 1/2 lb./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on beans in.

BROWN CATS—For control of flea and brown dog ticks, apply liberally and rub into skin, feet and legs. Feed sleeping quarters and elevated areas weekly. DO NOT treat kittens or puppies under 4 weeks old. DO NOT apply more than once weekly. For brown dog ticks be sure to also apply on legs, feet and between toes. Comb out dead ticks a few hours after treatment. Keep dust out of eyes. Feces and urine crumble and excreta. Sprinkle lightly on furniture, floors and rugs. Remove with a vacuum cleaner after an hour or two.

PHEASANTS—Chickens, Partridges, Pheasants and Pigeons. Northern field mice, chicken red mite, lice, fleas and bed bugs in houses or quarters. Apply 1 lb./40 sq. ft. of floor and litter, and to roosts and interior surfaces. DO NOT apply directly to eggs or nest litter. Do not house birds in treated houses or areas within 7 days of slaughter.

For Northern Field mice and lice on the birds, as a supplement to treatment of premises, apply 1 lb./100 birds. Sweep dust to spot and buff areas, and distribute throughout facilities. DO NOT repeat within 4 weeks. DO NOT APPLY within 7 days of slaughter.

DO NOT CONTAMINATE FEED, DRINKING WATER, FIBERS AND WASHING RECEPTACLES.

SPECIAL NOTICE

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity if several days duration following application. Do not control spider mites, but is compatible with most commonly used miticides.

*Trademark of Union Carbide Corporation

BEST AVAILABLE COPY