

Agrisect

BRAND

PARATHION 25% WETTABLE POWDER

ACTIVE INGREDIENTS:

PARATHION*

INERT INGREDIENTS:

	25%
	75%
TOTAL	100%

*O,O-Diethyl O-p-nitrophenyl thiophosphate

DANGER



POISON



DANGER

KEEP OUT OF REACH OF CHILDREN

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN! RAPIDLY ABSORBED THROUGH SKIN! DO NOT GET IN EYES OR ON SKIN. IN CASE OF CONTACT WASH IMMEDIATELY WITH SOAP & WATER. WASH HANDS, ARMS AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP & HOT WATER BEFORE RE-USE. WEAR NATURAL RUBBER GLOVES, PROTECTIVE CLOTHING AND GOGGLES. WEAR MASK OR RESPIRATOR OF TYPE PASSED BY U.S. DEPT OF AGRICULTURE FOR PARATHION PROTECTION. KEEP ALL UNPROTECTED PERSONS OUT OF OPERATING AREAS OR VICINITY OF DRIFT. VACATED AREAS SHOULD NOT BE RE-ENTERED UNTIL DRIFTING INSECTICIDE AND VOLATILE RESIDUES HAVE DISSIPATED.

THIS PRODUCT WILL KILL FISH & WILDLIFE. KEEP OUT OF LAKES, STREAMS & PONDS. BIRDS FEEDING ON TREATED AREAS MAY BE KILLED.

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HERewith.

net contents

pounds

FIRST AID: CALL PHYSICIAN IMMEDIATELY. UNTIL HE ARRIVES FOLLOW DIRECTIONS GIVEN BELOW:

INTERNAL: If swallowed make patient vomit by giving warm salt or soapy water. Give milk or water freely. Have patient lie down and keep warm. Do not give anything by mouth to unconscious patient. NOTE TO PHYSICIAN: Therapeutic doses of Atropine or 2-PAM appear to be effective as an antidote. EXTERNAL: Remove all contaminated clothing and flush with water for at least 15 minutes. EYES: Immediately flush with water for 15 minutes.

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL

Do not apply when weather conditions favor drift of spray from areas treated. Do not apply while trees are in bloom. Use agitation in spray tank.

APPLES, PEARS: To control mites, use $\frac{1}{2}$ to $\frac{1}{4}$ lb/100 gal. water not later than 14 days before harvest. For mealy bugs use 1#/100 gal. water not later than 14 days before harvest. For woolly aphids on apples use 1#/100 gal water early in growing season.

PEACHES: For control of cottony cushion scale use $\frac{1}{2}$ #/100 gal water for crawler only. Make two applications. Areas other than California: full coverage spray. Do not apply within 14 days of harvest. Do not apply more than 5# actual Parathion per acre per year. California: Do not apply within 21 days of harvest. Do not apply more than once after bloom. Do not apply more than 5# actual Parathion per acre between Jan. 1 and harvest.

TOMATOES, CARROTS: For aphids use $\frac{1}{2}$ #/100 gal water per acre not later than 10 days before harvest on tomatoes, 15 days on carrots. DO NOT USE TREATED CARROT TOPS FOR FOOD OR FEED.

RINSE EQUIPMENT AND CONTAINERS AND DISPOSE OF WASTES BY BURYING IN NON-CROP LANDS AWAY FROM WATER SUPPLIES. CONTAINERS SHOULD BE DISPOSED OF BY PUNCHING HOLES IN THEM AND BURYING WITH WASTES OR BY BURNING (KEEP OUT OF SMOKE).

USDA REG. # 8343- 44

GABRIEL
CHEMICALS LTD.

ROBBINSVILLE, NEW JERSEY

ACCEPTED WITH COMMENTS