

8378-12

04/30/87

1/2

NET WEIGHT 40 LBS. (18.2 Kg.)

THIS PRODUCT MUST NOT BE USED
ON GOLF COURSES AND SOD FARMS



ACCEPTED
MAR 9 1983
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 8378-12

ACCEPTED
APR 30 1987
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 8378-12

LAWN INSECT CONTROL PLUS FERTILIZER

28 - 3 - 9

Ready-to-use Contains 3.33% Diazinon Insecticide.

ACTIVE INGREDIENT: Diazinon*	
{O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)phosphorothioate} . . .	3.33%
INERT INGREDIENTS	96.67%
Total	100.00%

*Trademark of Ciba-Geigy

GUARANTEED PLANT FOOD ANALYSIS

Total Nitrogen (N)	28%
1.2% Ammoniacal Nitrogen	
20.8% Urea Nitrogen	
6.0% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	9%
Sulfur (S)	4.5%

Primary Plant Foods derived from: Urea, Muriate of Potash, Ammoniated Phosphates, and Sulfur Coated Urea. Sulfur derived from: Sulfur Coated Urea.

Potential Acidity 1,040 pounds Calcium Carbonate equivalent per ton.

KEEP OUT OF REACH OF CHILDREN!
CAUTION
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

EPA Reg. No. 8378-12

EPA Est. No. 8378-IN-1

12/22/86
IRB/RD

Shaw's
LAWN INSECT CONTROL PLUS FERTILIZER

Shaw's
LAWN INSECT CONTROL PLUS FERTILIZER



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling and before smoking or eating. Keep children and pets off treated areas until granules have been watered off grass blades and grass has been allowed to dry.

STATEMENT OF PRACTICAL TREATMENT

- If swallowed- Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
 - If inhaled- Remove victim to fresh air. Apply artificial respiration if indicated.
 - If on skin- Remove contaminated clothing and wash affected areas with soap and water.
 - If in eyes- Flush eyes with plenty of water. Get a physician if irritation persists.
- Note to Physician - if symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SHAW'S TURF CARE GRANULAR INSECT CONTROL KILLS:

Ants	Crickets	Springtails
Armyworms	Cutworms	Sod webworms
Bermuda mites	Digger wasps	Sowbugs
Billbugs	Earwigs	White grubs of:
Chiggers	Lawn moths	-Japanese beetle
Chinch bugs	Leafhoppers	-European chafer
Clover mites	Millipedes	-Southern chafer

OVERALL LAWN APPLICATION

SHAW'S Turf Care Granular Insect Control plus Lawn Food controls the following lawn insects on both grass and dichondra lawns: Chinch bugs - Earwigs - Ants - Lawn moths (Sod webworms) - Bermuda mites - Crickets - Armyworms - Chiggers - Clover mites - Cutworms - Leafhoppers - Sowbugs - Millipedes - Springtails (Collembola).

Apply SHAW'S Turf Care Granular Insect Control plus Lawn Food to your lawn with a fertilizer spreader at the rates shown. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

Apply at the rate of 3.0 lbs. per 1000 sq. ft. (an area 20' by 50').

BILL BUGS, WHITE GRUBS OF JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER

Apply 3.6 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

HYPERODES WEEVIL (NEW YORK ONLY)

Apply 3.0 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap container and put in trash collection.

SPREADER SETTINGS

	3 lbs. per 1000 sq. ft.	3.6 lbs. per 1000 sq. ft.
Scotts	5	5 1/2
Shaw's	3	5 1/2
Cyclone	3	5
Sears Broadcast	2 1/2	4

NOTE: Above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply 4.8 oz. per 100 sq. ft. for the 3 lbs. per 1000 sq. ft. rate or 5.7 oz. per 100 sq. ft. for the 3.6 lbs. per 1000 sq. ft. rate. Rate of walk, moisture and spreader condition can vary the setting necessary to apply the correct amount.

40 LBS. NET WT. (18.2 Kg.)

KNOX FERTILIZER & CHEMICAL COMPANY, INC.

KNOX INDIANA 46534