


Reg # 8518-1-1 PM-13 1991



# LAWN INSECT CONTROL Plus Fertilizer 28-3-9

## FOR USE ON HOME LAWNS

**This product must not be used on golf courses or sod farms**  
Ready To Use - Contains 3.33% Diazinon Insecticide

**ACTIVE INGREDIENT:**  
Diazinon [O,O-diethyl-O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate] - - - - 3.33%  
**INERT INGREDIENTS:** - - - - - 96.67%  
TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
STATEMENTS OF PRACTICAL TREATMENT  
**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists. **NOTE TO PHYSICIAN:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
EPA REG. NO. 8378-12 EPA EST. NO. 8378-IN-1  
NET WEIGHT 15 POUNDS (6.8 Kg)

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. May cause skin sensitization following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Where "watering in" is required, do not allow children or pets on those treated areas until granules are watered into the soil and grass is dry. Do not use on pets, animals or livestock.

**ENVIRONMENTAL HAZARDS**  
This product is highly toxic to birds and fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, NY between Nov. 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e. ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Runoff may be hazardous to aquatic organisms, in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

**DIRECTIONS FOR USE** It is a violation of Federal law to use this product in a manner inconsistent with its label.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage or disposal.  
**STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. **DISPOSAL:** Do not reuse empty bag. Wrap in several layers of newspaper and discard in trash.

**IMPORTANT** Read the entire Directions for Use and Conditions of Sale before using this product. Follow directions carefully.

### For Grass and Dichondra Lawns Feeds Lawn/Controls Surface and Soil Insects

## Surface Feeding Insects

**OVERALL LAWN APPLICATION**  
**CONTROLS:** Ants, Armyworms, Bermuda Mites, Billbugs, Brown Dog Ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Digger Wasps, Earwig, Fleas, Lawn Moths, Leafhoppers, Millipedes, Sod Webworms, Sowbugs, Springtails.

Apply at a rate of 3.0 pounds per 1,000 square feet. Supplies your lawn with a balanced plant food and controls surface feeding insects as listed above. Apply with a lawn fertilizer spreader at the rates shown when insect damage appears. Application should be made when grass is dry. Repeat as necessary.

## Soil Insects

**CONTROLS:** Billbugs, White Grubs of Japanese Beetle, European Chafer, Southern Chafer & Hyperodes Weevil\*.  
Apply 3.0 pounds per 1,000 square feet of lawn. Supplies your lawn with a complete plant food and controls soil insects as listed above. Apply with a lawn fertilizer spreader at the rates shown when insect damage appears. Water lawn thoroughly after application. For grub control, make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.\* (New York Only), treat problem areas in mid-April & in mid-May.

Spreader	Setting for 3.0 lbs. per 1,000 Sq.Ft. of Turf
Scott	5
Shaw's	3
Cyclone	3
Sears Broadcast	2 1/2

**IMPORTANT:** Spreader setting is approximate. Age, condition of spreader, and operator speed may cause a wide variation. Be sure to calibrate your spreader with each application.

Total Nitrogen (N)	28.0%
1.20% Ammoniacal Nitrogen	•••
20.80% Urea Nitrogen	•••••
6.00% Coated Slow Release Urea (N)	•••••
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	3.0%
Soluble Potash (K <sub>2</sub> O)	9.0%
Sulfur (S) (Free)	4.5%
Primary Plant foods derived from Sulfur Coated Urea, Urea, Ammonium Phosphate, and Muriate of Potash.	
Potential Acidity 1,040 lbs. Calcium Carbonate per ton.	

**NOTICE:** Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(EPA corrected Version dated 09-17-90 of 03-01-90b accepted copy)

