

PRE-EMERGE CRABGRASS CONTROL and FERTILIZER 25-5-8

SHAW'S PRE-EMERGE CRABGRASS CONTROL AND FERTILIZER 25-5-8

50 POUNDS COVERS 10,000 SQ.FT. KILLS CRABGRASS, FOXTAIL AND OTHER WEEDS LISTED AS THEY GERMINATE

FEEDS THE LAWN - CONTAINS DACTHAL

ACTIVE INGREDIENT DCPA (dimethyl tetrachloroterephthalate) INERT INGREDIENTS: --TOTAL 100.00%

Dacthal is a Registered Trademark of Fermenta Co.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENTS IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an uncorrelieus or convulsing person, IF IN EYES. Flush eyes with plenty of water. Call a physician if irritation persists IF ON SKIN. Wash aftertail area with scep and water.

SEE SIDE/BACK PANEL FOR ADDITIONAL PHOUSE TOWARY STATEMENTS EPA REG. NO. 8378-13 EFA EST. NO. 8378-IN-1

NET WEIGHT 50 POUNDS (22.68 Kg)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if ewallowed. Causes moderate eye irritation, Ayoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Do not contaminate feed or foodstuffs. Do not feed treated foliage to livestock or graze treated areas.

ENVIRONMENTAL HAZARDS

Collect and incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes, or potholes). Do not contaminate water when disposing of equipment wash-

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling,

USE ONLYWHEN WEARING THE FOLLOWING PROTECTIVE CLOTH-ING. Wear long-sleeved shirt and long-legged pants and chemical resistant gloves. IMPORTANT! If the pesticide comes in contact with skin wash off with soap and water. Always wash hands, face, and arms with scap and water before smoking, exting, drinking, or toileting. AFTERUSE: Before removing gloves, wash themn with soap and water. Take off all work clothes and shoes. Shower using soap and water, Do not reuse contaminated clothing. Clothing worn during use must be faundered separately from household articles.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal, STORAGE: Store product in original container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved wastes disposal facility. CONTAINER DISPOSAL: Completely empty beg into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities by burning. If burned stay out of smoke. Household-Do not reuse empty bay. Wrap and put in trash.

WHEN TO USE .

Early Spring application of Shaw's Pre-Emerge Crabgrass Control and Fertilizer before weeds germinate, provides full season control of crabgrass and adds soil nutrients essential to good grass growth. Crabgrass seed germination usually coincides with the time Forsythia blossom start to fall.

Also controls Carpetweed, Chickweed, Foxtail, Johnson Grass (from seed), Lambsquarter, Purslane, and Witchgrass

HOW MUCH TO USE

After making certain spreader is in good operating condition, apply this material at the rate of 5 pounds per 1,000 sq.ft. (see spreader settings) Treated areas should be watered lightly (1/3 to 1/2 inch of water if no rain falls within five days of application.

Reseeding should be delayed until 60 days after application.

NOTE FOR NEW LAWNS - Do not apply to newly seeded lawn until after lewn is completely green...when newly seeded grass is from 1 to 2 inches in height. Crabgrass already growing will not be controlled.

NOTE: Do not use on Putting Greens. Do not use on Bentgrasses when mowed at putting green height. Do not graze or feed grass clippings to livestock. Do not use on Dichondre lewris.

SPREADER SETTINGS

Sears Calibrated 3 3/4; Sears Broadcast 2 3/4; Cyclone 2 3/4; Jackson 3; Scotts Model #35 2 1/2; Central (Model 1662) 31/2.

MINIMUM GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N) --- --

1.96% Ammoniacal Nitrogen

17.04% Ures Nitrogen

6 00% Sulfur Coated Urea

0.00% Water Insoluble Nitrogen

AVAILABLE PHOSPHORIC ACID (P2OS) - 5.00% SOLUBLE POTASH(K6O) - - 8.00%

Primary Plant Foods derived from Ures, Diammonium Plicatrigite, and Muriate of Potesh. Potential scidity equivals x to 30% pounds of Culcium Carbonale per ton.

Notice: Conditions of use are beyond selects control. Selects liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use buyend claims on اعطعا

NET WEIGHT 50 POUNDS

KNOX FERTILIZER & CHEMICAL CO <u>KNOX, INDIANA 46534</u>

sammented for the posticade tegisters t under \$378-13 EPA Noy. No.