



United States Environmental Protection Agency Washington, DC 20460

Registration [ ] Amendment [ ] Other [x]

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number: 8378-19
2. EPA Product Manager: Joanne I. Miller
3. Proposed Classification: [x] None [ ] Restricted
4. Company/Product (Name): Shaw's Premium Lawn Food With Crabgrass Control
5. Name and Address of Applicant: Knox Fertilizer Company, P. O. Box, Knox, IN 46534
6. Expedited Review: EPA Reg. No. NA, Product Name NA

Section - II

Amendment - Explain below [ ]
Resubmission in response to Agency letter dated [ ]
Notification - Explain below [x]
Final printed labels in response to Agency letter dated [ ]
'Me Too' Application [ ]
Other - Explain below [ ]
NOTIFICATION FEB 28 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Notification of label change relative to PR Notice 2001-6. This notification is consistent with the guidance in PR Notice 2001-6 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product.

Section - III

1. Material This Product Will Be Packaged In:
Child-Resistant Packaging [ ] Yes [ ] No
Unit Packaging [ ] Yes [ ] No
Water Soluble Packaging [ ] Yes [ ] No
2. Type of Container: [ ] Metal [ ] Plastic [ ] Glass [ ] Paper [ ] Other (Specify)
3. Location of Net Contents Information: [ ] Label [ ] Container
4. Size(s) Retail Container
5. Location of Label Directions
6. Manner in Which Label is Affixed to Product: [ ] Lithograph Paper glued Stenciled [ ] Other

Section - IV

1. Contact Point: Name Robert F. Bischoff, Title Agent, TotalTurf Consulting LLC, Telephone No. (317) 873-0222
Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.
2. Signature: Robert F. Bischoff
3. Title: Agent, TotalTurf Consulting LLC
4. Typed Name: Robert F. Bischoff
5. Date: February 20, 2003
6. Date Application Received (Stamped)

NOTIFICATION  
FEB 28 2003

# Shaw's® Premium Lawn Food With Crabgrass Control

For Control of Crabgrass and Most Other Annual Weed Grasses on Home Lawns, Golf Courses, Parks and Other Ornamental & Recreational Turf. For Use On Established: Bluegrass (Perennial) - Ryegrass (Perennial) - Fescue - Bahiagrass - Bermudagrass - Centipedegrass - St. Augustinegrass - Soysigrass

## Preemergence Crabgrass Control Plus Fertilizer

### ACTIVE INGREDIENTS:

Benefin; N-butyl-N-ethyl-alpha,alpha,alpha,trifluoro-2,6-dinitro-p-toluidine .....	1.00%
Trifluralin; Alpha,alpha,alpha,trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine .....	0.50%
OTHER INGREDIENTS: .....	<u>98.50%</u>
<b>TOTAL</b>	<b>100.00%</b>

EPA Reg. No. 8378-19  
NET WEIGHT 15 LBS.

EPA EST. No. 8378-IN-1

### GUARANTEED FERTILIZER ANALYSIS

- Nitrogen-----XX%
- Ammonical (N)-----XX%
- Urea (N)-----XX%
- Sulfur Coated Urea (N)--XX%
- Phosphoric Acid (P2O5)-----XX%
- Soluble Potash (K2O)-----XX%

Primary Plant Food derived from Urea, Sulfur Coated Urea, Muriate of Potash, Ammoniated Phosphate.  
Potential Acidity XX lbs. Calcium Carbonate per ton.

## KEEP OUT OF REACH OF CHILDREN

### WARNING

### FIRST AID

**IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center, or doctor, or when going for treatment.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury. Harmful if absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged, or frequently repeated skin contact may cause allergic reactions in some individuals.

#### Personal Protective Equipment

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety recommendations

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine and estuarine fish and aquatic invertebrates, including shrimp, and oysters. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes, or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons either directly, or through drift. Only protected handlers may be in the area during application. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

#### Entry Restrictions

Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

### WHERE TO APPLY FOR CRABGRASS CONTROL

When applied according to these directions this product can be used safely on the following turfgrass varieties: Perennial Bluegrass varieties, Perennial Rye grass varieties, Bermudagrass, Bahiagrass, Centipedegrass, Fescue varieties, St. Augustinegrass and Zoysiagrass growing in home lawns, golf courses, parks and all other commercial, recreational and ornamental turfgrass areas. Will not harm most common ornamental trees and shrubs when used as directed.

### OTHER WEEDS CONTROLLED

This product contains a combination of Benefin and Trifluralin pre-emergence Herbicides plus a Lawn Food. It should be applied to established turf for the control of Crabgrass (Smooth and Hairy), Goosegrass (Silver Crab or Crowfoot), Barnyardgrass (Watergrass), Foxtail (Green and Yellow), and Annual Bluegrass (Poa Annuua) while providing a complete lawn fertilization.

### WHEN TO APPLY

Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

### WHEN AND HOW TO TIME THE APPLICATION

Successful pre-emergence control of the annual grassy weeds as listed requires proper application timing. It must be applied 1 to 2 weeks prior to the germination of the weed seeds. All summer annual weeds do not germinate at the same time. For this reason application timing must coincide with the normal germination period of any one of the annual grassy weeds. If not, poor and erratic weed control will result due to improper application.

### WATER THE LAWN AFTER APPLICATION

It is not necessary but optimum weed control performance will be obtained if treated areas are irrigated with  $\frac{1}{4}$  to  $\frac{1}{2}$  inch water soon after application. Any cultural practices that disturb the soil such as verticutting, should be done prior to application of this product.

### IMPORTANT PRECAUTIONS IN USING THIS PRODUCT

1. DO NOT apply in the spring to Bermuda grass which has been overseeded with winter grasses (annual rye) since thinning will occur.
2. DO NOT use on Dichondra, nor to newly sprigged areas of Bermuda grass, St. Augustine grass or Centipede grass, nor to Bentgrass or putting greens until these turfs are well established.
3. DO NOT apply in spring to turfgrass planted the previous fall. This product may thin established Annual Bluegrass (Poa Annuua) and Fine-Leafed Fescue lawns when used at higher rates than 133 lbs. per acre (2.0 lb. A.I. per Acre). Do not overdose.

### "SPRING GERMINATING"

#### ANNUAL GRASSES: CRABGRASS, FOXTAIL, BARNYARDGRASS

- A. Northern Cool Season Grasses - For preemergence control, apply 100 to 133 lbs. per acre (equal to 1.5 to 2 lbs. active ingredient/acre) applied in the spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 15 lb. bag will cover 5,000 to 6,500 sq. ft. Use for 5,000 sq. ft. when turf is completely overrun with weeds. Use for 6,500 sq. ft. when weeds are sparse and the turf is in a yearly lawn maintenance program.

- B. Southern Warm Season Turf grass - For preemergence control apply 133 to 200 lbs. per acre (equal to 2 to 3 lbs. active ingredient per acre), in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 15 lb. bag will cover 3,300 to 5,000 sq. ft. Use for 3,300 sq. ft. where turf is completely overrun by weeds. Use for 5,000 sq. ft. when weeds are sparse or the turf is in a yearly crabgrass control program.

### **"LATE SPRING-EARLY SUMMER GERMINATING" ANNUAL GOOSEGRASS (SILVER CRAB)**

Although classified as a summer annual grass, Goosegrass normally germinates 4 to 6 weeks later than the crabgrass, foxtails and other summer annual grass species. Applications of this product should therefore, be timed accordingly.

- A. Northern Cool Season Turf grasses - For preemergence control apply 133 lbs./acre (equal to 2 lbs. active ingredient per acre) of this product in late spring or early summer one to two weeks prior to the expected germination period of Goosegrass. Use 15 lbs. to cover 5,000 sq. ft.
- B. Southern Warm Season Turf grasses - For preemergence control, apply 133 to 200 lbs. per acre (equal to 2 to 3 lbs. active ingredient per acre) of this product in late spring or early summer one to two weeks, prior to the expected germination of Goosegrass. Use one 15 lb. bag to cover for 3,300 to 5,000 sq. ft. Use for 3,300 sq. ft. where turf is completely overrun by weeds. Use for 5,000 sq. ft. where weeds are sparse and the area is in a yearly lawn maintenance program.

### **APPLY TWICE FOR FULL SEASON CONTROL IN NORTHERN AND SOUTHERN TURFGRASSES**

In areas of the United States where preemergence control of both early germinating annual grasses (Crabgrass, Foxtail, and Barnyardgrass) and the later germinating Goosegrass is desired, two applications of this product at the recommended rate, spaced 8 to 10 weeks, may be made to provide full season control of these annual grass species.

### **POA ANNUA (ANNUAL BLUEGRASS) CONTROL**

In areas of heavy infestation of Annual Bluegrass, its elimination will leave areas of thin turf. Employ proper cultural practices to establish a new turf.

- A. Northern Cool Season Turf Grasses - For preemergence control apply 133 lbs. per acre (equal to 2 lbs. active ingredient per acre) of this product per acre in late summer or early fall and again in late winter or spring one to two weeks prior to expected germination of annual bluegrass. One 15 lb. bag will treat 5,000 sq. ft. of turf grass.
- B. Southern Warm Season Turf grasses - For preemergence control apply 200 lbs. per acre (equal to 3 lbs. active ingredient per acre) of this product per acre in the later summer or early fall one to two weeks prior to expected germination of poa annua. One 15 lb. bag will treat and feed 3,300 sq. ft. of turfgrass. In southern areas where germination continues for extended periods of time a second application at 200 lbs. per acre can be made 10 to 12 weeks after the initial treatment.

### **RESEEDING TREATED AREAS**

1. Delay reseeding - 6 weeks when applying at the rate of 100 or 133 lbs./acre equal to 1.5 to 2.0 lbs. active ingredient/acre, or when using one 15 lb. bag to cover 5,000 to 6,500 sq. ft.

2. Delay reseeding - 12 to 16 weeks when applying at rate of 200 lbs./acre equal to 3 lbs. of a.i. per acre or when using one 15 lb. bag to cover 3,300 sq. ft. of turfgrass. When reseeding increase the amount of seed put down by 15%.

**SPREADER SETTINGS**

The following spreader settings are for use with new equipment. Your spreader may need recalibrating since proper application is based on equipment condition with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven or improper application.

Spreader	15# bag per sq. ft.	Northern Lawns	Southern Lawns
		Crabgrass etc. Poa Annua	Crabgrass etc. Poa Annua
Scott #35	3,300	XX	XX
	5,000	XX	XX
	6,500	XX	XX
Cyclone B1	3,300	XX	XX
	5,000	XX	XX
	6,500	XX	XX

15 lbs. to treat 3,300 sq. ft. = 200 lbs./acre = 3.00 lbs. A.I./acre

15 lbs. to treat 5,000 sq. ft. = 133 lbs./acre = 2.00 lbs. A.I./acre

15 lbs. to treat 6,500 sq. ft. = 100 lbs./acre = 1.50 lbs. A.I./acre

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store in original, closed container only. PRODUCT DISPOSAL: IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

**KNOX FERTILIZER COMPANY, INC.**  
**P.O. BOX 248**  
**KNOX, IN 46534**  
**NET WEIGHT 15 LBS. (6.81 kg.)**

# Shaw's® Premium Lawn Food With Crabgrass Control

For Control of Crabgrass and Most Other Annual Weed Grasses on Home Lawns, Golf Courses, Parks and Other Ornamental & Recreational Turf. For Use On Established: Bluegrass (Perennial) - Ryegrass (Perennial) - Fescue - Bahiagrass - Bermudagrass - Centipedegrass - St. Augustinegrass - Soysiagrass

## Preemergence Crabgrass Control Plus Fertilizer

### ACTIVE INGREDIENTS:

Benefin; N-butyl-N-ethyl-alpha,alpha,alpha-trifluoro-2,6-dinitro-p-toluidine .....	1.00%
Trifluralin; Alpha,alpha,alpha-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine .....	0.50%
OTHER INGREDIENTS: .....	<u>98.50%</u>
<b>TOTAL</b>	<b>100.00%</b>

EPA Reg. No. 8378-19  
NET WEIGHT 15 LBS.

EPA EST. No. 8378-IN-1

### GUARANTEED FERTILIZER ANALYSIS

Nitrogen-----XX%  
 Ammonical (N)-----XX%  
 Urea (N)-----XX%  
 Sulfur Coated Urea (N)--XX%  
 Phosphoric Acid (P2O5)-----XX%  
 Soluble Potash (K2O)-----XX%

Primary Plant Food derived from Urea, Sulfur Coated Urea, Muriate of Potash, Ammoniated Phosphate.  
Potential Acidity XX lbs. Calcium Carbonate per ton.


## KEEP OUT OF REACH OF CHILDREN

## WARNING

### FIRST AID

**IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.



**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center, or doctor, or when going for treatment.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury. Harmful if absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged, or frequently repeated skin contact may cause allergic reactions in some individuals.

#### Personal Protective Equipment

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety recommendations

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine and estuarine fish and aquatic invertebrates, including shrimp, and oysters. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes, or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons either directly, or through drift. Only protected handlers may be in the area during application. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

#### Entry Restrictions

Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.



9/13

### WHERE TO APPLY FOR CRABGRASS CONTROL

When applied according to these directions this product can be used safely on the following turfgrass varieties: Perennial Bluegrass varieties, Perennial Rye grass varieties, Bermudagrass, Bahiagrass, Centipedegrass, Fescue varieties, St. Augustinegrass and Zoysiagrass growing in home lawns, golf courses, parks and all other commercial, recreational and ornamental turfgrass areas. Will not harm most common ornamental trees and shrubs when used as directed.

### OTHER WEEDS CONTROLLED

This product contains a combination of Benefin and Trifluralin pre-emergence Herbicides plus a Lawn Food. It should be applied to established turf for the control of Crabgrass (Smooth and Hairy), Goosegrass (Silver Crab or Crowfoot), Barnyardgrass (Watergrass), Foxtail (Green and Yellow), and Annual Bluegrass (Poa Annua) while providing a complete lawn fertilization.

### WHEN TO APPLY

Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

### WHEN AND HOW TO TIME THE APPLICATION

Successful pre-emergence control of the annual grassy weeds as listed requires proper application timing. It must be applied 1 to 2 weeks prior to the germination of the weed seeds. All summer annual weeds do not germinate at the same time. For this reason application timing must coincide with the normal germination period of any one of the annual grassy weeds. If not, poor and erratic weed control will result due to improper application.

### WATER THE LAWN AFTER APPLICATION

It is not necessary but optimum weed control performance will be obtained if treated areas are irrigated with  $\frac{1}{4}$  to  $\frac{1}{2}$  inch water soon after application. Any cultural practices that disturb the soil such as verticutting, should be done prior to application of this product.

### IMPORTANT PRECAUTIONS IN USING THIS PRODUCT

1. DO NOT apply in the spring to Bermuda grass which has been overseeded with winter grasses (annual rye) since thinning will occur.
2. DO NOT use on Dichondra, nor to newly sprigged areas of Bermuda grass, St. Augustine grass or Centipede grass, nor to Bentgrass or putting greens until these turfs are well established.
3. DO NOT apply in spring to turfgrass planted the previous fall. This product may thin established Annual Bluegrass (Poa Annua) and Fine-Leafed Fescue lawns when used at higher rates than 133 lbs. per acre (2.0 lb. A.I. per Acre). Do not overdose.

### "SPRING GERMINATING"

#### ANNUAL GRASSES: CRABGRASS, FOXTAIL, BARNYARDGRASS

- A. Northern Cool Season Grasses - For preemergence control, apply 100 to 133 lbs. per acre, (equal to 1.5 to 2 lbs. active ingredient/acre) applied in the spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 15 lb. bag will cover 5,000 to 6,500 sq. ft. Use for 5,000 sq. ft. when turf is completely overrun with weeds. Use for 6,500 sq. ft. when weeds are sparse and the turf is in a yearly lawn maintenance program.





11/13

2. Delay reseeding - 12 to 16 weeks when applying at rate of 200 lbs./acre equal to 3 lbs. of a.i. per acre or when using one 15 lb. bag to cover 3,300 sq. ft. of turfgrass. When reseeding increase the amount of seed put down by 15%.

### SPREADER SETTINGS

The following spreader settings are for use with new equipment. Your spreader may need recalibrating since proper application is based on equipment condition with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven or improper application.

Spreader	15# bag per sq. ft.	Northern Lawns		Southern Lawns	
		Crabgrass etc.	Poa Annua	Crabgrass etc.	Poa Annua
Scott #35	3,300		XX		XX
	5,000		XX		XX
	6,500		XX		XX
Cyclone B1	3,300		XX		XX
	5,000		XX		XX
	6,500		XX		XX

15 lbs. to treat 3,300 sq. ft. = 200 lbs./acre = 3.00 lbs. A.I./acre

15 lbs. to treat 5,000 sq. ft. = 133 lbs./acre = 2.00 lbs. A.I./acre

15 lbs. to treat 6,500 sq. ft. = 100 lbs./acre = 1.50 lbs. A.I./acre

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store in original, closed container only. In case of leak or spill, contain material and use as directed or dispose of as waste. **PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty bag. Wrap in several layers of newspaper and put in trash.

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

**KNOX FERTILIZER COMPANY, INC.**  
**P.O. BOX 248**  
**KNOX, IN 46534**  
**NET WEIGHT 15 LBS. (6.81 kg.)**

**Inserted: PRODUCT DISPOSAL: IF EMPTY:** Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.



