



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

APR 15 2010

Mr. Robert F. Bischoff Agent, Total Turf Consulting LLC Knox Fertilizer Company, Inc. P.O. Box 248 Knox, IN 46534

RE: Notification of Removal of Toll Free Number from the Disposal Instructions Section

EPA Registration Number: 8378-20 Date of Submission: March 23, 2010

Dear Mr. Bischoff:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-1 dated March 23, 2010, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 2007-1 and finds that the actions requested fall within the scope of PRN 2007-1. The label submitted with the application has been stamped "Notification" will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader

Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before compared to	m.	Form Appro	ve)MB No. 2	070-0060	્રિં? D. Approval expires 2-28-95
SEPA Environmental Pro- Washington,	tection Agen	icy	Registra Amendn Other	tion	OPP Identifier Number
Appl	ication for P	esticide - Section	on I		
1. Company/Product Number 8378- 20		2. EPA Product Manag Kathryn V. Montagi	er	3. Pro	posed Classification
Company/Product (Name) Shaw's 18-5-9 Turf Food with XL Crabgrass Co		PM# 23			
5. Name and Address of Applicant (Include ZIP Code) Knox Fertilizer Company, Inc. P. O. Box 248 Knox, IN 46534 Check if this is a new address		-	similar or identi		FIFRA Section 3(c)(3) mposition and labeling
	Secti	on - II	-		
Amendment - Explain below. Resubmission in response to Agency letter dated. Notification - Explain below.		Final printed la Agency letter "Me Too" App Other - Explain	dication.	NOTIF	1 5 2010
Explanation: Use additional page(s) if necessary. (For Notification of label change relative to PR Notice 2007-1. To regulations at 40 CFR 156.10 and 40 CFR 152.46, and no comproduct. I understand that it is a violation of 18 U.S.C. Sections is the remaining the subject to enforcement action and penalties under the subject to enforcement action and penalties.	his notification is co other changes have 1001 to willfully mal equirements of 40 C	nsistent with the guidand been made to the labeling se any false statement to FR 156.10 and 40 CFR	ng or the confident EPA. I further un	ial statem derstand t	ent of formula of this hat if this notification is not
	Secti	on - III			
1. Material This Product Will Be Packaged In: Child-Resistant Packaging Yes No If "Yes" No.	7	oluble Packaging /es lo No. per	2. Type of (Container Metal Plastic Glass Paper	
* Certification must be submitted Unit Packaging wgt. cont	ainer Package	wgt container	L	Other (S	pecify)
✓ Label Container	e(s) Retail Contain 50 lb	1 4	Location of Lab	el Directio	ns
6. Manner in Which Label is Affixed to Product	Lithograph Paper glued Stenciled	Other			
	Section	on - IV			
1. Contact Point (Complete items directly below for iden	tification of individ	ual to be contacted, if	necessary, to pro	cess this	application.)
Name Robert F. Bischoff	Title Agent, To	talTurf Consulting LL	<u> </u>	Telephone (317) 873	No. (Include Area Code) -0222
Ce i certify that the statements I have made on this fo I acknowledge that any knowlingly false or mislead both under applicable law.	rtification rm end ell ettechn ling statement me	ents thereto are true, o	accurate and con or imprisonment	plete. or	6. Date Application Received (Stamped)
2. Signature Rebet F. Bischoff	3. Title Agent, Tot	alTurf Consulting LLC			() () () () () () () () () ()
4. Typed Name Robert F. Bischoff	5. Date	March 23, 201		, CLUU C C C C	((((((((((((((((((((

TotalTurf Consulting LLC 26 A. A. Cooper Rd. Meigs, GA 31765

Direct: 843-384-2807 E-Mail: rbcooper@juno.com

Fax: 229-683-3426

March 23, 2010

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505P)
U. S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attn: Linda Arrington

Subject: Shaw's 18-5-9 Turf Food with XL Crabgrass Control

EPA Reg. No. 8378-20

Notification per PR Notice 2007-1

Removal of Toll Free Number from the Disposal Instructions Section

As allowed by PR Notice 2007-1 (Section II.C.3.), the subject product label has been revised to show the removal of the statement "or 1-800-CLEANUP" from the Disposal Instructions Section. The revised label is being submitted by notification on behalf of Knox Fertilizer Company, P. O. Box 248, Knox, IN 46534 by TotalTurf Consulting LLC, an authorized agent for Knox Fertilizer Company.

The following documents are included in the subject submission:

Cover Letter

Application for Pesticide (EPA Form No. 8570-1)

One Copy of Highlighted Final Printed Label (Coded as 20100323.Final Label (PRN2007-1))

If there are any questions or additional information is needed concerning this submission, do not hesitate to contact me at (317) 873-0222 or Raymond B. Cooper at (843) 384-2807.

Sincerely,

Robert F. Bischoff

Agent

TotalTurf Consulting LLC

Robert J. Bischoff

26 A. A. Cooper Road

Meigs, GA 31765

Shaw's® 18-5-9 Turf Food NOTIFICATION with XL Crabgrass Control APR 1 5 2010

A herbicide blend for effective long lasting crabgrass control. Contains BENFLURALIN-plus-TRIFLURALIN. For Use in Lawn and Turf Maintenance Programs which call for One or More Fertilizer plus pre-emergent herbicide applications. For control of crabgrass & other annual weeds as listed.

ACTIVE INGREDIENTS:

Benfluralin; N-butyl-N-ethyl-alpha,alpha,alpha,trifluoro-2,6-dinitro-p-toluidine	0.62%
Trifluralin; Alpha,alpha,alpha,trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.30%
OTHER INGREDIENTS:	99.08%
TOTAL 1	.00.00%

EPA Reg. No. 8378-20

EPA EST. No. 8378-IN-1

Shaw's® is a registered trademark of Knox Fertilizer Company, Inc.

NET WEIGHT 50 LBS. covers up to 13,300 sq. ft.

GUARANTEED FERTILIZER ANALYSIS

Nitrogen	XXX
Ammonical (N)	XXX
Urea (N)>	(X%
Sulfur Coated Urea (N):	XX%
Phosphoric Acid (P2O5)	XX%
Soluble Potash (K2O)	XX%

Primary Plant Food derived from Urea, Sulfur Coated Urea, Muriate of Potash, Ammoniated Phosphate. Potential Acidity XX lbs. Calcium Carbonate per ton.

KEEP OUT OF REACH OF CHILDREN WARNING

FIRST AID

<u>IF IN EYES:</u> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

<u>IF ON SKIN OR CLOTHING:</u> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

<u>IF SWALLOWED:</u> Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. <u>Do not give anything by mouth to an unconscious person.</u>

<u>IF INHALED</u>: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if absorbed through the skin. Avoid contact with skin. Prolonged, or frequently repeated skin contact may cause allergic skin reactions in some individuals.

Personal Protective Equipment

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves made with any waterproof material, shoes plus socks, and goggles, face shield, or safety glasses. Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine and estuarine fish and aquatic invertebrates, including shrimp, and oysters. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes, or estuaries to aerial drift. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

6666

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Only protected handlers may be in the area during application. This product must be watered in as soon as possible after application. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Entry Restrictions

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow people (or pets) to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow people (or pets) to enter the treated area until watering-in is complete and the surface is dry.

CAN BE USED ON THE FOLLOWING ESTABLISHED TURFGRASSES

When applied according to these directions this product can be used safely on the following turfgrass varieties: PERENNIAL BLUEGRASS VARIETIES, PERENNIAL RYE GRASS VARIETIES, BERMUDAGRASS, BAHIAGRASS, CENTIPEDEGRASS, FESCUE VARIETIES, ST. AUGUSTINEGRASS AND ZOYSIAGRASS turfgrasses and other ornamental turf. Will not harm most ornamental trees; shrubs when used as directed.

WILL CONTROL THE FOLLOWING GRASSY WEEDS

This product contains a combination of Benfluralin and Trifluralin preemergence herbicides plus a lawn food. It should be applied to established turf for the control of CRABGRASS (SMOOTH AND HAIRY), GOOSEGRASS (SILVERCRAB OR CROWFOOT), BARNYARDGRASS (WATERGRASS), FOXTAIL (GREEN AND YELLOW) AND ANNUAL BLUEGRASS (POA ANNUA) while providing fertilizer nutrients to the lawn.

SEE "WHEN TO APPLY" CHART FOR APPROPRIATE TIMES TO APPLY

Apply to turf 1 to 2 weeks prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turfgrass lawns.

IMPORTANT PRECAUTIONS IN USING THIS PRODUCT

- 1. DO NOT apply in the spring to Bermuda grass which as been overseeded with winter grasses (annual rye) since thinning will occur.
- 2. DO NOT use on Dichondra, nor to newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipedegrass, nor to Bentgrass or putting greens until these turfs are well established.
- 3. DO NOT apply in spring to turfgrass planted the previous fall. This product may thin established annual Bluegrass (Poa Annua) and Fine-Leafed Fescue lawns when used at higher rates than 163 lbs. per acre (1.5 lbs. A.I. per acre). Do not overdose.
- 4. DO NOT apply more than 440 lbs. per acre per year for use on cool season turfgrasses or 65% lbs. per acre per year for use on warm season turfgrasses.
- 5. DO NOT make more than 2 applications per year.

FOR BEST RESULTS APPLY AT THE PROPER TIME

Successful preemergence control of annual grassy weeds as listed requires proper application time. It must be applied 1 to 2 weeks prior to the germination of the weed seeds. All summer annual weeds do not germinate at the same time. For this reason application timing must coincide with the normal germination period of any one of the annual grassy weeds. If not, poor and erratic weed control will result due to improper application.

WATER THE TREATED AREA AFTER APPLICATION

Best results will be obtained if treated areas are irrigated with $\frac{1}{4}$ to $\frac{1}{2}$ inch water soon after application. Any cultural practices that disturb the soil such as verticutting, should be done prior to application of this product.

When to use lower and higher application rates in the following directions for use. In general, use the lower rates (i.e. 1.5 or 2 lbs. active ingredient/acre) when normal weed seed germination is expected. Use the higher rates (i.e. 2 or 3 lbs. active ingredient/acre) when weed seed germination is expected to be heavy.

"SPRING GERMINATING"

ANNUAL GRASSES: CRABGRASS, FOXTAIL, BARNYARDGRASS

- A. Northern Turf use 50 lbs. to treat 10,000 to 13,300 sq. ft. for Northern cool season turfgrasses For pre-emergence control, apply 163 to 220 lbs. per acre, (equal to 1.5 to 2 lbs. active ingredient/acre) applied in the spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 50 lb. bag will cover 10,000 to 13,300 sq. ft. Use for 10,000 sq. ft. when turf is completely overrun with weeds. Use for 13,300 sq. ft. when weeds are sparse and the turf is in a yearly lawn maintenance program.
- B. <u>Southern Turf</u> use 50 lbs. to treat 6,700 to 10,000 sq. ft. For Southern warm season turfgrasses For pre-emergence control apply 220 to 325 lbs. per acre (equal to 2 to 3 lbs. active ingredient per acre), in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination. One 50 lb. bag will cover 6,700 to 10,000 sq. ft. Use for 6,700 sq. ft. where turf is completely overrun by weeds. Use for 10,000 sq. ft. when weeds are sparse or the turf is in a yearly crabgrass control program.

"LATE SPRING-EARLY SUMMER GERMINATING" ANNUAL GOOSEGRASS (SILVER CRAB)

Although classified as a summer annual grass, Goosegrass normally germinates 4 to 6 weeks later than the crabgrass, foxtails and other summer annual grass species. Applications of this product should therefore, be timed accordingly.

A. Northern Turf - use 50 lbs. to treat 10,000 sq. ft. for Northern cool season turfgrasses - For preemergence control apply 220 lbs./acre (equal to 2 lbs. active ingredient per acre) of this product in late spring or early summer one to two weeks prior to the expected garmination period of Goosegrass.

 ϵ

B. <u>Southern Turf</u> - use 50 lbs. to treat 6,700 to 10,000 sq. ft. for Southern warm season turfgrasses - For preemergence control, apply 220 to 325 lbs. per acre (equal to 2 to 3 lbs. active ingredient per acre) of this product in late spring or early summer one to two weeks, prior to the expected germination of goosegrass. Use for 6,700 sq. ft. where turf is completely overrun by weeds. Use for 10,000 sq. ft. where weeds are sparse and the area is in a yearly lawn maintenance program.

APPLY TWICE FOR FULL SEASON CONTROL

In areas where preemergence control of both early germinating annual grasses (Crabgrass, Foxtail, and Barnyardgrass) and the later germinating Goosegrass is desired, two application of this product at the recommended rate, spaced 8 to 10 weeks apart, may be made to provide full season control of these annual grass species.

POA ANNUA (ANNUAL BLUEGRASS) CONTROL

In areas of heavy infestation of Annual Bluegrass, its elimination will leave areas of thin turf. Employ proper cultural practices to establish a new turf.

- A. Northern Cool Season Turfgrasses For preemergence control apply 220 lbs. per acre (equal to 2 lbs. active ingredient per acre) of this product per acre in late summer or early fall and again in late winter or spring one to two weeks prior to expected germination of annual bluegrass. One 50 lb. bag will treat 10,000 sq. ft. of turf grass.
- B. Southern Warm Season Turfgrasses For preemergence control apply 325 lbs. per acre (equal to 3 lbs. active ingredient per acre) of this product per acre in the later summer or early fall one to two weeks prior to expected germination of poa annua. In southern areas where germination continues for extended periods of time a second application at 325 lbs. per acre can be made 6 to 8 weeks after the initial treatment. One 50 lb. bag will treat and feed 6,700 sq. ft. of turfgrass.

TO RESEED TREATED AREAS

- 1. Delay reseeding 6 weeks when applying at the rate of 163 to 220 lbs./acre equal to 1.5 to 2.0 lbs. active ingredient/acre, or when using one 50 lb. bag to cover 10,000 to 13,300 sq. ft.
- 2. Delay reseeding 12 to 16 weeks when applying at rate of 325 lbs./acre equal to 3 lbs. of active ingredient per acre or when using one 50 lb. bag to cover 6,700 sq. ft. of turfgrass. When reseeding it may be wise to increase the amount of seed put down by 10%.

3.

WHEN TO APPLY

*Approximate "times of the year" when crabgrass etc. weed seeds germinate.

JAN. FEB. MAR. APR. MAY JUNE JULY AUG. SEPT. OCT. NOV. DEC.
NORTHERN TURF

1.///// 2.///// 3.///////
SOUTHERN TURF

4.///////// 5./////// 6.////

Soil moisture % & temperature degrees will change germination periods each year.

1. For crabgrass, foxtail, barnyardgrass in Northern Turf.

- 2. For goosegrass-Northern Turf.
- 3. For poa annua in Northern Turf.
- 4. For crabgrass, foxtail and barnyardgrass in Southern Turf.
- 5. For goosegrass-Southern Turf.
- 6. For poa annua in Southern Turf.

Two applications per year are usually required to control weeds when germination periods are extended.

SPREADER SETTINGS

The following spreader settings are for use with new equipment. Old spreaders may need recalibrating. Proper application is based on equipment condition with regard to age and use, also your walking speed and unevenness of the terrain. Do not overlap coverage. Poor control or turf damage can result from uneven or improper application.

Spreader	50# bag per sq. ft.	Northern Lawns Crabgrass etc. Poa Annua	Southern Lawns Crabgrass etc. Poa Annua
Scott #35	13,300	XX	XX
	10,000	XX	XX
	6,700	XX	XX
Cyclone B1 13,	13,300	XX	XX
•	10,000	XX	XX
	6,700	XX	XX

50 lbs. to treat 13,300 sq. ft. = 163 lbs./acre = 1.50 lbs. A.I./acre

50 lbs. to treat 10,000 sq. ft. = 220 lbs./acre = 2.00 lbs. A.I./acre

50 lbs. to treat 6,700 sq. ft. = 325 lbs./acre = 3.00 lbs. A.I./acre

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. <u>PESTICIDE STORAGE</u>: Store in original, closed container only. <u>PESTICIDE DISPOSAL</u>: <u>IF EMPTY</u>: Do not reuse this bag. Place in trash or offer for recycling if available. <u>IF PARTLY FILLED</u>: <u>Call your local solid waste agency for disposal instructions.</u> Never place unused product down any indoor or outdoor drain. <u>CONTAINER DISPOSAL</u>: Nonrefillable container. Do not reuse or refill this container, offer for recycling if available.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under acceptancy Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the labels and is reasonably fit for the purposes stated on the labels when used in strict accordence with the δ

^{*}Apply 1 to 2 weeks before seed germinate in your locale.

^{**}Two applications may be needed during this period to give adequate control.

directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, manner of application or other factors, all of which are beyond the control of the Seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount or product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller promptly notifies of such loss or damage in writing. To the extent consistent with applicable law, Seller shall not be liable for consequential or incidental damages or losses.

The terms of this Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

KNOX FERTILIZER COMPANY, INC.
P.O. BOX 248
KNOX, IN 46534
NET WEIGHT 50 LBS. (22.7 kg.)

Batch No.

