8378-23

20030220.amended label

07/15/2003


Shaw's Premium Weed & Feed "Formula 1"

Treats XX sq. ft.

This product is a premium grade herbicide and turf food combination which controls a wide spectrum of broadleaf weeds while providing both fast and long lasting action. This product may be used on bluegrass, fescue, rye, Bermuda, Bania, Zoysia and St. Augustinegrass lawns.

ACTIVE INGREDIENTS: *Dimethylamine Salt of

Dimenty drame Juli of	
2,4-Dichlorophenoxyacetic acid	0.313%
***Dimethylamine Salt of (+)-R-2-(2-Methyl	-4-
chlorophenoxyl) propionic acid	0.157%
****Dimethylamine Salt of (+)-R-2-(2,4-Dich	loro-
phenoxy) propionic acid	0.155%
INERT INGREDIENTS:	<u>99.375%</u>
TOTAL	100.00%
EPA Reg. No. 8378-23 EPA Est.	No. 8378-IN-1
Isomer Specific AOAC Method No.	6.321, Equivalent to:
*0.26% of 2,4-Dichlorophenoxyacetic Acid	•
**0.13% of (+)-R-2-(2-Methyl-4-Chlorophene	xy) propionic acid
***0.13% of (+)-R-2-(2,4-Dichloropheonxy) p	



TCONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product

container or label with you when calling a poison control center or doctor or when going for treatment.

You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation of dust and contact with eyes, skin or clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Do not allow people (other than applicator) or pets on treatment area during application.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or to areas where surface water is present or to interdial areas below the mean highwater mark. Do not contaminate water when disposing of equipment washwaters.

RE-ENTRY STATEMENT

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL	
STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contamina	ite water, foodstuffs, feed, or
seed by storage or disposal.	•
PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended	that alrof the contents of the bag
be used, carefully following label directions and precautions.	• • • • •
IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available.	••••
IF PARTLY FILLED: Call your local waste agency or 1-800-CLEANUP for disposal instructions. Neve indoor or outdoor drain.	er place unused product down any
<u>SPILL:</u> In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directi	er * * * *
- TALE IN CASE OF SPIN, Sweep of and dispose of Aldendard and to robot of DISPOSAL direction	

IMPORTANT: Read entire label before using this product.

Shaw's PREMIUM WEED & FEED Formula 1 is a stable mixture of three phenoxy herbicides. It is for use on Lawns, Ornamental Turf (not pastures), Golf Fairways, Aprons, Tees and Roughs. This product may be used on Bluegrass, Fescue, Rye, Bermuda, Bahia, Zoysia, and St. Augustinegrass lawns. DO NOT USE on Dichondra and Carpetgrass. DO NOT USE this product where desirable clovers are present or allow this product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

This product is designed for use in lawns for control of broadleaf weeds such as Black Medic, Buckhorn, Burdock, Chickory, Chickweed, Clover, Dock, Dandelion, English Daisy, Ground Ivy, Heal-All, Knotweed, Lambsquarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Ragweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Onion, Wild Lettuce, Wild Garlic, Yarrow and other similar broadleaf weeds.

NORTHERN GRASSES

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. IMPORTANT: Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf killing action, avoid unnecessary disturbances, including water, on treated areas for 48 hours after application. Watering will wash off weed killing material. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawns until after 3 mowings.

SOUTHERN GRASSES

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in the spring. IMPORTANT: Apply when grass is moist (after rain, sprinkling or dew). DO NOT water lawns after material is applied, as watering will wash off weed killing material. For most effective broadleaf killing action, avoid unnecessary disturbances of treated areas for 48 hours after application. Avoid contact with valuable shrubs, plants, or flowers. Possible retreatment may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not over apply. Under conditions of adequate moisture, high temperature, and high humidity, over-application may cause minor transitory discoloration of Centipede and St. Augustinegrass. CAUTION: It is recommended that during hot, dry conditions, application rates should be reduced by one half to avoid possible dehydration or browning of grass. Do not reseed for at least three or four weeks after application. Do not apply to newly seeded or sodded lawns until after three mowings.

HOW TO APPLY: The maximum rate to turf is 2 pounds of 2,4-D acid equivalent per acre, per application site. The maximum number of broadcast applications per treatment site is 2 per year. Apply evenly with a lawn spreader either as one pass application or a two pass application, criss-crossing the area using a half rate each time. Lawns that are heavily over-run with weeds and are in poor condition may require more than one application of this product. A suggested second application should be made in approximately 30 days.

SPREADER SETTINGS

The following settings are approximate for applying contents of this package. Settings are based on average walking speed. Reduce setting for slower speed and raise setting for higher speed.

Cyclone xx-xx K-Mart xx-xx

Scotts xx-xx Montgomery Wards xx-xx

Central xx-xx

These settings are approximate. Check your spreader on 250 sq. ft. and adjust at the rate of 0.9 pound per 250 sq. ft. Clean spreader thoroughly after each use.

GUARANTEED FERTILIZER ANALYSIS

418-5----

......

TOTAL NITROGEN(N)	XX%		1
AVAILABLE PHOSPHATE(P2O5)	XX%		• • •
SOLUBLE POTASH (K2O)	XX%		∎ • 4 v ·
Primary plant foods derived from: u	rea, ammoniated phosphate and muriate of potash.	•••••••	•
	l risk and liability of use, storage and/or handling of this	material not is ac	
terms of this label.		• •	16 × 3 × 17
	NET WEIGHT XX POUNDS	• • • •	• • • •
	KNOX FERTILIZER COMPANY, INC.		*
	P.O. BOX 248		
	KNOX, IN 46534		\$ /
			* *
			, د . •