toric failer,

SHAW'S

#### PREMIUM WEED AND FEED

## FORMULA 3

FOR USE ON HOME LAWNS, PARKS, GOLF COURSES AND OTHER RECREATIONAL TURF AREAS TREATS 5,000 SQ. FEET (50'  $\times$  10')

THIS PRODUCT IS A PREMIUM GRADE TURF FOOD AND HERBICIDE COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG LASTING ACTION

## ACTIVE INGREDIENTS:

Isooctyl Ester of 2-(2-Methyl-4-chlorophenoxy)propionic	Acid*	1.0%
Isooctyl Ester of 2-Methyl-4-chlorophenoxyacetic Acid**		1.0%
INERT INGREDIENTS:		. 98.0%
	<b>ጥ</b> ንፑል፣	<u> 100 09</u>

## BY ISOMER SPECIFIC AOAC ASSAY, EQUIVALENT TO:

\* .62% of 2-(2-methyl-4-chlorophenoxy)propionic Acid

\*\* .60% of 2-methyl-4-chlorophenoxyacctic Acid

## GUARANIEED ANALYSIS:

Total Nitrogen (N)

1.0% Ammoniacal Nitrogen

24.0% Urea Nitrogen
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>)

Soluble Potash (K<sub>2</sub>O)

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Grea, Urea, Triple Super Phosphate, and Muriate of Potash

Chlorine (Cl) Maximum ..... 2.5%

The urea nitrogen in this product has been coated to provide 6.8% coated slow release urea nitrogen.

## KEEP OUT OF REACH OF CHILDREN

## CAUTTON

IF SWALLOWED: Get Medical Attention.

IF IN EYES: Flush

Flush Eyes with Water for 15 Minutes and Get Medical Attention.

IF ON SKIN: Wash Skin with Soap and Water.

SEE REAR PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 8378-

EPA. EST. NO.8378-5N-1

#### MANUFACTURED BY

KNOX FERTILIZER AND CHEMICAL COMPANY, INC. Knox, Indiana 46534

NET WEIGHT - 18 Lbs.

M 23

8378-25

(Rear Panel)

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTTON

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly after handling. Remove contaminated clcthing and wash before reuse.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when conditions favor drift from target areas.

Shaw's Premium Weed and Feed - Formula 3 is a remarkably effective lawn fertilizer which combines both fast and long lasting feeding action plus effective weed control. Weeds contacted by this product gradually die and disappear from the lawn. It is for use on lawns and ornamental turf.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL before using this product.

This product may be used on Bahia, Bermuda, Blue Grass, Fescue, Rye, and Zoysia Lawns. USE CAUTION: Some bent varieties such as Arlington, Congressional and Seaside may be injured and some injury may occur to St. Augustine. DO NOT USE on Dichondra or where desirable clovers are present. Do not allow this product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees, or other desirable plants.

This product was designed for control of broadleaf weeds in lawns such as: Beggarstick, Bindweed, Black Medic, Burcucumber, Burdock, Buttercup, Canadian Thistle, Carpetweed, Catsear, Chervil, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dragon Heat Mint, Fat Hen, Fennel, Gelinsoga, Goatsbeard, Goldenrod, Ground Ivy, Halbert-leaved Salt Brush, Hempnettle, Henbit, Hoary Cress, Honeysuckle, Jimson Weed, Kochia, Knotweed, Lambsquarter, Marsh Elder, Merkican Poppy, Morning Glory, Mustard, Nettles, Pennycress, Peppergrass, Pepperweed, Pigweed, Plantain, Poison Hemlock, Poison Ivy, Poison Oak, Prickly Lettuce, Puncturevine, Purslane, Ragweed, Red Root, Redstem, Shepard's Purse, Sticklepod, Sneezeweed, Sow Thistle, Spanish Needle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Vetch, Whitebrush, White Top, Wild Carrot, Wild Gooseberry, Wild Jute, Wild Marigold, Wild Petunia, Wild Radish, Wild Sage, Wild Garlic, Wintercress, Witchweed, Yellow Charlock, Yellow Daisy, Yellow Rocker,

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. IMPORTANT: Apply when grass is moist (after a rain, sprinkling or dew).

For most effective broadleaf weed killing action avoid unnecessary disturbance, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not reseed for at least 3 to 4 weeks after application.

Do not apply to newly seeded or sodded lawn until after 3 mowings.

This 18 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreadar, either as one pass application or a two pass application, criss-crossing the area at half the rate each time. The following settings are approximate for applying contents of this package at the above suggested rate. Check your spreader on 250 square feet and adjust to apply at the rate of .9 lb. per 250 square feet. Settings can vary with type, age and condition of spreader. Settings are based on an average walking speed. Reduce

(Rear Panel cont.)



setting for slower speed; raise setting for higher\_speed. Clean spreader thoroughly after use.

## SPREADER SETTINGS

MAKE	SETTING'	. MAKE	SETTING
Cyclone	5 1/4 - 5 3/4	Montgomery Wards	
Scotts 35	5 1/2	1091, 1093, 1095	6 3/4
Scotts C-1	J - L	Montgomery Wards	
Central	7 - 8	$108\overline{0}$ , $1081$	5
K-Mart	7 - 8	Precision	6

#### STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in an area inaccessible to children and domestic animals. Do not store near food or food-stuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

WARRANTY: Knox Fertilizer and Chemical Company warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.



