JUL 1 2 1995

Knox Fertilizer Company, Inc. c/o Mr. H. R. McLane H. R. McLane, Inc. 7210 Red Road, Suite 206 Miami, Florida 33143-5321

Subject: Change of Product Name--Revised Labeling
Dursban 92 with Plant Food
(Formerly Dursban 92 + Fertilizer)
EPA Registration Number 8378-26
Your Application Dated March 25, 1994
and Submission Dated April 14, 1995

Dear Mr. McLane:

We have noted that you previously requested that the official name of record of this product be changed to

Dursban 92 with Plant Food.

The change is acceptable and we have emended our records to include the new name.

The labeling referred to above, originally included with your application dated March 25, 1994, and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

Revise the last sentence in the Environmental Hazards section to read: "Do not contaminate water by cleaning of equipment or disposal of waste."

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr. Product Manager (19) Insecticide-Rodenticide Branch Registration Division (7505C)

Enclosure



## DURSBAN®92 with Plant Food

50 LBS TREATS & FEEDS UP TO 20,500 SQ.FT. OF LAWN

20-5-10 LAWN FOOD LAWN AND TURF INSECT CONTROL

KILLS SOD WEBWORMS, WHITE GRUBS, ANTS, FLEAS, BROWN DOG TICKS, CRICKETS, CHINCH BUGS, CUTWORMS, EARWIGS, SOWBUGS, AND ARMYWORMS EUROPEAN CRANE FLY LARWAE HYPERODES (TURFGRASS) WEEVIL IN HOME LAWNS AND ORNAMENTAL TURF with COMMENTS in EPA Letter Dated

**ACTIVE INGREDIENT:** 

Chlorpyrifos: [O,O,diethyl O(3,5,6 trichloro-2-pyridyl) phosphorothioate]. 0.92% INERT INGREDIENTS:

Total DURSBAN® is a Registered Trademark of The DowElanco Products Company

> KEEP OUT OF REACH OF CHILDREN CAUTION

JUL 1 2 1995

99.08 6
100.00% Under the Federal Insceticide, mpany Fungicide, and Rodenticide Act as amended, for the peaticide registered under EPA Reg. No.

STATEMENT OF PRACTICAL TREATMENT

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing and wash affected area with soap & warm water. Wash clothing before reuse, IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention immediately. NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. May be injurious to eyes and skin. Do not get on skin or in eyes. Wash thoroughly after handling. Do not wear contaminated clothing. Wash contaminated clothing before reuse. Avoid breathing dust. Keep away from food, feedstuffs, and water supplies. Where watering-in is required keep children and pets off treated areas until grass is dry. ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment of disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal, STORAGE: Store product in its original container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposed from the use of this product may be disposed of on site or at an approved waste disposed from the use of this product may be CONTAINER DISPOSAL; Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. HOUSEHOLD: Do not reuse empty container. Wrap container and put in trash.

**GUARANTEED FERTILIZER ANALYSIS** 

NITROGEN (N) ...... UREA (N) variable
AMMONIATED NITROGEN variable % variable % AVAILABLE PHOSPHORIC ACID (P2O5) variable % SOLUBLE POTASH (K2O) .... variable % Derived from Urea, Ammoniated Phosphale, Muriate of Potash. Potential acidity =

Calcium Carbonate per ton.

## LAWNS AND ORNAMENTAL TURF SURFACE FEEDING INSECTS

For control of Chinch Bugs, Sod Webworms (Lawn Moth), Ants, Fleas, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs and Millipedes apply 50 lbs. to treat 20,500 Sq.Ft. (refer to spreader settings.) Treat in the spring or when insects first appear troublesome and repeat as needed for control. Additional treatments at 30 day intervals may be needed for season long control of Sod Webworms. woms

FOR ROOT FEEDING INSECTS

For Billbug Larvae, and the "White Grub" larvae of (Japanese Beetle, European Chafer, Southern Chafer), and Hyperodes Weevil apply 50 lbs. to treat 10,250 Sq.Ft. of lawn.

Sq.Ft. or lawn.

<u>European Crane Fly larvae</u> in turf will be controlled at this same rate, (50 lbs/10,250 Sq.Ft.), with an individual application or if present, controlled simultaneously with the same application being applied for "White Grubs". Consult your Agricultural Extension Agent for proper time to apply.

For the <u>Hyperodes (Turfgrass) Weevil</u> apply in mid-April and again in mid-May using 50 lbs. per 10,250 Sq.Ft. (refer to the spreader settings chart). Apply uniformly over the area and immediately follow with a thorough watering to wash the insecticide into the soil where insects are feeding. For best control apply when the

insecticide into the soil where insects are feeding. For best control apply when the 'grubs" are young and actively feeding near the soil surface in mid July & early August.

**IMPORTANT** 

The following spreader settings are offered for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreading equipment may require adjustments from the following figures. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 Sq.Ft. into your spreader and apply to your premeasured 500 Sq.Ft.(5 ft. wide by 100 ft long). Faulty and/or erratic application will result in poor insect control!

## SPREADER SETTINGS

SPREADER	AMOUNT TO APPLY	WEBWORMS AND OTHER SURFACE FFEDING INSECTS SO IF PER 20,502 SQ FT 2,5 lbs / 1,000 Sq (2,	N'HTE GRUBS AND ROOT FEEDING INSECTS 30 # PER 10,250 SO FT. 5 0 lbs /1,000 SQ FT.
Scott	50 pounds	4.5	#6
Cyclone	50 pounds	4.5	

NOTICE: Buyer assumes all risk and liability of use storage and/or handling of this malerial not in accordance with the terms of this label (8378-26.PM5 \ 03-21-94)

EPA REG. NO. 8378-26 NET WEIGHT 25 POUNDS (50 LB.) EPA EST. NO. 8378-IN-1

KNOX FERTILIZER COMPANY, INC. KNOX, INDIANA 46534