

8378-39

1/21/2003

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Robert F. Bischoff
Authorized Agent for Knox Fertilizer Company
TotalTurf Consulting LLC
P.O. Box 7928
Hilton Head, SC 29938

JAN 21 2003

Knox Fertilizer Company
P.O. Box 248
Knox, IN 46534

Dear Mr. Bischoff:

Subject: Shaw's Turf Food/Crabgrass Control Barricade® 20
EPA Reg. No. 8378-38
Shaw's Turf Food/Crabgrass Control Barricade® 28
EPA Reg. No. 8378-39
Shaw's Turf Food/Crabgrass Control Barricade® 38
EPA Reg. No. 8378-40
Shaw's Turf Food Crabgrass Control Barricade® 50
EPA Reg. No. 8378-56
Applications and Letters Dated December 11, 2002,
Requesting Amendments To Registrations: To Comply
with EPA PR-Notice 2001-1 and 2001-6, by Updating First
Aid Statements and Storage and Disposal Statements,
Respectively; and To Revise Spreader Setting Tables,
Revise Crabgrass Seed Germination Dates Sections,
Amend Container Sizes and Amend Warranty and Liability
Sections

The propose labels submitted with your applications and letters dated December 11, 2002 have been reviewed and found acceptable as amendments to the registration of the subject pesticide products under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, provided that you:


1. Delete the word "Selective" used to describe these products, on front panels and in text of all labeling.
2. Move the "Application Directions" section to the end of "Directions of Use" and follow the statement: "Apply uniformly with suitable, calibrated application equipment." with a heading: "Spreader Recommendations" and the table of spreaders. Follow the table of spreaders with: "Calibrate your spreader according to the manufacturer's directions. Initial spreader

setting may require adjustment to deliver the recommended Application rate under actual application conditions. Apply this product uniformly over the area to be treated. Avoid streaking, skips and overlaps during application. Calibration of spreader may be needed if conditions of area to be treated changes and rate or speed of application are changed".

3. On all supplemental labeling revise the "Application Directions" section by adding the statement: "See "Spreader Recommendations" section for spreader setting and calibration instructions."
4. Revise the claim under "Tolerant Ornamental Species" that reads: "This product may be applied over the top of these species." by modifying the claim to read: "....., provided spray residue on shoots of plants is washed to the soil by over the top irrigation or by adequate rain soon after application".
5. Submit one (1) copy of each of the final printed labels prior to your shipment of these pesticide products under the enclosed revised stamped labels.

If these conditions are not complied with, these registrations will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment under these labels constitutes acceptance of these conditions. A stamped copy of each these labels are enclosed for your records.

Sincerely yours,


for Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure (8)

3/12

Shaw's Turf Food/Crabgrass Control Barricade® 28

For Selective Preemergence Control of Grass and Broadleaf Weeds in Established Turf Grasses, (excluding golf course putting greens) and Lawns - Also for Landscape Ornamentals Including Established Perennials and Wildflower Plantings. (See Supplemental Label)

ACTIVE INGREDIENT:

Proflaminate [N³,N³-Di-*n*-propyl-2,4-dinitro-6-(trifluoromethyl)-*m*-phenylenediamine] 0.28%

INERT INGREDIENTS: 99.72%

TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

Barricade® is a registered trademark of Syngenta

EPA Reg. No. 8378-39

EPA Est. No. 8378-IN-1

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	XX%
Available Phosphoric Acid (P2O5)	XX%
Soluble Potash (K2O)	XX%

Ingredients: Ureaformaldehyde, Urea, Sulfur Coated Urea, Ammonium Phosphate, Muriate of Potash

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. No not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center, or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides. PRODUCT

DISPOSAL: IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. If product is spilled, sweep up and use as per label instructions or dispose of as indicated above.

4/12

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is a turf fertilizer and a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least $\frac{1}{2}$ inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown: (1) for sale or other commercial use; (2) for commercial seed production or; (3) for research purposes. For use on plants intended for esthetic purposes or climatic modification and being grown in ornamental gardens, or parks, or on golf courses or lawns and grounds. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.


Application Directions

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least $\frac{1}{2}$ inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.

Crabgrass Seed Germination Dates



Approximate Date	
	After May 30
	After May 10
	After April 20
	After March 20
	January 1 to March 20

USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

5/12

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year Of Product by Turf Grass Species ⁽¹⁾		
Turf Species	Lbs. Product/A	Lbs. a.i./A
Creeping Bentgrass	233 lbs.	0.65 lb.
Creeping Red Fescue	267 lbs.	0.75 lb.
Buffalograss	357 lbs.	1.00 lb.
Kentucky Bluegrass	"	"
Perennial Ryegrass	"	"
Bermudagrass ⁽²⁾	535 lbs.	1.50 lb.
Bahiagrass	"	"
Centipedegrass	"	"
Seashore Paspalum	"	"
St. Augustinegrass	"	"
Tall Fescue (including turf type)	"	"
Zoysia	"	"

/A= per acre

(1) These are the maximum rates per calendar year by species limitations. (2) May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 180 lbs./A. Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations. Do not apply more than 535 lbs. of this product per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates section.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Henbit	Purslane, common
Bluegrass, Annual (Poa Annua)	Itchgrass	Pursley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass ⁽³⁾
Chickweed, Common	Junglerice	Shepherd's Purse ⁽²⁾
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (Large, Smooth)	Kochia	Speedwell, Persian
Crowfoot grass	Lambsquarter, common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Witchgrass
Goosegrass ⁽¹⁾	Pigweed	Woodsorrel, Yellow (from seed)

(1) In many areas a single application of 233 to 535 lbs/A will control goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 233 to 357 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate section. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate section. (2) Application for this weed should be made in late summer, fall, or winter prior to germination. (3) Suppression Only. Sequential application may be made as long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Spreader settings for 1000 sq. ft.	180 lbs. Product/A	267 lbs. Product/A	357 lbs. Product/A	535 lbs. Product/A
Scott's Evergreen 18"	XX	XX	XX	XX
Scott's Speedy Brdct.	XX	XX	XX	XX
Red Devil Broadcast	XX	XX	XX	XX
Republic EZ Drop 18"	XX	XX	XX	XX
Republic EZ Broadcast	XX	XX	XX	XX
Republic EZ Trctr. Brdct.	XX	XX	XX	XX

6/12

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Product rate lbs./Acre		Months Before Overseeding		
Product	a.i./A	North	Transition	South
180	0.50	4	4	4
233	0.65	5	4	4
267	0.75	6	5	5
285	0.80	-	6	6
357	1.00	-	7	7
407	1.14	-	-	9
465	1.30	-	-	10
535	1.50	-	-	12

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. If these terms are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid.

***FOR USE ON LANDSCAPE ORNAMENTALS**

(including Established Perennials and Wildflower Plantings). This product may be applied for residual preemergence weed control in ornamental plantings.

FOR COMPLETE DETAILS AND DIRECTIONS FOR USE ON ORNAMENTAL PLANTS, REFER TO THE SUPPLEMENTAL LABEL FOR USE OF THIS PRODUCT ON LANDSCAPE ORNAMENTALS.

NET WEIGHT XX LBS.
KNOX FERTILIZER COMPANY, INC.
P.O. BOX 248
KNOX, IN 46534

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JAN 21 2003

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

8378-39

7/12

Supplemental labeling

Shaw's Turf Food/Crabgrass Control Barricade® 28

For Selective Preemergence Control of Grass and Broadleaf Weeds in Landscape
Ornamentals Including Established Perennials and Wildflower Plantings.

ACTIVE INGREDIENT:

Prodiamine [N3,N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] 0.28%

INERT INGREDIENTS: 99.72%

TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA REG. NO. 8378-39

EPA EST. NO. 8378-IN-1

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.
Barricade® is a registered trademark of Syngenta

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) XX%
Available Phosphoric Acid (P2O5) XX%
Soluble Potash (K2O) XX%
Ingredients: Ureaformaldehyde, Urea, Sulfur Coated Urea, Ammonium Phosphate, Muriate of Potash

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. No not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center, or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides. **PRODUCT**

DISPOSAL: IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **If product is spilled,** sweep up and use as per label instructions or dispose of as indicated above.

8/12

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is a turf fertilizer and a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, landscape ornamentals, established perennials and wildflower plantings. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least $\frac{1}{2}$ inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown: (1) for sale or other commercial use; (2) for commercial seed production or; (3) for research. For use on plants intended for esthetic purposes or climatic modification and being grown in ornamental gardens, or parks, or on golf courses or lawns and grounds. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

Application Directions

Apply uniformly through suitable, calibrated, hand held, push or pull type granular spreader application equipment.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. The length of time of residual weed control provided by this product is related to the rate applied.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Henbit	Purslane, common
Bluegrass, Annual (Poa Annua)	Itchgrass	Pursley, Florida
Carpeweed	Johnsongrass	Rescuegrass ⁽³⁾
Chickweed, Common	Junglerice	Shepherd's Purse ⁽²⁾
Chickweed, Mauseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (large, smooth)	Kochia	Speedwell, Persian
Crowfoot grass	Lambsquarter, common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Witchgrass
Goosegrass ⁽¹⁾	Pigweed	Woodsorrel, Yellow (from seed)

(1) In many areas a single application of 233 to 535 lbs./A will control goosegrass. However, under goosegrass pressure and/or on extended growing season, most effective weed control may be obtained by making an initial application of 233 to 357 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed a total of 535 lbs. per acre, per season. (2) Application for this weed should be made in late summer, fall, or winter prior to germination. (3) Suppression Only. Sequential application may be made as long as the total amount of product applied does not exceed the maximum annual application rates of 535 lbs. per acre. All applications must be made prior to germination of the weed seeds.

LANDSCAPE ORNAMENTALS

Including Established Perennials and Wildflower Plantings

This product may be applied for residual preemergence weed control in ornamentals.

Use Rates

Apply at 5.35 to 12.25 lbs./1,000 sq. ft. (equiv. To 233 to 535 lbs./A) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 535 lbs./acre.

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product into the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

9/12

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

<u>Scientific Name</u>	<u>Common Name</u>
Abies spp.	Fir species** (Balsam, Fraser, Noble, etc.)
Abelia grandiflora	Abelia: Sherwood
Acer palmatum	Japanese Maple
A. platanoides	Norway maple
Achillea, sp.	Yarrow: King Edward
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebia quintata	Five-Leaf or Chocolate vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia sp.	Aquilegia; Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemisia sp.	Wormwood; Silver Mound, Castle
Aster sp.	Aster: Bonny Blue, Purple Dome
Aster X Frikartii	
Athrium Filiz-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia sp.	Fibrous Begonia; Hardy Grandis
Berberis gladwynensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. thunbergii	Japanese Barberry
B. Verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Bougainvillea spp.	Bougainvillea
Buddleia davidii	Butterfly-bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Creeper, Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Chrysanthemum nipponicum	
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis sp.	Coreopsis (Calliopsis); Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaeus spp.	Hawthorne
Crococsmia sp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma sp.	Cooperi Pink

10/12

<u>Scientific Name</u>	<u>Common Name</u>
Delphinium sp.	Larkspur; Blue Elf
Dianthus deltoides	Dianthus; Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea pupurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Japanese Spindle Tree (Evergreen Euonymus)
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia sp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine, Radicans
Gaura sp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.	Gladiolus species**
Gypsophila repens	Baby's Breath
Hedera helix	English Ivy
Hellanthenum sp.	Sunrose
Hemerocallis sp.	Daylily: Aztec Gold, Stella De Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus sp.	Mallow: Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Searsucker"
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly; Helleri
I. opaca	American Holly
I. pernyi	Holly
I. vomitoria	Yaupon Holly; Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet
Iris spp.	Iris species**
Jasminum nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. Conferta	Shore Juniper; Blue Pacific
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans sp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender sp.	Lavender; Munstead
Leontopodium aplanum	Edelweiss
Ligustrum sinense	Chinese Privet; Variegata
Ligustrum amurense	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (waxleaf)
Lillum sp.	Lily: Jazz
Liriope muscari	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
Liriope Muscari var. variegata	Liriope, Variegated
Liliope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink

11/12

Scientific Name

Lonicera japonica
 L. tatarica
 Loropetalum chinense
 Lythrum sp.
 Magnolia spp.
 Maleophora luteola
 Malus sp.
 Miscanthus sinensis
 Nandina domestica
 Narcissus spp.
 Nerium oleander
 Olea europaea
 Ophiopogon japonicus
 Osmanthus heterophyllus
 Osteopermum fruticosum
 Oxydendrum arboreum
 Paeonia suffruticosa
 Pennisetum setaceumi
 Perovskia atriplicifolia
 Persea americana
 Photnia fraseri
 Physotegia Virginiana
 Picea spp.
 Pieris japonica
 Pinus brutia
 P. canariensis
 P. eliottii
 P. halepensis
 P. nigra
 P. palustrus
 P. radiata
 P. thunbergiana
 P. strobus
 P. sylvestris
 P. taeda
 P. virginiana
 Pistachio sp.
 Pittosporum rhombilfolium
 P. tobira
 Podocarpus macrophyllus
 Prunus laurocerasus
 Prunus sp.
 Pseudotsuga menziesii
 Pyracantha coccinea
 P. fortuneana
 P. koidzumii
 Pyrus sp.
 Quercus Shumardii
 Quercus spp.
 Raphiolepis indica
 Raphiolepis umbellata
 Rhododendron
 (including Azalea)

Common Name

Japanese Honeysuckle
 Tatarian Honeysuckle
 Burgundy
 Loosestrife; Modern Pink
 Magnolia species**
 Ice Plant
 Crabapple*
 Yaku Jima, Silberfeder
 Heavenly Bamboo
 Narcissus species**
 Oleaner
 Olive*
 Mondo Grass
 Osmanthus (False Holly): Gulf Tide
 Trailing African Daisy
 Sourwood
 Tree Peony
 Fountain Grass (Dwarf)

 Avocado*
 Frasier's Photinia (Redtip)
 Dragonhead, False; Vivid
 Spruce species** (Colorado Blue, Norway, etc.)
 Lily-of-the-Valley Bush
 Calabrian Pine
 Canary Island Pine
 Slash Pine
 Aleppo Pine
 Austrian Pine
 Longleaf Pine
 Monterey Pine
 Japanese Black Pine
 Eastern White Pine
 Scotch Pine
 Loblolly Pine
 Virginia Pine
 Pistachio*
 Queensland Pittosporum
 Japanese Pittosporum
 Japanese Yew
 English Laurel
 Almond, Apricot, Nectarine, Peach, Plum and Prune*
 Douglas Fir**
 Firethorn (Scarlet)
 Firethorn
 Firethorn
 Bradford Pear sp.
 Oak, Shumard's Red
 Oak species
 Indian Hawthorne
 Yedda Hawthorn
 'Coral Bells'
 'Delaware Valley White'
 'Flame Creeper'
 'Formosa'
 'Girard Crimson'
 'George L. Tabor'
 'Hino-crimson'
 'PJM'
 'Roseum Elegans'

12/12

<u>Scientific Name</u>	<u>Common Name</u>
	'Wakeiebisu'
	'White Gumbo'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia sp.	Black-eyed Susan; Goldstrum
Santolina, virens	Santolina
Saxifraga sp.	Saxifrage; Purple Dome
Scabiosa sp.	Pincushion Flower
Sedum album	Stonecrop
Sedum cauticola	Stonecrop; Lidakense
Sedum dasyphyllum	Stonecrop
Sedum spurium	Stonecrop; Dragon's Blood
Spiraea Bumalda	Spirea; Anthony Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium sp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica sp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowwood Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca minor	Common Periwinkle, Vinca
Vitis sp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filamentosa	Adam's Needle, Yucca

* Do not use on food producing trees, vines or plants.

** Not for use on container grown plants.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. If these terms are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid.

NET WEIGHT XX POUNDS
Knox Fertilizer Company, Inc.
P.O. Box 248
Knox, IN 46534

ACCEPTED
with COMMENTS
In EPA Letter Dated

JAN 21 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8378-39