

8378-41

12/23/2002

1/6

DEC 23 2002

H. R. McLane
 H.R. McLane Inc.
 7210 Red Road, Suite 206A
 Miami, FL 33143-5321

Dear Mr. McLane:

SUBJECT: Label Revision Amendment - First Aid Statement
 Shaw's Annual Grass Control 147
 EPA Registration No. 8378-41
 Application Dated September 23, 2002

The labeling, referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided the following change is made:

Under "FIRST AID" section, remove the last sentence "Have this product container . . ." and place it as an independent entry above the "IF SWALLOWED" paragraph.

The amended label supersedes all previously accepted ones. A stamped copy is enclosed for your records. Please submit one copy of your printed label before you release the product for shipment.

Sincerely yours,

Joanne I. Miller
 Product Manager (23)
 Herbicide Branch
 Registration Division (7505C)

Enclosure

CONCURRENCES							
SYMBOL ▶	7505C						
SURNAME ▶	JIMiller						
DATE ▶	Dec 23, 2002						

Shaw's Annual Grass Control 147

Contains TREFLAN® The Flower and Garden Weed Preventer

1 pound covers up to 1250 square feet

Prevents Crabgrass and Most Other Annual Weed Grasses in Flower Beds, Home Vegetable Gardens and Around Ornamental Trees, Shrubs and Roses.

ACTIVE INGREDIENT:

Trifluralin; Alpha, alpha, alpha, trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 1.47%

INERT INGREDIENTS: 98.53%

TOTAL 100.00%

EPA Reg. No. 8378-41

EPA Est. No. 8378-IN-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED: Call a physician or poison control center immediately. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

See additional Precautionary Statements on this label

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or contact with skin or clothing. This product may cause skin sensitization reactions in some people.

Personal Protective Equipment

Users should wear protective clothing such as long pants, long-sleeved shirt, shoes and socks and household gloves

ACCEPTED
with COMMENTS
In EPA Letter Dated

DEC 23 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

008378-41

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment wash waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

Entry Restrictions

Do not allow people or pets to enter the treated area until dusts have settled. If watering in is required after the application do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering is complete and the surface is dry. Do not apply this product in a way that will contact any person or pet either, directly or through drift. Keep people and pets out of the area during application.

Users Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Do not reuse bag. Place bag in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

This product prevents the listed broadleaf and grassy weeds from becoming established around flowers, roses, shrubs, trees and vegetables. It eliminates the need for extensive hand weeding. Without weeds, plants, can grow larger, stronger and healthier. Do not apply to Ajuga, Myrtle, Pachysandra as injury may result.

WHEN TO APPLY

Apply as early as possible in the growing season before weed seeds germinate to take advantage of season-long weed control. Apply to soil in plant beds anytime during the growing season around listed plants and anytime prior to setting transplants (see section under Vegetables For use Before transplanting "sets") Do not use on Flower Seeds. Follow these directions carefully.

WEEDS CONTROLLED

Annual Bluegrass, Barnyardgrass (watergrass), Brachiaria (Signal Grass), Bromegrass, Cheat, Crabgrass, Foxtail, Goosegrass, Johnsongrass (from seed), Junglerice, Panicum (fall), Panicum (Texas), Sandbur, Sprangletop, Stinkgrass, Woolly Cupgrass. Also control these broadleaf weeds: Carelessweed, Carpetweed, Chickweed, Florida Pusley, Goosefoot, Knotweed, Kochia, Lambsquarter, Pigweed, Puncture Vine, Purslane, Russian Thistle, Stinging Nettle.

HOW TO USE FOR ORNAMENTAL PLANTS

Roses, Shrubs, Flowers and Trees

For plants such as roses, shrubs, and flowers and trees remove existing weeds before applying this product. Till the soil to prepare a smooth soil planting surface. For plants which are 2 to 3 ft. in height sprinkle.

FOR ESTABLISHED PLANTS

1 LB. EVENLY OVER 320 SQ. FT. OF SURROUNDING SOIL SURFACE. (1 OZ./20 SQ. FT.)

Do not apply to plant foliage. Best results will be obtained by lightly sprinkling the granules into the soil surface after application. For optimum weed control plan on a second application in 4 weeks.

PRE-PLANTING APPLICATION

For "light" soils apply 1 lb./1250 sq. ft.

On "medium loam" soil apply 1 lb. to 960 sq. ft.

For "heavy" soils apply 1 lb./640 sq. ft.

Mix into soil surface. Not recommended for "muck" soils. Small gladiola corms less than 1 inch in diameter may be injured by pre-plant applications. Use only on 2 to 3 feet high listed transplants. Do not use on flower seeds. Best results will be obtained when lightly watered in. Plan on a second application in 4 weeks.

ORNAMENTAL PLANTS

Use on the following lists of plants out of doors. Do not use in enclosed areas such as greenhouses.

ROSES AND OTHER ESTABLISHED FLOWERS

Achillea, Ageratum, Arctotis, Aster, Balsam, Calendula, California Poppy, Forget-Me-Not, Four O'Clock, Gaillardia, Gladiolus, Ixora, Lobelia, Lupinus, Marigold, Morning Glory, Nasturtium, Nicotiana, Perwinkle, Petunia, Phlox, Portulaca, Rudbeckia, Salvia, Scabiosa, Shasta Daisy, Snapdragon, Stocks, Snow on the Mountain, Sunflower, Sweet Alyssum, Sweet Pea, Sweet William, Zinnia.

ESTABLISHED SHRUBS

Arborvitae, Azalea, Barberry, Boxwood, Camellia, Cherry Laurel, Cinquefoil, Common Lilac, Cotoneaster, Deutzia, Elaeagnus, Euonymus, Forsythia, Holly, Honeysuckle, India Hawthorn, Juniper, Mock Orange, Mountain Laurel, Pieris Japonica, Pineapple guava, Privet, Rhododendron, Rubira, Sakaki, Spirea, Viburnum, Weiglea Bristol Ruby, Willow, Yew.

ESTABLISHED ORNAMENTAL TREES

Apple, Bald Cypress, Balsam Fir, Black Gum, Black Locust, Canada Hemlock, Chinese Chestnut, Cottonwood, Dogwood, Douglas Fir, European White Birch, Honey Locust, Japanese Larch, London

Plane Trees, Maple, Oak, Pine, Redbud, Spruce, Stone Fruits, Sycamore, Sweetgum, Tulip Tree, White Ash.

ESTABLISHED ORNAMENTAL GROUND COVERS

Algerian Ivy, Asparagus, Bellflower, Big Blue Lilyturf, Bunge, California Ivy, Ceanothus, Cotoneaster, Crown Vetch, Dwarf Coyote Brush, Dwarf Peter Pan, Gazania, Germander, Ice Plant, Marigold, Mascarene grass, Myoporum, Needlepoint Ivy, Rock Rose, Rosemary, Rose of Sharon, Rupture Wort, St. John's Wort, Sedum, Snow-in Summer, Speedwell, Strawberry, Thrift-Sea Pink, Tickseed, Trailing African Daisy, Verbena, Wire Plant, Woolly Yarrow.

HOW TO USE FOR VEGETABLES FOR USE IN HOME GARDENS ONLY.

Apply and mix into the soil before planting to provide long lasting control of many annual grasses and broadleaf weeds as listed.

For "light" soils apply 1 lb./1250 sq. ft.

On "medium loam" soil apply 1 lb. to 960 sq. ft.

For "heavy" soils apply 1 lb./640 sq. ft.

Controls weeds as they germinate. Will not control established weeds. Application may be made up to 10 weeks prior to planting direct seeded vegetables and anytime prior to setting transplants. Remove all existing weeds first then lightly till the soil to prepare a smooth seed bed.

For vegetables from seed: Treat soil prior to planting seed - Use for Carrots, Celery, Okra, Southern Peas, Green Peas, Lima Beans, Snap Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Turnip Greens, Mustard Greens, Collards, and Kale. May be applied up to 10 weeks before planting direct seeded vegetables.

May be used on Cantaloupes, Cucumbers, Watermelons after plants are up in the 3 to 4 leaf stage. Asparagus (established) - Apply in winter or early spring after ferns are removed, but prior to spear emergence.

For use before transplanting "sets" - Celery, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Peppers, Tomatoes. Make the application to the prepared soil bed and mix into the top 1 to 3 inches of soil before "sets" are placed.

For potatoes: May be applied to soil surface after planting "seed pieces" before or after emergence. Use care not to damage seed pieces while cultivation nor allow treated soil to contact emerged plant foliage. In all cases mix and cultivate granules into the soil after application, using care not to damage emerged crops.

DO NOT apply this product to "muck" soils. Do not over-dose! Injury may occur to subsequent crops which are sensitive to Trifluralin. Do not use with crops other than those listed on this label. This product is not intended for use on lawns. Refer to other EPA registered products for that use. Do not use in enclosed areas such as greenhouses.

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product.

NOTICE: Knox Fertilizer Company, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use, subject to the inherent risks referred to above. Knox Fertilizer Company, Inc. makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Knox Fertilizer Company, Inc. be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

6/6

20020918.amendmentPR2001-01&06

Distributed By:
Knox Fertilizer Company, Inc.
PO Box 248
Knox, Indiana 46534

Net Weight 10 lbs. (4.5 kg.)

Shaw's is a registered trademark of Knox Fertilizer Company, Inc.
Treflan is a registered trademark of Dow AgroSciences LLC