

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JUN 12 2003

H.R. McLane
H.R. McLane, Inc.
7210 Red Road, Suite 206
Miami, FL 331434-5321

Dear Mr. McLane:

Subject: Label Amendment
Shaw's X-X-X Fertilizer Plus Preemergent Weed Control (High end range)
EPA Reg. No. 8378-45
Submission dated September 18, 2002

The labeling, referred to above, submitted in connection with the application under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the following changes:

1. Under "If Inhaled in First Aid";

Move the last sentence "Have this product . . . , or when going for treatment" to A new paragraph. Also, add the following sentence:
"You may contact 1-800-xxx-xxxx for emergency medical treatment information."

2. Under "Precautionary Statements";

It is recommended that you add the following statements:

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

3. Under "Direction for Use";

- a. It is recommended that you require persons applying this product to wear long-sleeved shirt and long pants, and shoes plus socks.
- b. Add the following statements:
"Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application."

2/5

Page 2
EPA Registration No. 8378-45


4. Und"Storage and Disposal";

Change "Storage" to "Pesticide Storage," and "Disposal" to "Pesticide Disposal."

5. The Agency has not reviewed or accepted the warranty and liability statements. It may need to be changed at a later date.
6. The Agency no longer accepts multi range active ingredient (AI) registrations. You must submit another application for the "low end range" (0.50 % AI).

The amended label supersedes all previously accepted labels. A stamped copy of the revised label is enclosed for your records. Please submit one copy of your final printed label before you release the product for shipment.

Sincerely yours,



Jim Tompkins
Product manager (25)
Herbicide Branch
Registration Division (7505C)

Enclosures

CONCURRENCES							
SYMBOL ▶	7505C						
SURNAME ▶	C. Bien						
DATE ▶	6-10-2003						

EPA Form 1320-1 (12-70) OFFICIAL FILE COPY

High end range

Shaw's X-X-X Fertilizer Plus Preemergent Weed Control

ACCEPTED
with COMMENTS
in EPA Letter Dated

Under the Federal Insecticide
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
and Spurge. Supplies quick

Preemergent Control of Crabgrass, Chickweed, Foxtail, Goosegrass, Knotweed, Oxalis, Poa Annua,
greening and extended feeding.

~~8378-45~~

ACTIVE INGREDIENT:

Pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine)..... 1.79%

INERT INGREDIENTS: 98.21%
100.00%

JUN 12 2003

EPA REG. NO. 8378-45

EPA EST. NO. 8378-IN-1

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control, or doctor, or when going for treatment.

See additional Precautionary Statements on this label.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not feed clippings to livestock. Do not graze treated areas.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, sweep up and use per label instructions or dispose of as directed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

STORAGE: Store product in original bag in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all the contents of the bag be used, carefully following label directions and precaution. **IF EMPTY:** Do not reuse bag. Place bag in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

This product is a turf fertilizer and a selective preemergence herbicide for use on established turfgrass including Kentucky Bluegrass, Perennial Ryegrass, Fine Fescue, Tall Fescue, Bentgrass and Poa Annua (except putting greens and tees), Bermudagrass (non "winter-overseeded"), St. Augustinegrass, Bahiagrass, Zoysiagrass, Centipedegrass and blends of these grasses. Refer to label section entitled "Bermudagrass, Bentgrass and Poa Annua Precautions". Do not apply this product through any type of irrigation system. Do not apply aerially.

WEEDS CONTROLLED

When used at the recommended rates (See Turf Species and Rate Table) this product will prevent sprouting of the following grassy and dicot weeds and provides a nitrogen feeding.

Grassy Weeds

Poa Annua
Barnyardgrass
Crabgrass-(Large, Smooth)
Foxtail-(Yellow, Green)
Goosegrass
Fall panicum

Dicot Weeds

Common Chickweed
Knotweed
Oxalis
Prostrate Spurge

To prevent Goosegrass, Crabgrass, Foxtail, Barnyardgrass, Fall panicum, Oxalis, Knotweed, Prostrate Spurge, apply in the spring before they sprout. To prevent Poa Annua and fall germinating chickweed, apply product to dry foliage in late summer or early fall before seeds germinate. Repeat application yearly for continued control.

ESTABLISHED TURF

When applied prior to germination of weeds this product will control weeds listed in WEEDS CONTROLLED section, in established Northern and Southern turf.

NORTHERN Cool Season GRASSES

Apply 1.3 to 2.2 lbs. per 1,000-sq. ft. prior to germination of weeds. (equiv. to 1.0 to 1.75 lbs. of Active Ingredient/acre)

SOUTHERN Warm Season GRASSES

Apply 1.3 to 2.6 lbs. per 1,000-sq. ft. prior to germination of weeds. (equiv. To 1.0 to 2.0 lbs. of Active Ingredient/acre)

Where light weed infestations are expected, the lower rates may be used. For extended weed control, repeat application is recommended. For best results, water after all application (1/4 to 1.2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical herbicide barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying the turf after applying this product.

PRECAUTIONS IN USING

Do not use on Bentgrass and Poa Annua putting greens and tees, or on dichondra. Where Bentgrass and Bentgrass-containing turf and Poa Annua are desired species, a lower rate application only is recommended. On Cool Season Turfgrass avoid application when temperatures are regularly in the 80's. This product may cause discoloration and thinning where over applications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperatures-below 40 degrees F, scalping, and excessive soil moisture).

BERMUDAGRASS, BENTGRASS AND POA ANNUA PRECAUTIONS

Do not apply to dormant Bermudagrass, Bentgrass or Poa Annua (turf that is not growing actively or has frost injury). May be applied to dormant Bermudagrass if it has been maintained at $\frac{1}{2}$ inch or more high cut and is growing in medium fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury of Bermudagrass do not make an application until turf has greened up, is growing vigorously, and has filled in the dead areas. Application to Bermudagrass overseeded with cool season grass may injure and thin the overseeded variety. Do not apply to Bentgrass or Poa Annua after late summer.

OTHER PRECAUTIONS

Delay seeding 4 months and sprigging by 5 months after an application. For best results, the soil surface should be worked (verticut and/or etc.) to destroy any remaining chemical residue before planting. For newly planted areas, wait until the turf has been mowed up to 4 times, and is growing vigorously before application. Do not apply to the same area more than twice per year. Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5 to 10 miles per hour) due to its yellow staining potential.

APPLICATION EQUIPMENT AND INSTRUCTIONS SPREADER SETTINGS

Your used spreader may need calibrating. Since proper application is based on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreading equipment may require adjustments from the following figures. If you have reason to believe that your spreader is not calibrated correctly, test apply by placing the amount for 500 sq. ft. into your spreader and apply to your pre-measured 500 sq. ft. (5 ft. wide by 100 ft. long). The spreader settings are approximate. Different product analysis or particle size will affect bulk density and actual spreader setting needs.

Spreader Setting	Spread width	Pounds per 1,000 sq. ft.				
		1.3	1.6	1.9	2.2	2.6
Scotts R8A	7'	xx	xx	xx	xx	xx
Lesco high wheel	7'	xx	xx	xx	xx	xx
Spyker 24-22	7'	xx	xx	xx	xx	xx
Prizelawn CBR-11	7'	xx	xx	xx	xx	xx
Earthway 2400-SS	7'	xx	xx	xx	xx	xx
Vicon	5'	xx	xx	xx	xx	xx
Lely	4 1/2'	xx	xx	xx	xx	xx

IMPORTANT: PLEASE READ BEFORE USE

By using this product the user accepts the following: **LIMITED WARRANTY:** Knox Fertilizer Company warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to herein, when used in accordance with its directions; and (c) that the directions, warnings cautions, and other statements on this label are based upon responsible experts evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, of residues on food crops, and upon reports of field experience. Testing has not been performed on all varieties of Bentgrass and Poa Annua, in all states, or under all application, weather and crop conditions. The user should exercise judgement and caution when using this product on a given variety until familiar with the performance under his growing conditions.

There are no express warranties other than those set forth herein. Knox neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. Knox expressly excluded and disclaims all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality or performance.

This warranty does not extend to, and the user shall be solely responsible for, any loss or damage which results from the use of this product in any manner which is inconsistent with this label's directions, warning or cautions.

User's exclusive remedy and Knox's or seller's exclusive liability for any claim, loss, damage, or injury resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort, or otherwise, shall be limited, at Knox's option, to replacement of, or repayment of the purchase price for, the quantity of product with respect to which damages are claimed. In no event shall Knox or Seller be liable for special, indirect, or consequential damages resulting from the use or handling of this product.

Guaranteed Analysis	
X - X - X	
Total Nitrogen (N)	XX%
Available Phosphorus (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Primary Plant Foods Derived from: Urea, Muriate of Potash, and Ammoniated Phosphate.	

Distributed by: Knox Fertilizer Company, Inc.
PO Box 248
Knox, Indiana 46534

NET WEIGHT 50 LBS. (22.7 kg.)